

# SMOOTHING PROPERTIES IN MULTISTEP BACKWARD DIFFERENCE METHOD AND TIME DERIVATIVE APPROXIMATION FOR LINEAR PARABOLIC EQUATIONS

YUBIN YAN

Received 11 May 2004 and in revised form 6 December 2004

A smoothing property in multistep backward difference method for a linear parabolic problem in Hilbert space has been proved, where the operator is selfadjoint, positive definite with compact inverse. By using the solutions computed by a multistep backward difference method for the parabolic problem, we introduce an approximation scheme for time derivative. The nonsmooth data error estimate for the approximation of time derivative has been obtained.

## 1. Introduction

Consider the nonhomogeneous linear parabolic equation

$$u_t + Au = f, \quad \text{for } t > 0, \text{ with } u(0) = v, \quad (1.1)$$

in a Hilbert space  $H$  with norm  $\|\cdot\|$ , inner product  $(\cdot, \cdot)$ , where  $u_t = du/dt$  and  $A$  is a linear, selfadjoint, positive definite, not necessarily bounded operator with a compact inverse, densely defined in  $\mathcal{D}(A) \subset H$ , where  $v \in H$  and  $f$  is a function of  $t$  with values in  $H$ .

Since  $A^{-1}$  is compact, we assume that  $A$  has eigenvalues  $\{\lambda_j\}_{j=1}^{\infty}$  and a corresponding basis of orthonormal eigenfunctions  $\{\varphi_j\}_{j=1}^{\infty}$ . For any arbitrary function  $g(\lambda)$ , defined on the spectrum  $\sigma(A) = \{\lambda_j\}_{j=1}^{\infty}$  of  $A$ , the operator norm of  $g(A)$  can be computed by

$$\|g(A)\| = \sup_j |g(\lambda_j)| = \sup_{\lambda \in \sigma(A)} |g(\lambda)|, \quad (1.2)$$

which will be used frequently in this paper.

Let  $U^n$ ,  $n \geq 0$ , be an approximation of the solution  $u(t_n)$  of (1.1) at time  $t_n = nk$ , where  $k$  is the time step. We introduce the backward difference operator  $\bar{\partial}_p$ ,  $p \geq 1$ , by

$$\bar{\partial}_p U^n = \sum_{j=1}^p \frac{k^{j-1}}{j} \bar{\partial}^j U^n, \quad \text{where } \bar{\partial} U^n = \frac{U^n - U^{n-1}}{k}. \quad (1.3)$$

It is easy to see that, for any smooth real-valued function  $u$ ,

$$u_t(t_n) = \bar{\partial}_p u^n + O(k^p), \quad \text{as } k \rightarrow 0, \text{ with } u^n = u(t_n). \quad (1.4)$$

With  $U^0, \dots, U^{p-1}$  given, we define our approximate solution  $U^n$  by

$$\bar{\partial}_p U^n + AU^n = f^n, \quad \text{for } n \geq p, \text{ where } f^n = f(t_n). \quad (1.5)$$

It is well known from the theory for numerical solution of ordinary differential equations, see, for example, Hairer and Wanner [4], that this method is  $A(\theta)$ -stable for some  $\theta = \theta_p > 0$  when  $p \leq 6$ . The theory of stability and error estimates for the approximation of the solution of (1.1) by a multistep method in both constant and variable time-step cases have been well developed, see Becker [1], Bramble et al. [2], Crouzeix [3], Hansbo [6], LeRoux [7, 8], Palencia and Garcia-Archilla [9], Savaré [10], and Thomée [11], and the references therein.

The purpose of this paper is to consider the smoothing property in multistep backward difference method and time derivative approximation of (1.1). The similar results in single-step methods for homogeneous parabolic problems in general Banach space have been studied, for example, by Hansbo [5, 6] and Yan [12, 13].

We obtain, in Theorem 2.1, the following smoothing property in multistep backward difference method: if  $U^n$  is the solution of (1.5) with  $f = 0$ , then we have, with  $p \leq 6$ ,

$$\|\bar{\partial}_p U^n\| \leq Ct_n^{-1} \sum_{j=0}^{p-1} \|U^j\|, \quad \text{for } n \geq 2p, \quad U^0, U^1, \dots, U^{p-1} \in H. \quad (1.6)$$

We introduce the norm  $|\nu|_s = (A^s \nu, \nu)^{1/2}$ ,  $s \in \mathbb{R}$ , defined by

$$|\nu|_s^2 = \sum_{j=1}^{\infty} \lambda_j^s (\nu, \varphi_j)^2 < \infty, \quad \text{for } s \in \mathbb{R}, \quad (1.7)$$

where  $\{\lambda_j, \varphi_j\}_{j=1}^{\infty}$  is the eigensystem of the operator  $A$ . We see that  $|\cdot|_0 = \|\cdot\|$ .

It is natural to approximate the time derivative  $u_t(t_n)$  of the solution of (1.1) by  $\bar{\partial}_p U^n$  ( $n \geq 2p$ ), where  $U^n$ ,  $n \geq p$ , is computed by the multistep backward difference method (1.5). Approximating  $u_t(t_n)$  by  $\bar{\partial}_p U^n$ , we obtain, in Theorem 3.3, with  $n \geq 2p$ ,

$$t_n^{2p+2} \|\bar{\partial}_p U^n - u_t(t_n)\|^2 \leq C \sum_{j=p}^{2p-1} \left( \|U^j - u^j\|_{-2p}^2 + k^{2p+2} \|A(U^j - u^j)\|^2 \right) + Ck^{2p} G(u), \quad (1.8)$$

where

$$G(u) = \int_0^{t_n} \left( \|u^{(p+1)}(s)\|_{-2p-1}^2 + s^{2p+2} \|u^{(p+1)}(s)\|_1^2 + s^2 \|u_t(s)\|_1^2 \right) ds + t_{2p}^3 \|u_t(t_{2p})\|_1^2. \quad (1.9)$$

In the case of  $f \equiv 0$ , if the discrete initial values satisfy, with  $U^0 = \nu$ ,

$$\|U^j - u^j\|_{-2p} + k^{p+1} \|A(U^j - u^j)\| \leq Ck^p \|\nu\|, \quad \text{for } p \leq j \leq 2p-1, \quad \nu \in H, \quad (1.10)$$

for some suitable discrete starting values  $U^0, U^1, \dots, U^{p-1}$ , then, in Corollary 3.4, we get

$$\|\bar{\partial}_p U^n - u_t(t_n)\| \leq C k^p t_n^{-p-1} \|v\|, \quad \text{for } n \geq 2p. \quad (1.11)$$

We also discuss the starting value approximation in the case of  $p = 2$ . For suitable initial approximation  $U^0, U^1$ , we can prove, with  $U^0 = v \in H$ ,

$$\sum_{j=2}^3 (|U^j - u^j|_{-4} + k^3 \|A(U^j - u^j)\|) \leq C k^2 \left( \|v\| + \|f(0)\| + \int_0^{t_j} \|f'(\tau)\| d\tau \right). \quad (1.12)$$

Thus, in the case of  $p = 2$ , our error estimate reads, with  $U^0 = v \in H$ ,

$$\|\bar{\partial}_2 U^n - u_t(t_n)\| \leq C k^2 t_n^{-3} \left( \|v\| + \|f(0)\| + \int_0^{t_3} \|f'(\tau)\| d\tau + K(u) \right), \quad n \geq 4, \quad (1.13)$$

where

$$K(u)^2 = \int_0^{t_n} \left( |u^{(3)}(s)|_{-5}^2 + s^6 |u^{(3)}(s)|_1^2 + s^2 |u_t(s)|_1^2 \right) ds + t_4^3 |u_t(t_4)|_1^2. \quad (1.14)$$

By  $C$  and  $c$  we denote large and small positive constants independent of the functions and parameters concerned, but not necessarily the same at different occurrences. When necessary for clarity, we distinguish constants by subscripts.

## 2. Smoothing properties

In this section, we will show the smoothing properties for the multistep backward difference method. Before showing this, we first discuss some properties of the backward difference operator  $\bar{\partial}_p$  defined by (1.3). We first note that (1.3) can be written in another form as

$$\bar{\partial}_p U^n = k^{-1} \sum_{\nu=0}^p c_\nu U^{n-\nu}, \quad (2.1)$$

where the coefficients  $c_\nu$  are independent of  $k$ . Introducing  $P(x) = \sum_{\nu=0}^p c_\nu x^\nu$ , it is easy to check that (1.4) is equivalent to

$$P(e^{-\lambda}) - \lambda = O(\lambda^{p+1}), \quad \text{as } \lambda \rightarrow 0. \quad (2.2)$$

In fact, with  $u(t) = e^t$  in (1.4), we have

$$P(e^{-k}) - k = O(k^{p+1}), \quad \text{as } k \rightarrow 0, \quad (2.3)$$

replacing  $k$  by  $\lambda$ , we show (2.2). On the other hand, if (2.2) holds, (1.4) follows from Taylor expansion of  $\bar{\partial}_p u^n$  at  $t_n$ .

For  $p = 1$ , (1.5) reduces to the backward Euler method

$$\frac{U^n - U^{n-1}}{k} + AU^n = f^n, \quad \text{for } n \geq 1, \quad (2.4)$$

and the starting value is  $U^0 = v$ .

For  $p = 2$ , we have

$$\frac{(3/2)U^n - 2U^{n-1} + (1/2)U^{n-2}}{k} + AU^n = f^n, \quad \text{for } n \geq 2, \quad (2.5)$$

and both  $U^0$  and  $U^1$  are needed to start the procedure.

Bramble et al. [2] obtain the following stability result, that is, with  $U^n$  the solution of (1.5),

$$\|U^n\| \leq C \sum_{j=0}^{p-1} \|U^j\| + Ck \sum_{j=p}^n \|f^j\|, \quad \text{for } n \geq p. \quad (2.6)$$

In this paper, we first show the smoothing property for the multistep backward difference method.

**THEOREM 2.1.** *Let  $p \leq 6$ . Then there is a constant  $C$ , independent of the positive definite operator  $A$ , such that for the solution  $U^n$  of (1.5) with  $f = 0$ ,*

$$\|\bar{\partial}_p U^n\| \leq Ct_n^{-1} \sum_{j=0}^{p-1} \|U^j\|, \quad \text{for } n \geq 2p. \quad (2.7)$$

To prove this theorem, we need the following lemma from Thomée [11, Lemma 10.3].

**LEMMA 2.2.** *The solution of (1.5) may be written, with  $f = 0$ , as*

$$U^n = \sum_{s=0}^{p-1} \beta_{ns}(kA) U^s, \quad \text{for } n \geq p, \quad (2.8)$$

where with  $\lambda > 0$ ,  $P(\zeta) = \sum_{\nu=0}^p c_\nu \zeta^\nu$ , the  $\beta_{ns}(\lambda)$  are defined by,

$$\beta_{ns}(\lambda) = \sum_{j=p-s}^p \beta_{n-s-j}(\lambda) c_j, \quad \sum_{j=0}^{\infty} \beta_j(\lambda) \zeta^j := (P(\zeta) + \lambda)^{-1}, \quad (2.9)$$

where it is assumed that  $\beta_{n-s-j}(\lambda) = 0$  in the case  $n - s - j < 0$ .

If  $p \leq 6$ , there are positive constants  $c$ ,  $C$ , and  $\lambda_0$  such that

$$|\beta_j(\lambda)| \leq \begin{cases} Ce^{-cj\lambda}, & \text{for } 0 < \lambda \leq \lambda_0, \\ C\lambda^{-1}e^{-cj}, & \text{for } \lambda \geq \lambda_0. \end{cases} \quad (2.10)$$

*Proof of Theorem 2.1.* By (2.8) and (2.1), we find that

$$\bar{\partial}_p U^n = k^{-1} \sum_{\nu=0}^p c_\nu \sum_{s=0}^{p-1} \beta_{(n-\nu)s}(kA) U^s = \sum_{s=0}^{p-1} \left[ k^{-1} \sum_{\nu=0}^p c_\nu \beta_{(n-\nu)s}(kA) \right] U^s. \quad (2.11)$$

By (2.8), we see that  $n - \nu \geq p$  for any  $0 \leq \nu \leq p$ , which implies that  $n$  must be larger than or equal to  $2p$ , that is,  $n \geq 2p$ . Since  $\bar{\partial}_p U^n$  is linearly dependent on  $U^s$  ( $0 \leq s \leq p-1$ ), it suffices to consider separately the cases when  $U^l \neq 0$ ,  $0 \leq l \leq p-1$ , and  $U^s = 0$ ,  $0 \leq s \leq p-1$ ,  $s \neq l$ . In other words, we need to show, for  $U^l \neq 0$ ,  $0 \leq l \leq p-1$ , and  $U^s = 0$ ,  $0 \leq s \leq p-1$ ,  $s \neq l$ ,

$$\|\bar{\partial}_p U^n\| \leq C t_n^{-1} \|U^l\|. \quad (2.12)$$

We first consider the case  $0 < l \leq p-1$ . By Lemma 2.2, we have

$$\begin{aligned} \bar{\partial}_p U^n &= k^{-1} \sum_{\nu=0}^p c_\nu (\beta_{(n-\nu)l} U^l) \\ &= k^{-1} \sum_{\nu=0}^p c_\nu \left( \sum_{j=p-l}^p \beta_{n-\nu-l-j}(kA) c_j \right) U^l \\ &= k^{-1} \sum_{j=p-l}^p \left( \sum_{\nu=0}^p c_\nu \beta_{n-\nu-l-j}(kA) \right) c_j U^l, \quad \text{for } 0 < l \leq p-1. \end{aligned} \quad (2.13)$$

We remark that  $n - \nu - l - j$  may be negative. In this case, we assume that  $\beta_{n-\nu-l-j}(\lambda) = 0$ . By Lemma 2.3, we have

$$\sum_{\nu=0}^p c_\nu \beta_{n-\nu-l-j}(\lambda) = -\lambda \beta_{n-l-j}(\lambda). \quad (2.14)$$

Thus

$$\bar{\partial}_p U^n = k^{-1} \sum_{j=p-l}^p (-kA \beta_{n-l-j}(kA)) c_j U^l, \quad (2.15)$$

and (2.12) will follow from

$$\left\| k^{-1} \sum_{j=p-l}^p (-kA \beta_{n-l-j}(kA)) c_j \right\| \leq C t_n^{-1}. \quad (2.16)$$

In view of (1.2), (2.16) may be written as, for  $0 < l \leq p-1$ ,

$$\left| n \lambda \sum_{j=p-l}^p \beta_{n-l-j}(\lambda) \right| \leq C, \quad \text{for } \lambda \in \sigma(kA). \quad (2.17)$$

Now we show (2.17). By (2.10). We have, for small  $\lambda$ ,  $0 \leq \lambda \leq \lambda_0$ ,

$$\left| n\lambda \sum_{j=p-l}^p \beta_{n-l-j}(\lambda) \right| \leq C \sum_{j=p-l}^p n\lambda e^{-c(n-l-j)\lambda} \leq C, \quad (2.18)$$

and, for large  $\lambda$ ,  $\lambda > \lambda_0$ ,

$$\left| n\lambda \sum_{j=p-l}^p \beta_{n-l-j}(\lambda) \right| \leq C \sum_{j=p-l}^p ne^{-c(n-l-j)} \leq C, \quad \text{for } \lambda \geq \lambda_0. \quad (2.19)$$

Hence (2.17) holds.

It remains to consider the case  $l = 0$ . We have, by Lemma 2.2,

$$\bar{\partial}_p U^n = k^{-1} \sum_{\nu=0}^p c_\nu (\beta_{(n-\nu)0} U^0) = k^{-1} \left( \sum_{\nu=0}^p c_\nu \beta_{n-\nu-p}(kA) \right) c_p U^0. \quad (2.20)$$

Thus in this case, (2.12) will follow from

$$\left\| k^{-1} \sum_{\nu=0}^p (c_\nu \beta_{n-\nu-p}(kA)) c_p \right\| \leq C t_n^{-1}. \quad (2.21)$$

By (1.2), (2.21) may be written as

$$n \left| \sum_{\nu=0}^p c_\nu \beta_{n-\nu-p}(\lambda) \right| \leq C, \quad \text{for } \lambda \in \sigma(kA), n \geq 2p, \quad (2.22)$$

or, using (2.14),

$$|n\lambda \beta_{n-p}(\lambda)| \leq C, \quad \text{for } \lambda \in \sigma(kA), n \geq 2p, \quad (2.23)$$

which we will now prove. For small  $0 \leq \lambda < \lambda_0$ , we have, by (2.10),

$$|n\lambda \beta_{n-p}(\lambda)| \leq (n\lambda e^{-cn\lambda}) e^{cp\lambda} \leq C. \quad (2.24)$$

For  $\lambda > \lambda_0$ , using again (2.10), we have,

$$|n\lambda \beta_{n-p}(\lambda)| \leq C(ne^{-cn}) e^{cp} \leq C. \quad (2.25)$$

Thus (2.23) holds. Together these estimates complete the proof of Theorem 2.1.  $\square$

LEMMA 2.3. *Let  $p \leq 6$ . Let  $\beta_m(\lambda)$ ,  $m \geq 0$ , and  $c_m$ ,  $0 \leq m \leq p$ , be defined as in Lemma 2.2. Then, for any  $n \geq 2p$ ,  $0 < l \leq p-1$ ,  $p-l \leq j \leq p$ ,*

$$(c_0 + \lambda) \beta_{n-l-j}(\lambda) + c_1 \beta_{n-l-j-1}(\lambda) + \cdots + c_p \beta_{n-l-j-p}(\lambda) = 0, \quad (2.26)$$

where it is assumed that  $\beta_{n-l-j-\nu}(\lambda) = 0$  in the case of  $n-l-j-\nu < 0$  for  $0 \leq \nu \leq p$ .

*Proof.* By (2.9), we have

$$(P(\zeta) + \lambda) \left( \sum_{j=0}^{\infty} \beta_j(\lambda) \zeta^j \right) = 1, \quad (2.27)$$

that is,

$$(c_0 + \lambda + c_1 \zeta^1 + c_2 \zeta^2 + \dots) (\beta_0(\lambda) + \beta_1(\lambda) \zeta + \beta_2(\lambda) \zeta^2 + \dots) = 1. \quad (2.28)$$

Comparing with the coefficients of  $\zeta^j$ ,  $j = 0, 1, 2, \dots$ , we get

$$\begin{aligned} (c_0 + \lambda) \beta_0(\lambda) &= 1, \\ (c_0 + \lambda) \beta_1(\lambda) + c_1 \beta_0(\lambda) &= 0, \\ &\vdots \\ (c_0 + \lambda) \beta_p(\lambda) + c_1 \beta_{p-1}(\lambda) + \dots + c_p \beta_0(\lambda) &= 0, \\ (c_0 + \lambda) \beta_{p+1}(\lambda) + c_1 \beta_p(\lambda) + \dots + c_p \beta_1(\lambda) &= 0, \\ &\vdots \end{aligned} \quad (2.29)$$

which implies that, for any  $m \geq 1$  (not  $m = 0$ ),

$$(c_0 + \lambda) \beta_m(\lambda) + c_1 \beta_{m-1}(\lambda) + \dots + c_p \beta_{m-p}(\lambda) = 0, \quad (2.30)$$

where  $\beta_{m-\nu} = 0$  if  $m - \nu < 0$  for  $1 \leq \nu \leq p$ .

Note that  $n - l - j \geq 1$  since  $n \geq 2p$ ,  $0 < l \leq p - 1$ ,  $p - l \leq j \leq p$ . Thus we obtain (2.26) by replacing  $m$  with  $n - l - j$  in (2.30). The proof is complete.  $\square$

### 3. Error estimates

In this section, we will show the error estimates for the approximation  $\bar{\partial}_p U^n$  of the time derivative  $u_t(t_n)$  in nonsmooth data cases.

We first recall the following stability result, see Thomée [11, Theorem 10.4].

LEMMA 3.1. *Let  $p \leq 6$  and  $s \geq 0$ , and let  $U^n$  be the solution of (1.5). Then, with  $C$  independent of the positive definite operator  $A$ ,*

$$\begin{aligned} t_n^s \|U^n\|^2 + k \sum_{j=p}^n t_j^s \|U^j\|_1^2 &\leq C \sum_{j=0}^{p-1} \left( \|U^j\|_{-s}^2 + k^s \|U^j\|^2 \right) \\ &\quad + Ck \sum_{j=p}^n \left( \|f^j\|_{-s-1}^2 + t_j^s \|f^j\|_{-1}^2 \right), \quad \text{for } n \geq p. \end{aligned} \quad (3.1)$$

By shifting the origin, we have the following generalization of Lemma 3.1.

LEMMA 3.2. *Let  $p \leq 6$  and  $s \geq 0$ , and let  $U^n$  be the solution of (1.5). Assume that  $m \geq p$  and  $U^{m-p}, \dots, U^{m-1}$  are given. Then, with  $C$  independent of the positive definite operator  $A$ ,*

$$\begin{aligned} t_n^s \|U^n\|^2 + k \sum_{j=m}^n t_j^s \|U^j\|_1^2 &\leq C \sum_{j=m-p}^{m-1} \left( \|U^j\|_{-s}^2 + k^s \|U^j\|^2 \right) \\ &\quad + Ck \sum_{j=m}^n \left( \|f^j\|_{-s-1}^2 + t_j^s \|f^j\|_{-1}^2 \right), \quad \text{for } n \geq m. \end{aligned} \quad (3.2)$$

We have the following error estimate of time derivative approximation in nonsmooth data case.

THEOREM 3.3. *Let  $p \leq 6$  and let  $U^n$  and  $u$  be the solutions of (1.5) and (1.1), respectively. Then, with  $n \geq 2p$ ,*

$$t_n^{2p+2} \|\bar{\partial}_p U^n - u_t(t_n)\|^2 \leq C \sum_{j=p}^{2p-1} \left( \|U^j - u^j\|_{-2p}^2 + k^{2p+2} \|A(U^j - u^j)\|^2 \right) + Ck^{2p} G(u), \quad (3.3)$$

where

$$G(u) = \int_0^{t_n} \left( \|u^{(p+1)}(s)\|_{-2p-1}^2 + s^{2p+2} \|u^{(p+1)}(s)\|_1^2 + s^2 \|u_t(s)\|_1^2 \right) ds + t_{2p}^3 \|u_t(t_{2p})\|_1^2. \quad (3.4)$$

*Proof.* The error  $\varepsilon^n = \bar{\partial}_p U^n - u_t(t_n)$  ( $n \geq p$ ) satisfies

$$\bar{\partial}_p \varepsilon^n + A \varepsilon^n = -\tau^n, \quad \text{where } \tau^n = A(\bar{\partial}_p u(t_n) - u_t(t_n)), \text{ for } n \geq 2p. \quad (3.5)$$

Applying Lemma 3.2 with  $s = 2p+2$ ,  $m = 2p$ , we have, for  $n \geq 2p$ ,

$$t_n^{2p+2} \|\varepsilon^n\|^2 \leq C \sum_{j=p}^{2p-1} \left( \|\varepsilon^j\|_{-2p-2}^2 + k^{2p+2} \|\varepsilon^j\|^2 \right) + Ck \sum_{j=2p}^n \left( \|\tau^j\|_{-2p-3}^2 + t_j^{2p+2} \|\tau^j\|_{-1}^2 \right). \quad (3.6)$$

We now estimate the term  $k \sum_{j=2p}^n \|\tau^j\|_{-2p-3}^2$ . We will show that, with any norm  $\|\cdot\|$  in  $H$ ,

$$\|\bar{\partial}_p u(t_j) - u_t(t_j)\| \leq Ck^{p-1} \int_{t_{j-p}}^{t_j} \|u^{(p+1)}(s)\| ds, \quad \text{for } j \geq 2p. \quad (3.7)$$

Assuming this, we have

$$\|\tau^j\|_{-2p-3}^2 \leq Ck^{2p-1} \int_{t_{j-p}}^{t_j} \|u^{(p+1)}(s)\|_{-2p-1}^2 ds, \quad \text{for } j \geq 2p. \quad (3.8)$$

Thus

$$\begin{aligned} k \sum_{j=2p}^n |\tau^j|_{-2p-3}^2 &\leq Ck^{2p} \sum_{j=2p}^n \int_{t_{j-p}}^{t_j} |u^{(p+1)}(s)|_{-2p-1}^2 ds \\ &\leq Ck^{2p} \int_0^{t_n} |u^{(p+1)}(s)|_{-2p-1}^2 ds. \end{aligned} \quad (3.9)$$

It remains to estimate  $k \sum_{j=2p}^n t_j^{2p+2} |\tau^j|_{-1}^2$ . If  $j \neq 2p$ , we have, by (3.7) with norm  $\|A^{1/2} \cdot\|$ ,

$$k \sum_{j=2p+1}^n t_j^{2p+2} |\tau^j|_{-1}^2 \leq Ck^{2p} \sum_{j=2p+1}^n t_j^{2p+2} \int_{t_{j-p}}^{t_j} |u^{(p+1)}(s)|_1^2 ds. \quad (3.10)$$

Here we have  $t_j \leq cs$  for  $s \in [t_{j-p}, t_j]$ ,  $j \geq 2p+1$ , which follows from

$$t_j \leq s \frac{t_j}{t_{j-p}} \leq s \frac{t_{2p+1}}{t_{p+1}} \leq cs, \quad \text{for } j \geq 2p+1. \quad (3.11)$$

Hence

$$k \sum_{j=2p+1}^n t_j^{2p+2} |\tau^j|_{-1}^2 \leq Ck^{2p} \sum_{j=2p+1}^n \int_{t_{j-p}}^{t_j} s^{2p+2} |u^{(p+1)}(s)|_1^2 ds. \quad (3.12)$$

For  $j = 2p$ , we write, since  $\sum_{\nu=0}^p c_\nu = 0$ ,

$$\tau^{2p} = k^{-1} A \left( \sum_{\nu=0}^p c_\nu u(t_{2p-\nu}) - u_t(t_{2p}) \right) = k^{-1} A \left( \sum_{\nu=0}^p c_\nu \int_{t_p}^{t_{2p-\nu}} u_t(s) ds - u_t(t_{2p}) \right), \quad (3.13)$$

and we obtain

$$k |\tau^{2p}|_{-1}^2 \leq C \int_{t_p}^{t_{2p}} |u_t(s)|_1^2 ds + k |u_t(t_{2p})|_1^2, \quad (3.14)$$

which follows from

$$\begin{aligned} |\tau^{2p}|_{-1}^2 &\leq C \left( k^{-2} \sum_{\nu=0}^p \left| \int_{t_p}^{t_{2p-\nu}} u_t(s) ds \right|_1^2 + |u_t(t_{2p})|_1^2 \right) \\ &\leq Ck^{-2} \sum_{\nu=0}^p (pk) \int_{t_p}^{t_{2p-\nu}} |u_t(s)|_1^2 ds + |u_t(t_{2p})|_1^2 \\ &\leq Ck^{-1} \int_{t_p}^{t_{2p}} |u_t(s)|_1^2 ds + |u_t(t_{2p})|_1^2. \end{aligned} \quad (3.15)$$

Thus, we get

$$\begin{aligned} kt_{2p}^{2p+2} |\tau^{2p}|_{-1}^2 &\leq Ck^{2p+2} \left( \int_{t_p}^{t_{2p}} |u_t(s)|_1^2 ds + k |u_t(t_{2p})|_1^2 \right) \\ &\leq Ck^{2p} \left( \int_{t_p}^{t_{2p}} s^2 |u_t(s)|_1^2 ds + t_{2p}^3 |u_t(t_{2p})|_1^2 \right). \end{aligned} \quad (3.16)$$

It remains to estimate (3.7). We write, by Taylor expansion around  $t_{j-p}$ ,

$$u(t) = \sum_{l=0}^p \frac{u^{(l)}(t_{j-p})}{l!} (t - t_{j-p})^l + \frac{1}{p!} \int_{t_{j-p}}^t (t-s)^p u^{(p+1)}(s) ds \equiv Q(t) + R(t). \quad (3.17)$$

By (1.4) and since  $Q(t)$  is a polynomial of degree  $p$ , we have  $\bar{\partial}_p Q(t) - Q_t(t) = 0$ . Thus, by (2.1),

$$\bar{\partial}_p u(t_j) - u_t(t_j) = \bar{\partial}_p R(t_j) - R_t(t_j) = k^{-1} \sum_{\nu=0}^p c_\nu R(t_{j-\nu}) - R_t(t_j). \quad (3.18)$$

Noting that

$$\begin{aligned} \|R(t_{j-\nu})\| &\leq Ck^p \int_{t_{j-p}}^{t_j} \|u^{(p+1)}(s)\| ds, \quad \text{for } 0 \leq \nu \leq p, \quad j \geq 2p, \\ \|R_t(t_j)\| &= \frac{1}{(p-1)!} \left\| \int_{t_{j-p}}^{t_j} (t_j - s)^{p-1} u^{(p+1)}(s) ds \right\| \leq Ck^{p-1} \int_{t_{j-p}}^{t_j} \|u^{(p+1)}(s)\| ds, \end{aligned} \quad (3.19)$$

we complete the proof of (3.7).

Together these estimates complete the proof.  $\square$

In the homogeneous case, that is,  $f = 0$ , we have the following nonsmooth data error estimates.

**COROLLARY 3.4.** *Let  $p \leq 6$  and let  $U^n$  and  $u$  be the solutions of (1.5) and (1.1), respectively. Assume that  $f = 0$  and the discrete initial values satisfy, with  $U^0 = v$ ,*

$$|U^j - u^j|_{-2p} + k^{p+1} \|A(U^j - u^j)\| \leq Ck^p \|v\|, \quad \text{for } p \leq j \leq 2p-1. \quad (3.20)$$

Then, with  $C$  independent of the positive definite operator  $A$ ,

$$\|\bar{\partial}_p U^n - u_t(t_n)\| \leq Ck^p t_n^{-p-1} \|v\|, \quad \text{for } n \geq 2p. \quad (3.21)$$

*Proof.* For the solution  $u$  of homogeneous parabolic equation, it is easy to show that

$$\int_0^{t_n} |u^{(p+1)}(s)|_{-2p-1}^2 ds \leq C\|v\|^2, \quad \int_0^{t_n} s^{2p+2} |u^{(p+1)}(s)|_1^2 ds \leq C\|v\|^2, \quad (3.22)$$

and  $t_{2p}^3 |u_t(t_{2p})|_1^2 \leq C\|v\|^2$ . Applying Theorem 3.3, we complete the proof.  $\square$

Next we will consider the starting value approximation. In Theorem 3.3, we see that it is necessary to define starting approximations  $\{U^j\}_{j=0}^{p-1}$  such that

$$|U^j - u^j|_{-2p} + k^{p+1} \|A(U^j - u^j)\| = O(k^p), \quad \text{for } p \leq j \leq 2p-1. \quad (3.23)$$

Here we will consider the cases  $p = 1, 2$ . The approach can be extended to the general case for  $p > 2$ , but the proof is more complicated.

In the case of  $p = 1$ , the approximate solution is defined by the backward Euler method

$$\bar{\partial}_1 U^n + AU^n = f^n, \quad \text{for } n \geq 1, \text{ with } U^0 = \nu, \quad (3.24)$$

or, with  $r(\lambda) = 1/(1+\lambda)$ ,

$$U^n = r(kA)U^{n-1} + kr(kA)f^n, \quad \text{for } n \geq 1, \text{ with } U^0 = \nu. \quad (3.25)$$

We then have the following lemma, see Thomée [11, Theorem 9.1].

LEMMA 3.5. *Let  $U^1$  and  $u$  be the solutions of (3.24) and (1.1), respectively. Then, with  $u^1 = u(t_1)$ ,  $U^0 = u^0 = \nu$ ,*

$$\begin{aligned} & |U^1 - u^1|_{-2} + k^2 \|A(U^1 - u^1)\| \\ & \leq Ck \|\nu - A^{-1}f(0)\| + Ck \int_0^k \|A^{-1}f'(\tau)\| d\tau + Ck^2 \int_0^k \|f'(\tau)\| d\tau. \end{aligned} \quad (3.26)$$

In particular, if  $f = 0$ , then

$$|U^1 - u^1|_{-2} + k^2 \|A(U^1 - u^1)\| \leq Ck \|\nu\|. \quad (3.27)$$

We now turn to the case  $p = 2$ . In this case, we need two starting values  $U^0$  and  $U^1$ . We will use the backward Euler method to compute  $U^1$ , that is, the approximation  $U^n$  of the solution  $u(t_n)$  of (1.1) is defined by

$$\bar{\partial}_2 U^n + AU^n = f^n, \quad \text{for } n \geq 2, \quad \bar{\partial} U^1 + AU^1 = f^1, \quad \text{with } U^0 = \nu. \quad (3.28)$$

We have the following lemma.

LEMMA 3.6. *Let  $U^j$ ,  $j = 2, 3$  and  $u$  be the solutions of (3.28) and (1.1), respectively. Then, with  $u^j = u(t_j)$ ,  $U^0 = u^0 = \nu$ ,*

$$|U^j - u^j|_{-4} + k^3 \|A(U^j - u^j)\| \leq Ck^2 \left( \|\nu\| + \|f(0)\| + \int_0^{t_j} \|f'(\tau)\| d\tau \right), \quad j = 2, 3. \quad (3.29)$$

In particular, if  $f = 0$ , then

$$|U^j - u^j|_{-4} + k^3 \|A(U^j - u^j)\| \leq Ck^2 \|\nu\|, \quad j = 2, 3. \quad (3.30)$$

*Proof.* Here we only prove the case  $j = 2$ , that is, we will show that

$$|U^2 - u^2|_{-4} + k^3 \|A(U^2 - u^2)\| = O(k^2). \quad (3.31)$$

The proof for the case  $j = 3$  is similar.

Since  $\bar{\partial}_2 U^2 = k^{-1}((3/2)U^2 - 2U^1 + (1/2)U^0)$ , we may write

$$U^2 = q_1(kA)U^1 + q_2(kA)U^0 + kP(kA)f^2, \quad (3.32)$$

where

$$q_1(\lambda) = \frac{2}{3/2 + \lambda}, \quad q_2(\lambda) = \frac{-1/2}{3/2 + \lambda}, \quad P(\lambda) = \frac{1}{3/2 + \lambda}. \quad (3.33)$$

Thus, noting that  $u^2 = e^{-2kA}\nu + \int_0^{2k} e^{-(2k-s)A}f(s)ds$ , we have

$$U^2 - u^2 = q_1(kA)(U^1 - u^1) + q_2(kA)(U^0 - u^0) + E_2. \quad (3.34)$$

Here, by simple calculation,

$$\begin{aligned} E_2 &= q_1(kA)u^1 + q_2(kA)u^0 + kP(kA)f^2 - u^2 \\ &= Q(kA)\nu + kb_0(kA)f(0) + kR(f), \end{aligned} \quad (3.35)$$

where

$$\begin{aligned} Q(\lambda) &= q_1(\lambda)e^{-\lambda} + q_2(\lambda) - e^{-2\lambda}, \\ b_0(\lambda) &= q_1(\lambda) \int_0^1 e^{-(1-s)\lambda} ds + P(\lambda) - 2 \int_0^1 e^{-2(1-s)\lambda} ds, \\ R(f) &= q_1(kA) \int_0^1 e^{-(1-s)kA} \left( \int_0^{ks} f'(\tau) d\tau \right) ds \\ &\quad + P(kA) \int_0^{2k} f'(\tau) d\tau - 2 \int_0^1 e^{-2(1-s)kA} \left( \int_0^{2ks} f'(\tau) d\tau \right) ds. \end{aligned} \quad (3.36)$$

We first show that

$$k^3 \|A(U^2 - u^2)\| \leq Ck^2 \left( \|\nu\| + k\|f(0)\| + k \int_0^{2k} \|f'(\tau)\| d\tau \right). \quad (3.37)$$

In fact, noting  $U^0 = u^0 = \nu$ ,

$$\begin{aligned} k^3 \|A(U^2 - u^2)\| &\leq k^3 \|Aq_1(kA)(U^1 - u^1)\| + k^3 \|AQ(kA)\nu\| \\ &\quad + k^3 \|kAb_0(kA)f(0)\| + k^3 \|kAR(f)\| \\ &= I + II + III + IV. \end{aligned} \quad (3.38)$$

Since  $|\lambda Q(\lambda)| \leq C$  and  $|\lambda b_0(\lambda)| \leq C$  for  $0 \leq \lambda < \infty$ , we have

$$II \leq Ck^2 \|\nu\|, \quad III \leq Ck^3 \|f(0)\|. \quad (3.39)$$

By (3.36), it is easy to see that

$$IV \leq Ck^3 \int_0^{2k} \|f'(\tau)\| d\tau. \quad (3.40)$$

Now we turn to  $I$ . Note that

$$\begin{aligned} U^1 - u^1 &= (r(kA) - e^{-kA})v + kr(kA)f^1 - \int_0^k e^{-(k-s)A} f(s)ds \\ &= (r(kA) - e^{-kA})v + kr(kA) \left( f(0) + \int_0^k f'(\tau) d\tau \right) \\ &\quad - k \int_0^1 e^{-(1-s)kA} \left( f(0) + \int_0^{ks} f'(\tau) d\tau \right) ds. \end{aligned} \quad (3.41)$$

We find

$$I = k^3 \|Aq_1(kA)(U^1 - u^1)\| \leq Ck^2 \left( \|v\| + k\|f(0)\| + k \int_0^k \|f'\| d\tau \right). \quad (3.42)$$

Combining this with the estimates for  $II, III$ , and  $IV$ , we obtain (3.37).

We next show that

$$|A(U^2 - u^2)|_{-6} \leq Ck^2 \left( \|v\| + \int_0^{2k} \|f'(\tau)\| d\tau \right). \quad (3.43)$$

As in the proof of (3.37), we write

$$\begin{aligned} |A(U^2 - u^2)|_{-6} &\leq |Aq_1(kA)(U^1 - u^1)|_{-6} + |AQ(kA)v|_{-6} \\ &\quad + |kAb_0(kA)f(0)|_{-6} + |kAR(f)|_{-6} \\ &= I' + II' + III' + IV'. \end{aligned} \quad (3.44)$$

Since  $|\lambda^{-2}Q(\lambda)| < C$  and  $|\lambda^{-1}b_0(\lambda)| \leq C$  for  $0 \leq \lambda < \infty$ , we get

$$\begin{aligned} II' &= k^2 \|(kA)^{-2}Q(kA)v\| \leq Ck^2 \|v\|, \\ III' &= k^2 \|(kA)^{-1}b_0(kA)A^{-1}\| \leq Ck^2 \|f(0)\|. \end{aligned} \quad (3.45)$$

Further, we easily find that

$$\begin{aligned} IV' &\leq Ck^2 \int_0^{2k} \|f'(\tau)\| d\tau, \\ I' &\leq Ck^2 \left( \|v\| + \|f(0)\| + \int_0^k \|f'(\tau)\| d\tau \right). \end{aligned} \quad (3.46)$$

Hence (3.43) follows.

Together, these estimates show (3.31). The proof is complete.  $\square$

Combining Theorem 3.3 with Lemma 3.6, we get the following error estimate in the case  $p = 2$ .

**COROLLARY 3.7.** *Let  $U^n$  and  $u$  be the solutions of (3.28) and (1.1), respectively. Then, with  $U^0 = u^0 = v$ ,*

$$\|\bar{\partial}_2 U^n - u_t(t_n)\| \leq Ck^2 t_n^{-3} \left( \|v\| + \|f(0)\| + \int_0^{t_3} \|f'(\tau)\| d\tau + K(u) \right), \quad n \geq 4, \quad (3.47)$$

where

$$K(u)^2 = \int_0^{t_n} \left( |u^{(3)}(s)|_{-5}^2 + s^6 |u^{(3)}(s)|_1^2 + s^2 |u_t(s)|_1^2 \right) ds + t_4^3 |u_t(t_4)|_1^2. \quad (3.48)$$

## References

- [1] J. Becker, *A second order backward difference method with variable steps for a parabolic problem*, BIT **38** (1998), no. 4, 644–662.
- [2] J. H. Bramble, J. E. Pasciak, P. H. Sammon, and V. Thomée, *Incomplete iterations in multistep backward difference methods for parabolic problems with smooth and nonsmooth data*, Math. Comp. **52** (1989), no. 186, 339–367.
- [3] M. Crouzeix, *On multistep approximation of semigroups in Banach spaces*, J. Comput. Appl. Math. **20** (1987), no. Special Issue, 25–35.
- [4] E. Hairer and G. Wanner, *Solving Ordinary Differential Equations. II*, Springer Series in Computational Mathematics, vol. 14, Springer, Berlin, 1991, Stiff and differential-algebraic problems.
- [5] A. Hansbo, *Nonsmooth data error estimates for damped single step methods for parabolic equations in Banach space*, Calcolo **36** (1999), no. 2, 75–101.
- [6] ———, *Strong stability and non-smooth data error estimates for discretizations of linear parabolic problems*, BIT **42** (2002), no. 2, 351–379.
- [7] M.-N. Le Roux, *Semi-discrétisation en temps pour les équations d'évolution paraboliques lorsque l'opérateur dépend du temps*, RAIRO Anal. Numér. **13** (1979), no. 2, 119–137.
- [8] ———, *Semidiscretization in time for parabolic problems*, Math. Comp. **33** (1979), no. 147, 919–931.
- [9] C. Palencia and B. García-Archilla, *Stability of linear multistep methods for sectorial operators in Banach spaces*, Appl. Numer. Math. **12** (1993), no. 6, 503–520.
- [10] G. Savaré,  *$A(\Theta)$ -stable approximations of abstract Cauchy problems*, J. Numer. Math. **65** (1993), no. 3, 319–335.
- [11] V. Thomée, *Galerkin Finite Element Methods For Parabolic Problems*, Springer Series in Computational Mathematics, vol. 25, Springer, Berlin, 1997.
- [12] Y. Yan, *Smoothing A properties and approximation of time derivatives for parabolic equations: variable time steps*, BIT **43** (2003), no. 3, 647–669.
- [13] ———, *Smoothing properties and approximation of time derivatives for parabolic equations: constant time steps*, IMA J. Numer. Anal. **23** (2003), no. 3, 465–487.

Yubin Yan: Department of Mathematics, The University of Manchester, M13 9PL, Manchester, UK  
 Current address: Department of Automatic Control and Systems Engineering, Faculty of Engineering, The University of Sheffield, Mappin Street, Sheffield, S1 3JD, UK

E-mail address: y.yan@sheffield.ac.uk

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)