

STARLIKENESS ASSOCIATED WITH PARABOLIC REGIONS

ROSIHAN M. ALI

Received 29 April 2004 and in revised form 15 September 2004

A parabolic starlike function f of order ρ in the unit disk is characterized by the fact that the quantity $zf'(z)/f(z)$ lies in a given parabolic region in the right half-plane. Denote the class of such functions by $PS^*(\rho)$. This class is contained in the larger class of starlike functions of order ρ . Subordination results for $PS^*(\rho)$ are established, which yield sharp growth, covering, and distortion theorems. Sharp bounds for the first four coefficients are also obtained. There exist different extremal functions for these coefficient problems. Additionally, we obtain a sharp estimate for the Fekete-Szegö coefficient functional and investigate convolution properties for $PS^*(\rho)$.

1. Introduction

Let A denote the class of analytic functions f in the open unit disk $U = \{z : |z| < 1\}$ and let f be normalized so that $f(0) = f'(0) - 1 = 0$. In [4], Goodman introduced the class UCV of uniformly convex functions consisting of convex functions $f \in A$ with the property that for every circular arc γ contained in U , with center also in U , the image arc $f(\gamma)$ is a convex arc. He derived a two-variable characterization of functions in UCV , that is, $f \in A$ belongs to UCV if and only if for every pair $(z, \varsigma) \in U \times U$,

$$1 + \Re \left\{ (z - \varsigma) \frac{f''(z)}{f'(z)} \right\} \geq 0. \quad (1.1)$$

Ma and Minda [6] and Rønning [10] independently developed a one-variable characterization that $f \in UCV$ if and only if for every $z \in U$,

$$\left| \frac{zf''(z)}{f'(z)} \right| < \Re \left(1 + \frac{zf''(z)}{f'(z)} \right). \quad (1.2)$$

Rønning [10] also showed that $f \in UCV$ if and only if the function $zf' \in PS^*$, where PS^* is the class of functions $g \in A$ satisfying

$$\left| \frac{zg'(z)}{g(z)} - 1 \right| < \Re \frac{zg'(z)}{g(z)}, \quad z \in U. \quad (1.3)$$

Several authors have studied the classes above, amongst which the authors of [4, 6, 7, 8, 9, 10, 12].

In [9], the class PS^* was generalized by looking at functions $f \in A$ satisfying

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \Re \frac{zf'(z)}{f(z)} - \alpha, \quad z \in U. \quad (1.4)$$

In this paper, we continue the investigation of this generalized class but under a slight modification of parameter. For $0 \leq \rho < 1$, let Ω_ρ be the parabolic region in the right half-plane

$$\Omega_\rho = \{w = u + iv : v^2 < 4(1 - \rho)(u - \rho)\} = \{w : |w - 1| < 1 - 2\rho + \Re w\}. \quad (1.5)$$

The class of *parabolic starlike functions of order ρ* is the subclass $PS^*(\rho)$ of A consisting of functions f such that $zf'(z)/f(z) \in \Omega_\rho$, $z \in U$. Thus $f \in PS^*(\rho)$ if and only if for $z \in U$,

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < 1 - 2\rho + \Re \frac{zf'(z)}{f(z)}. \quad (1.6)$$

Similarly, a function $f \in A$ belongs to $UCV(\rho)$ if and only if for every pair (z, ς) in the polydisk $U \times U$,

$$1 + \Re \left\{ (z - \varsigma) \frac{f''(z)}{f'(z)} \right\} > 2\rho - 1. \quad (1.7)$$

A function $f \in UCV(\rho)$ is called an *uniformly convex function of order ρ* . Thus the classes discussed earlier correspond to $UCV = UCV(1/2)$ and $PS^* = PS^*(1/2)$. In [5], Lee showed that

$$g \in UCV(\rho) \iff f = zg' \in PS^*(\rho), \quad (1.8)$$

that is,

$$g \in UCV(\rho) \iff \left| \frac{zg''(z)}{g'(z)} \right| < 2(1 - \rho) + \Re \frac{zg''(z)}{g'(z)}. \quad (1.9)$$

In the present paper, we continue the study of $PS^*(\rho)$ realized by Ali and Singh [3], and more recently by Aghalary and Kulkarni [1]. We give examples of functions in the class $PS^*(\rho)$, and establish subordination results, which yield sharp growth, covering and distortion theorems. Sharp bounds on the first four coefficients are also obtained. There exist different extremal functions for these coefficient problems. Additionally, we obtain a sharp estimate for the Fekete-Szegö coefficient functional and examine convolution properties for $PS^*(\rho)$.

2. Preliminary results

From its definition, it is clear that the class $PS^*(\rho)$ is contained in the class $S^*(\rho)$ of starlike functions of order ρ , that is, $\Re(zf'(z)/f(z)) > \rho$, $z \in U$. It is also fairly immediate

that $\text{PS}^*(\rho)$ is related to the class of strongly starlike functions, where a function $f \in A$ is said to be strongly starlike of order α , $0 < \alpha \leq 1$, if f satisfies $|\text{Arg}zf'(z)/f(z)| < \pi\alpha/2$, $z \in U$. We state the relation in the theorem below.

THEOREM 2.1. *If $f \in \text{PS}^*(\rho)$, then f is strongly starlike of order γ , where $(\pi/2)\gamma = \tan^{-1} \sqrt{(1-\rho)/\rho}$. In other words, for $z \in U$,*

$$\left| \text{Arg} \frac{zf'(z)}{f(z)} \right| \leq \frac{\pi\gamma}{2}. \quad (2.1)$$

A sufficient condition for a function f to be parabolic starlike of order ρ is given by the following theorem.

THEOREM 2.2. *If $f \in A$ satisfies*

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < 1 - \rho, \quad (2.2)$$

then $f \in \text{PS}^(\rho)$.*

Proof. The given condition implies that

$$\Re \frac{zf'(z)}{f(z)} - \left| \frac{zf'(z)}{f(z)} - 1 \right| + 1 - 2\rho \geq 2(1-\rho) - 2 \left| \frac{zf'(z)}{f(z)} - 1 \right| > 0. \quad (2.3)$$

□

The following two examples are now easily established from Theorem 2.2.

Example 2.3. The function $f(z) = z + \alpha z^n \in \text{PS}^*(\rho)$ if and only if $|\alpha| \leq (1-\rho)/(n-\rho)$.

Example 2.4. The generalized hypergeometric function is defined by

$$F(a_1, \dots, a_p; b_1, \dots, b_q; z) = 1 + \sum_{n=1}^{\infty} \frac{(a_1)_n \cdots (a_p)_n}{(b_1)_n \cdots (b_q)_n} \frac{z^n}{n!}, \quad b_j \neq 0, -1, \dots, \quad (2.4)$$

where $(\lambda)_n$ is the Pochhammer symbol defined by

$$(\lambda)_n = \begin{cases} 1, & n = 0, \\ \lambda(\lambda+1)(\lambda+2) \cdots (\lambda+n-1), & n = 1, 2, \dots. \end{cases} \quad (2.5)$$

If $|zF'(z)/F(z)| < 1 - \rho$, then $zF \in \text{PS}^*(\rho)$.

Ali and Singh [3] showed that the normalized Riemann mapping function q_ρ from U onto Ω_ρ is given by

$$q_\rho(z) = 1 + \frac{4(1-\rho)}{\pi^2} \left[\log \frac{1+\sqrt{z}}{1-\sqrt{z}} \right]^2 = 1 + \sum_{n=1}^{\infty} B_n z^n. \quad (2.6)$$

Here

$$B_1 = \frac{16(1-\rho)}{\pi^2}, \quad B_n = \frac{16(1-\rho)}{n\pi^2} \sum_{k=0}^{n-1} \frac{1}{2k+1}, \quad n = 2, 3, \dots \quad (2.7)$$

Since the latter sum is bounded above by $1 + (1/2)\log(2n-1)$ (see [6]) an upper bound for each coefficient is given by

$$B_n < \frac{16(1-\rho)}{n\pi^2} \left(1 + \frac{1}{2} \log(2n-1) \right). \quad (2.8)$$

However these bounds do not yield sharp coefficient estimates for the class $PS^*(\rho)$. We will return to the coefficient problem in the next section.

Let $k \in PS^*(\rho)$ be defined by $k(0) = k'(0) - 1 = 0$ and

$$\frac{zk'(z)}{k(z)} = q_\rho(z). \quad (2.9)$$

In [8], Ma and Minda established a general result that leads to the following result.

THEOREM 2.5 [8]. *If $f \in PS^*(\rho)$, then*

- (a) $zf'(z)/f(z) \prec zk'(z)/k(z)$ and $f(z)/z \prec k(z)/z$,
- (b) $-k(-r) \leq |f(z)| \leq k(r)$, $|z| \leq r < 1$,
- (c) $|\operatorname{Arg}(f(z)/z)| \leq \max_{|z|=r} |\operatorname{Arg}(k(z)/z)|$, $|z| \leq r < 1$,
- (d) $k'(-r) \leq |f'(z)| \leq k'(r)$, $|z| \leq r < 1$.

Equality in (b), (c), and (d) holds for some $z \neq 0$ if and only if f is a rotation of k .

Since the function k is continuous in \overline{U} , $-k(-1) = \lim_{r \rightarrow 1^-} -k(-r)$ and $k(1) = \lim_{r \rightarrow 1^-} k(r)$ exist. Rønning [9] established the following corollary.

COROLLARY 2.6 [9]. *(a) Let $f \in PS^*(\rho)$. Then either f is a rotation of k or $f(U) \supset \{w : |w| \leq -k(-1)\}$, where the Koebe constant is $-k(-1) = e^{-(1-\rho)(1.25475)}$.*

(b) The functions in $PS^(\rho)$ are uniformly bounded by the sharp constant $k(1) = e^{3.41023(1-\rho)}$.*

3. Coefficient bounds

We first give another sufficient condition for a function f to belong to $PS^*(\rho)$.

THEOREM 3.1. *If $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ satisfies $\sum_{n=2}^{\infty} (n-1)|a_n| \leq (1-\rho)/(2-\rho)$, then $f \in PS^*(\rho)$. The constant $(1-\rho)/(2-\rho)$ cannot be replaced by a larger number.*

Proof. Let $g(z) = \int_0^z (f(\xi)/\xi) d\xi = z + \sum_{n=2}^{\infty} (a_n/n) z^n$. In view of (1.8), it suffices to show that $g \in UCV(\rho)$. Since

$$\sum_{n=2}^{\infty} |a_n| \leq \frac{1-\rho}{2-\rho}, \quad (3.1)$$

it follows that

$$1 + \Re(z - \zeta) \frac{g''(z)}{g'(z)} \geq 1 - \frac{\sum_{n=2}^{\infty} (n-1) |a_n| |z|^{n-2}}{1 - \sum_{n=2}^{\infty} |a_n| |z|^{n-1}} |z - \zeta| \geq 2\rho - 1. \quad (3.2)$$

Thus $g \in \text{UCV}(\rho)$. The function $f(z) = z + ((1-\rho)/(2-\rho))z^2$ in Example 2.3 shows that the constant $(1-\rho)/(2-\rho)$ is the best possible. \square

We next consider the problem of finding

$$A_n = \max_{f \in \text{PS}^*(\rho)} |a_n|. \quad (3.3)$$

If $f(z) = z + a_2 z^2 + a_3 z^3 + \dots \in \text{PS}^*(\rho)$ and $h(z) = zf'(z)/f(z)$, then there exists a Schwarz function w defined in U with $w(0) = 0$, $|w(z)| < 1$, and satisfying

$$h(z) = \frac{zf'(z)}{f(z)} = q_\rho(w(z)). \quad (3.4)$$

If $h(z) = 1 + b_1 z + b_2 z^2 + \dots$, the first equality in (3.4) implies that

$$(n-1)a_n = \sum_{k=1}^{n-1} a_k b_{n-k}. \quad (3.5)$$

Since q_ρ is univalent in U and $h \prec q_\rho$, the function

$$p(z) = \frac{1 + q_\rho^{-1}(h(z))}{1 - q_\rho^{-1}(h(z))} = 1 + c_1 z + c_2 z^2 + \dots \quad (3.6)$$

belongs to the class P consisting of analytic functions p in the unit disk U with positive real part such that $p(0) = 1$ and $\Re p(z) > 0$, $z \in U$. In other words,

$$h(z) = q_\rho \left(\frac{p(z) - 1}{p(z) + 1} \right). \quad (3.7)$$

While (3.5) gives a_n in terms of the coefficients b_k , (3.7) expresses the b_k 's in terms of the coefficients c_m 's and B_m 's. It is now easily established that

$$\begin{aligned} a_2 &= \frac{8(1-\rho)}{\pi^2} c_1, \\ a_3 &= \frac{8(1-\rho)}{2\pi^2} \left[c_2 - \left(\frac{1}{6} - \frac{8(1-\rho)}{\pi^2} \right) c_1^2 \right], \\ a_4 &= \frac{8(1-\rho)}{3\pi^2} \left[c_3 - \left(\frac{1}{3} - \frac{12(1-\rho)}{\pi^2} \right) c_1 c_2 + \left(\frac{2}{45} - \frac{2(1-\rho)}{\pi^2} + \frac{32(1-\rho)^2}{\pi^4} \right) c_1^3 \right]. \end{aligned} \quad (3.8)$$

Thus the coefficient estimates for $\text{PS}^*(\rho)$ may be viewed in terms of nonlinear coefficient problems for the class P .

We now introduce the following functions in $\text{PS}^*(\rho)$. Define $k_n, G, H \in A$, respectively, by

$$\frac{zk'_n(z)}{k_n(z)} = q_\rho(z^{n-1}), \quad \frac{zH'(z)}{H(z)} = q_\rho\left(\frac{z(z-r)}{1-rz}\right), \quad \frac{zG'(z)}{G(z)} = q_\rho\left(-\frac{z(z-r)}{1-rz}\right), \quad 0 \leq r \leq 1. \quad (3.9)$$

It is clear from (3.4) that $k_n, G, H \in \text{PS}^*(\rho)$, and that $k_2(z) = k(z)$. Since

$$k_n(z) = z + \frac{16(1-\rho)}{(n-1)\pi^2} z^n + \dots, \quad (3.10)$$

we find that

$$A_n \geq \frac{16(1-\rho)}{(n-1)\pi^2}. \quad (3.11)$$

On the other hand, Ali and Singh [3] proved that

$$(n-1)A_n \leq 2\sqrt{2}(1-\rho)e^{4(1-\rho)^2}, \quad (3.12)$$

which also yields the sharp order of growth $|a_n| = O(1/n)$.

From a result of Ma and Minda [8], we can also deduce the following solution to the Fekete-Szegő coefficient functional over the class $\text{PS}^*(\rho)$. We will omit the details.

THEOREM 3.2. *Let $f(z) = z + a_2 z^2 + a_3 z^3 + \dots \in \text{PS}^*(\rho)$. Then*

$$|a_3 - ta_2^2| \leq \begin{cases} \frac{16(1-\rho)}{3\pi^4} [24(1-\rho)(1-2t) + \pi^2], & t \leq \frac{1}{2} - \frac{\pi^2}{96(1-\rho)}, \\ \frac{8(1-\rho)}{\pi^2}, & \frac{1}{2} - \frac{\pi^2}{96(1-\rho)} \leq t \leq \frac{1}{2} + \frac{5\pi^2}{96(1-\rho)}, \\ \frac{16(1-\rho)}{3\pi^4} [24(1-\rho)(2t-1) - \pi^2], & t \geq \frac{1}{2} + \frac{5\pi^2}{96(1-\rho)}. \end{cases} \quad (3.13)$$

If $1/2 - \pi^2/96(1-\rho) < t < 1/2 + 5\pi^2/96(1-\rho)$, equality holds if and only if $f = k_3$ or one of its rotations. If $t < 1/2 - \pi^2/96(1-\rho)$ or $t > 1/2 + 5\pi^2/96(1-\rho)$, equality holds if and only if $f = k_2$ or one of its rotations. If $t = 1/2 - \pi^2/96(1-\rho)$, equality holds if and only if $f = H$ or one of its rotations, while if $t = 1/2 + 5\pi^2/96(1-\rho)$, then equality holds if and only if $f = G$ or one of its rotations.

The above estimates can be used to determine sharp upper bounds on the second and third coefficients, respectively, which we will state below. In addition, the sharp bound on the fourth coefficient A_4 is determined with the aid of the following lemma.

LEMMA 3.3 [2]. *Let $p(z) = 1 + \sum_{k=1}^{\infty} c_k z^k \in P$. If $0 \leq \beta \leq 1$ and $\beta(2\beta-1) \leq \delta \leq \beta$, then*

$$|c_3 - 2\beta c_1 c_2 + \delta c_1^3| \leq 2. \quad (3.14)$$

In particular,

$$|c_3 - 2\beta c_1 c_2 + \beta c_1^3| \leq 2. \quad (3.15)$$

When $\beta = 0$, equality holds if and only if

$$p(z) := p_3(z) = \sum_{k=1}^3 \lambda_k \frac{1 + \epsilon e^{-2\pi i k/3} z}{1 - \epsilon e^{-2\pi i k/3} z}, \quad |\epsilon| = 1, \lambda_k \geq 0, \quad (3.16)$$

with $\lambda_1 + \lambda_2 + \lambda_3 = 1$. If $\beta = 1$, equality holds if and only if p is the reciprocal of p_3 . If $0 < \beta < 1$, equality holds if and only if

$$p(z) = \frac{1 + \epsilon z}{1 - \epsilon z}, \quad |\epsilon| = 1 \quad \text{or} \quad p(z) = \frac{1 + \epsilon z^3}{1 - \epsilon z^3}, \quad |\epsilon| = 1. \quad (3.17)$$

THEOREM 3.4. Let $f(z) = z + a_2 z^2 + a_3 z^3 + \dots \in \text{PS}^*(\rho)$. Then

$$|a_2| \leq \frac{16(1-\rho)}{\pi^2}, \quad (3.18)$$

with equality if and only if $f = k$ or its rotations. Further

$$|a_3| \leq \begin{cases} \frac{8(1-\rho)}{\pi^2} \left(\frac{2}{3} + \frac{16(1-\rho)}{\pi^2} \right), & 0 \leq \rho \leq 1 - \frac{\pi^2}{48}, \\ \frac{8(1-\rho)}{\pi^2}, & 1 - \frac{\pi^2}{48} \leq \rho < 1. \end{cases} \quad (3.19)$$

For $0 \leq \rho < 1 - \pi^2/48$, equality holds if and only if $f = k$ or its rotations. For $1 - \pi^2/48 < \rho < 1$, equality holds if and only if $f = k_3$ or its rotations. If $\rho = 1 - \pi^2/48$, equality holds if and only if $f = H$ or its rotations. Additionally,

$$|a_4| \leq \begin{cases} \frac{16(1-\rho)}{3\pi^2} \left[\frac{128(1-\rho)^2}{\pi^4} + \frac{16(1-\rho)}{\pi^2} + \frac{23}{45} \right], & 0 \leq \rho \leq 1 + \frac{\pi^2}{16} \left(1 - \sqrt{\frac{89}{45}} \right), \\ \frac{16(1-\rho)}{3\pi^2}, & 1 + \frac{\pi^2}{16} \left(1 - \sqrt{\frac{89}{45}} \right) \leq \rho < 1. \end{cases} \quad (3.20)$$

Equality holds in the upper expression of the right inequality if and only if $f = k$ or its rotations, while equality holds in the lower expression of the right inequality if and only if $f = k_4$ or its rotations.

Proof. In the light of Theorem 3.2, we are left to finding an estimate on the fourth coefficient. The relation (3.8) gives

$$\begin{aligned} a_4 &= \frac{8(1-\rho)}{3\pi^2} \left[c_3 - \left(\frac{1}{3} - \frac{12(1-\rho)}{\pi^2} \right) c_1 c_2 + \left(\frac{2}{45} - \frac{2(1-\rho)}{\pi^2} + \frac{32(1-\rho)^2}{\pi^4} \right) c_1^3 \right] \\ &:= \frac{8(1-\rho)}{3\pi^2} E. \end{aligned} \quad (3.21)$$

We will apply Lemma 3.3 with

$$2\beta = \frac{1}{3} - \frac{12(1-\rho)}{\pi^2}, \quad \delta = \frac{2}{45} - \frac{2(1-\rho)}{\pi^2} + \frac{32(1-\rho)^2}{\pi^4}. \quad (3.22)$$

The conditions on β and δ are satisfied if

$$1 + \frac{\pi^2}{16} \left(1 - \sqrt{\frac{89}{45}} \right) \leq \rho < 1. \quad (3.23)$$

Thus $|a_4| \leq 16(1-\rho)/3\pi^2$, with equality if and only if the function p in (3.7) is given by $p(z) = (1+\epsilon z^3)/(1-\epsilon z^3)$. This implies that $f = k_4$.

In view of the fact that $0 < \delta < 1$, and that $\delta - \beta \geq 0$ provided

$$1 + \frac{\pi^2}{16} \left(1 - \sqrt{\frac{89}{45}} \right) \geq \rho, \quad (3.24)$$

Lemma 3.3 yields

$$\begin{aligned} |E| &\leq |c_3 - 2\delta c_1 c_2 + \delta c_1^3| + 2(\delta - \beta) |c_1 c_2| \\ &\leq 2 + 8 \left(\frac{32(1-\rho)^2}{\pi^4} + \frac{4(1-\rho)}{\pi^2} - \frac{11}{90} \right) \\ &= 2 \left(\frac{128(1-\rho)^2}{\pi^4} + \frac{16(1-\rho)}{\pi^2} + \frac{23}{45} \right). \end{aligned} \quad (3.25)$$

Equality holds if and only if the function p in (3.7) is given by $p(z) = (1+\epsilon z)/(1-\epsilon z)$, that is, $f = k$. This completes the proof. \square

THEOREM 3.5. *Let $f(z) = z + a_2 z^2 + a_3 z^3 + \dots \in \text{PS}^*(\rho)$. For $\mu \in C$ and*

$$\begin{aligned} \lambda(\mu) &= \frac{1}{3} + \frac{16(1-\rho)}{\pi^2} (2\mu - 1), \\ |a_3 - \mu a_2^2| &\leq \begin{cases} \frac{16(1-\rho)}{3\pi^4} |24(1-\rho)(1-2\mu) + \pi^2|, & |\lambda(\mu) - 1| \geq 1, \\ \frac{8(1-\rho)}{\pi^2}, & |\lambda(\mu) - 1| \leq 1. \end{cases} \end{aligned} \quad (3.26)$$

Equality holds in the upper expression of the right inequality if $f = k$ or its rotations, while equality holds in the lower expression of the right inequality if $f = k_3$ or its rotations.

Proof. From the relation (3.8), we get

$$a_3 - \mu a_2^2 = \frac{4(1-\rho)}{\pi^2} \left[c_2 - \frac{\lambda(\mu)}{2} c_1^2 \right]. \quad (3.27)$$

The well-known estimate

$$\left| c_2 - \frac{1}{2} c_1^2 \right| \leq 2 - \frac{1}{2} |c_1|^2 \quad (3.28)$$

leads to

$$\left| c_2 - \frac{\lambda(\mu)}{2} c_1^2 \right| \leq \left| c_2 - \frac{1}{2} c_1^2 \right| + \frac{|1 - \lambda(\mu)|}{2} |c_1|^2 \leq 2 + \frac{|\lambda(\mu) - 1| - 1}{2} |c_1|^2, \quad (3.29)$$

which yields the desired result. \square

4. Convolution properties

The convolution of $f(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=0}^{\infty} b_n z^n$ is defined to be the function $(f * g)(z) = \sum_{n=0}^{\infty} a_n b_n z^n$. For $\alpha < 1$, denote by R_{α} the class of prestarlike functions of order α consisting of $f \in A$ such that $f * ((z)/(1-z)^{2-2\alpha}) \in S^*(\alpha)$. Here $S^*(\alpha)$ is the class of starlike functions of order α . An important result in convolution is contained in the following lemma of Ruscheweyh.

LEMMA 4.1 [11, page 54]. *If $f \in R_{\alpha}$, $g \in S^*(\alpha)$, and H is an analytic function in U , then*

$$\frac{f * g H}{f * g}(U) \subset \overline{\text{co}}H(U), \quad (4.1)$$

where $\overline{\text{co}}H(U)$ is the closed convex hull of $H(U)$.

THEOREM 4.2. *If $f \in R_{\rho}$ and $g \in PS^*(\rho)$, then $f * g \in PS^*(\rho)$.*

Proof. Since g also belongs to $S^*(\rho)$ and $H(z) = zg'(z)/g(z) \prec q_{\rho}(z)$, Lemma 4.1 yields

$$\frac{z(f * g)'}{f * g}(U) = \frac{f * zg'}{f * g}(U) = \frac{f * g(zg'/g)}{f * g}(U) \subset \overline{\text{co}} \frac{zg'}{g}(U) \subset \overline{\Omega}_{\rho}, \quad (4.2)$$

and hence, $f * g \in PS^*(\rho)$. \square

Since $R_{1/2} = S^*(1/2)$ (see [11]), and $R_0 = C$, where C is the class of convex functions in A , a similar proof also yields the following result.

COROLLARY 4.3. (a) *If $f, g \in PS^*(\rho)$ for $\rho \geq 1/2$, then $f * g \in PS^*(\rho)$.*

(b) *If $f \in C$ and $g \in PS^*(\rho)$, then $f * g \in PS^*(\rho)$.*

Acknowledgments

This research was supported by a Universiti Sains Malaysia Fundamental Research Grant. The author is greatly indebted to Professor V. Ravichandran for his helpful comments in the preparation of this paper.

References

- [1] R. Aghalary and S. R. Kulkarni, *Certain properties of parabolic starlike and convex functions of order ρ* , Bull. Malays. Math. Sci. Soc. (2) **26** (2003), no. 2, 153–162.
- [2] R. M. Ali, *Coefficients of the inverse of strongly starlike functions*, Bull. Malays. Math. Sci. Soc. (2) **26** (2003), no. 1, 63–71.
- [3] R. M. Ali and V. Singh, *Coefficients of parabolic starlike functions of order ρ* , Computational Methods and Function Theory 1994 (Penang), Ser. Approx. Decompos., vol. 5, World Scientific Publishing, New Jersey, 1995, pp. 23–36.

570 Starlikeness associated with parabolic regions

- [4] A. W. Goodman, *On uniformly convex functions*, Ann. Polon. Math. **56** (1991), no. 1, 87–92.
- [5] S. K. Lee, *Characterizations of parabolic starlike functions and the generalized uniformly convex functions*, Master's thesis, Universiti Sains Malaysia, Penang, Malaysia, 2000.
- [6] W. C. Ma and D. Minda, *Uniformly convex functions*, Ann. Polon. Math. **57** (1992), no. 2, 165–175.
- [7] ———, *Uniformly convex functions. II*, Ann. Polon. Math. **58** (1993), no. 3, 275–285.
- [8] ———, *A unified treatment of some special classes of univalent functions*, Proceedings of the Conference on Complex Analysis (Tianjin, 1992), Conf. Proc. Lecture Notes Anal., I, International Press, Massachusetts, 1994, pp. 157–169.
- [9] F. Rønning, *On starlike functions associated with parabolic regions*, Ann. Univ. Mariae Curie-Skłodowska Sect. A **45** (1991), 117–122.
- [10] ———, *Uniformly convex functions and a corresponding class of starlike functions*, Proc. Amer. Math. Soc. **118** (1993), no. 1, 189–196.
- [11] S. Ruscheweyh, *Convolutions in Geometric Function Theory*, Séminaire de Mathématiques Supérieures, vol. 83, Presses de l'Université de Montréal, Quebec, 1982.
- [12] T. N. Shanmugam and V. Ravichandran, *Certain properties of uniformly convex functions*, Computational Methods and Function Theory 1994 (Penang), Ser. Approx. Decompos., vol. 5, World Scientific Publishing, New Jersey, 1995, pp. 319–324.

Rosihan M. Ali: School of Mathematical Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia

E-mail address: rosihan@cs.usm.my

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk