

*Research Article*  
**Integer Powers of Arcsin**

Jonathan M. Borwein and Marc Chamberland

Received 1 November 2006; Revised 28 February 2007; Accepted 18 April 2007

Recommended by Ahmed Zayed

New simple nested-sum representations for powers of the arcsin function are given. This generalization of Ramanujan's work makes connections to finite binomial sums and polylogarithms.

Copyright © 2007 J. M. Borwein and M. Chamberland. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

This note discovers, derives, and then studies *simple* closed-form Taylor series expressions for integer powers of  $\arcsin(x)$ . Specifically, we show that, for  $|x| \leq 2$  and  $N = 1, 2, \dots$ ,

$$\frac{\arcsin^{2N}(x/2)}{(2N)!} = \sum_{k=1}^{\infty} \frac{H_N(k)}{\binom{2k}{k} k^2} x^{2k}, \quad (1.1)$$

where  $H_1(k) = 1/4$  and

$$H_{N+1}(k) := \frac{1}{4} \sum_{n_1=1}^{k-1} \frac{1}{(2n_1)^2} \sum_{n_2=1}^{n_1-1} \frac{1}{(2n_2)^2} \cdots \sum_{n_N=1}^{n_{N-1}-1} \frac{1}{(2n_N)^2}, \quad (1.2)$$

and also that, for  $|x| \leq 2$  and  $N = 0, 1, 2, \dots$ ,

$$\frac{\arcsin^{2N+1}(x/2)}{(2N+1)!} = \sum_{k=0}^{\infty} \frac{G_N(k) \binom{2k}{k}}{2(2k+1)4^{2k}} x^{2k+1}, \quad (1.3)$$

where  $G_0(k) = 1$  and

$$G_N(k) := \sum_{n_1=0}^{k-1} \frac{1}{(2n_1+1)^2} \sum_{n_2=0}^{n_1-1} \frac{1}{(2n_2+1)^2} \cdots \sum_{n_N=0}^{n_{N-1}-1} \frac{1}{(2n_N+1)^2}. \quad (1.4)$$

The convention is that the sum is zero if the starting index exceeds the finishing index.

Nested sums are not new. The last decade saw many interesting results concerning *Euler sums* or *multizeta values*, wonderful generalizations of the classical  $\zeta$ -function, whose discovery can be traced to a letter from Goldbach to Euler [1, pages 99-100] and [2, Chapter 3]—a letter that played a seminal role in the discovery of the  $\zeta$ -function.

When Gauss was criticized for the lack of motivation in his writings, he remarked that the architects of great cathedrals do not obscure the beauty of their work by leaving the scaffolding in place after the construction has been completed. While we find (1.1) and (1.3) worthy of undistracted attention, in truth their discovery was greatly facilitated by the use of *experimental mathematics*—the relatively new approach to doing mathematical research with the intelligent use of computers. This perspective is elucidated throughout this paper. It is also illustrative of the changing speed of mathematical communication that the special cases (2.1), (2.2), and (2.3), given below, are already online at [3].

## 2. Experiments and proofs

The first identity below is very well known:

$$\arcsin^2\left(\frac{x}{2}\right) = \frac{1}{2} \sum_{k=1}^{\infty} \frac{x^{2k}}{\binom{2k}{k} k^2}. \quad (2.1)$$

It is explored at some length in [4, pages 384–386]. While it is seen in various calculus books (see [5, pages 88–90], where the series for  $\arcsin^3(x)$  is also proven), it dates back to at least two centuries and was given by Ramanujan among many others; see [6, pages 262–263]. As often in Mathematics, history is complicated. Equation (2.1) has been rediscovered repeatedly. For example, an equivalent form is elegantly solved as a 1962 MAA Monthly problem [7, Problem E 1509, Page 232]. We quote in extenso, the editors' attempts to trace the history of the formula:

*The series was located in the Smithsonian Mathematical Formulae and Tables of Elliptic Functions, 6.42 No. 5, p. 122; Chrystal, Algebra, vol. 2. 1906 ed, Ex. xx, No. 7, p. 335, (cites Pfaff as source); Bromwich, An Introduction to the Theory of Infinite Series, 1908 ed, Prob. 2, page 197 (claims known to Euler); Knopp, Theory and Application of Infinite Series, Ex. 123, Chap. VIII, p.271; Schuh, Leerboek der Differiaal en Integraalrekening, vol. 2, pp. 154–156; Hobson, Treatise on Plane Trigonometry, eqs. 20, 21, 22, pp. 279–280; M.R. Speigel, this Monthly, 60 (1953) 243–247; Taylor, Advanced Calculus, p. 632; Edwards, Differential Calculus for Beginners (1899), p.78.*

Note the appearance in the Monthly itself in 1953.

The second identity, slightly rewritten (see [6]), is less well known,

$$\arcsin^4\left(\frac{x}{2}\right) = \frac{3}{2} \sum_{k=1}^{\infty} \left\{ \sum_{m=1}^{k-1} \frac{1}{m^2} \right\} \frac{x^{2k}}{\binom{2k}{k} k^2}, \quad (2.2)$$

and when compared, they hint at the third and fourth identities below—subsequently confirmed numerically—the prior flimsy pattern and numerical exploration suggest that

$$\begin{aligned} \arcsin^6\left(\frac{x}{2}\right) &= \frac{45}{4} \sum_{k=1}^{\infty} \left\{ \sum_{m=1}^{k-1} \frac{1}{m^2} \sum_{n=1}^{m-1} \frac{1}{n^2} \right\} \frac{x^{2k}}{\binom{2k}{k} k^2}, \\ \arcsin^8\left(\frac{x}{2}\right) &= \frac{315}{2} \sum_{k=1}^{\infty} \left\{ \sum_{m=1}^{k-1} \frac{1}{m^2} \sum_{n=1}^{m-1} \frac{1}{n^2} \sum_{p=1}^{n-1} \frac{1}{p^2} \right\} \frac{x^{2k}}{\binom{2k}{k} k^2}. \end{aligned} \quad (2.3)$$

Reassured by this confirmation, we conjectured that in general

$$\frac{\arcsin^{2N}(x/2)}{(2N)!} = \sum_{k=1}^{\infty} \frac{H_N(k)}{\binom{2k}{k} k^2} x^{2k}, \quad (2.4)$$

where  $H_{N+1}(k)$  is as in (1.1). Subsequently, we were naturally led to discover the corresponding odd formulae.

We next provide a proof of both (1.1), equivalently (2.4), and (1.3).

*Proof of (1.1) and (1.3).* The formulae for  $\arcsin^k(x)$  with  $2 \leq k \leq 4$  are given in [6, pages 262–263], and Berndt comments that [5] is the best source he knows for  $k = 2$  and 3. Berndt’s proof also implicitly gives our desired result since he establishes, via a differential equation argument, that, for all real parameters  $a$ , one has

$$e^{a\arcsin(x)} = \sum_{n=0}^{\infty} c_n(a) \frac{x^n}{n!}, \quad (2.5)$$

where

$$c_{2n+1}(a) := a \prod_{k=1}^n (a^2 + (2k-1)^2), \quad c_{2n}(a) := \prod_{k=1}^n (a^2 + (2k-2)^2). \quad (2.6)$$

Now, expanding the power of  $a^n$  on each side of (2.5) provides the asserted formula. Note that (2.5) is equivalent to the somewhat-less-elegant if more-concise [8, Formula 10.49.33], specifically,

$$\sum_{k=0}^{\infty} \frac{(ia)_{k/2}}{k! (ia+1)_{-k/2}} (-ix)^k = \exp \left[ 2a \sin^{-1} \left( \frac{x}{2} \right) \right]. \quad (2.7)$$

Another proof can be obtained from the hypergeometric identity

$$\frac{\sin(ax)}{a \sin(x)} = {}_2F_1\left(\frac{1+a}{2}, \frac{1-a}{2}; \frac{3}{2}; \sin^2(x)\right) \quad (2.8)$$

given in [4, Exercise 16, page 189].  $\square$

*Maple* can also prove identities such as (2.5) as the following code shows:

```
> ce:=n->product(a^2+(2*k)^2,k=0..n-1):
> co:=n->a*product(a^2+(2*k+1)^2,k=0..n-1):
> sum(ce(n)*x^(2*n)/(2*n)!,n=0..infinity) assuming x>0;
          cosh(a arcsin(x))

> simplify(expand(sum(co(n)*x^(2*n+1)/(2*n+1)!,n=0..infinity)))
assuming x>0;
          sinh(a arcsin(x))
```

A necessarily equivalent formula for powers of  $\arcsin$  is listed by Hansen in [8, Formula 88.2.2],

$$\begin{aligned} & \sum_{n=0}^{\infty} x^{2n} \prod_{k=1}^{m-1} \left\{ \sum_{n_k=0}^{n_{k-1}} \frac{(2n_{k-1} - 2n_k)!}{[(n_{k-1} - n_k)!]^2 (2n_{k-1} - 2n_k + 1)} 2^{2n_k - 2n_{k-1}} \right\} \frac{(2n_{m-1})!}{(n_{m-1}!)^2 (2n_{m-1} + 1)} 2^{-2n_{m-1}} \\ &= \left( \frac{\sin^{-1} x}{x} \right)^m, \end{aligned} \quad (2.9)$$

but its relation to (1.1) and (1.3) is not obvious; our nested-sum representations are definitely more elegant. While Ramanujan listed only the first two cases in his notebooks [6, pages 262-263], his previous entries suggest that an approach for any power was being assembled. One can only imagine how much farther his intuition would have taken him if he had today's computational power!

Powers of  $\arcsin$  play an important role in analytical calculations of massive Feynman diagrams, see [9], and in the construction of Laurent expansions of different types of hypergeometric functions with respect to small parameters. In [9], series expansions for small powers of  $\arcsin$  are given, but they do not have the compact nested-sum form seen in (1.1) and (1.3). Likewise, formula (2.5) has proven central to recent work on effective asymptotic expansions for Laguerre polynomials and Bessel functions, [10]. These results are again motivated largely by applications in mathematical physics and in prime computation.

### 3. Properties of coefficient functions

It is of some independent interest, at least to the present authors, to determine a few properties of  $H_N(k)$  and  $G_N(k)$ .

**COROLLARY 3.1.** *The following properties obtain:*

- (a)  $H_N(k) = G_N(k) = 0$  if  $N > k$ ;

- (b)  $H_N(k) = \sum_{j=N-1}^{k-1} (H_{N-1}(j)/(2j)^2)$ ,  $G_N(k) = \sum_{j=N-1}^{k-1} (G_{N-1}(j)/(2j+1)^2)$ ;
- (c)  $H_k(k) = 1/4^k(k-1)!^2$ ,  $G_k(k) = 4^k k!^2/(2k)!^2$ ;
- (d)  $\sum_{N=1}^k (-4)^N H_N(k) = 0$  for  $k \geq 2$ ,  $\sum_{N=0}^k (-1)^N G_N(k) = 0$  for  $k \geq 1$ .

*Proof.* Parts (a)–(c) follow from the definition of  $H_N(k)$  and  $G_N(k)$ . To prove part (d), first note that (1.1) may be rewritten as

$$x^{2N} = \frac{(2N)!}{4^N} \sum_{k=1}^{\infty} \frac{4^{k+N} H_N(k)}{\binom{2k}{k} k^2} (\sin x)^{2k}. \quad (3.1)$$

Use this to obtain

$$\begin{aligned} 1 - 2 \sin^2 x &= \cos(2x) = 1 + \sum_{N=1}^{\infty} \frac{(-1)^N 4^N x^{2N}}{(2N)!} \\ &= 1 + \sum_{N=1}^{\infty} (-1)^N \sum_{k=1}^{\infty} \frac{4^{k+N} H_N(k)}{\binom{2k}{k} k^2} (\sin x)^{2k} \\ &= 1 + \sum_{k=1}^{\infty} \frac{4^k (\sin x)^{2k}}{\binom{2k}{k} k^2} \sum_{N=1}^k (-4)^N H_N(k). \end{aligned} \quad (3.2)$$

Now, matching powers of  $(\sin x)^2$  implies the first part of (d). Equation (1.3) may be similarly manipulated to give the second part of (d).  $\square$

Correspondingly, we have the following corollary.

**COROLLARY 3.2.** *For nonnegative integers  $N$  and  $k$ , one has*

$$\frac{4^{2k} H_N(k)}{\binom{2k}{k} k^2} = - \sum_{j=0}^{k-1} \frac{\binom{2j}{j} \binom{2k-2j}{k-j}}{2j-1} G_N(k-j), \quad (3.3)$$

$$\frac{\binom{2k-2}{k-1} G_N(k-1)}{2^{2k-1} (2k-1)} = - \sum_{j=0}^{k-1} \frac{\binom{2j}{j} 4^{k-2j}}{(k-j)(2j-1) \binom{2k-2j}{k-j}} H_{N+1}(k-j). \quad (3.4)$$

*Proof.* Differentiate (1.3) to obtain

$$\arcsin^{2N} \left( \frac{x}{2} \right) = \sqrt{1 - \left( \frac{x}{2} \right)^2} (2N)! \sum_{k=0}^{\infty} \frac{G_N(k) \binom{2k}{k}}{16^k} x^{2k}. \quad (3.5)$$

Using the binomial theorem and comparing to (1.1) gives (3.3). Similarly, differentiating (1.1) leads to (3.4).  $\square$

Even the simplest case, with  $N = 1$ , yields the nonobvious identity

$$\frac{4^{2k-1}}{\binom{2k}{k} k^2} = - \sum_{j=0}^{k-1} \frac{\binom{2j}{j} \binom{2k-2j}{k-j}}{2j-1} \sum_{n=0}^{k-j-1} \frac{1}{(2n+1)^2}. \quad (3.6)$$

Asking a student to prove this identity directly is instructive, particularly from an experimental mathematical perspective. While *Maple* stares dumbly at the right-hand side, it immediately redeems itself after one interchanges the order of summation, producing

$$\sum_{n=0}^{k-1} \frac{1}{(2n+1)^2} \sum_{j=0}^{k-n-1} \frac{\binom{2j}{j} \binom{2k-2j}{k-j}}{2j-1} = -\frac{\sqrt{\pi} 4^k \Gamma(k)}{4k \Gamma(k+1/2)}, \quad (3.7)$$

which may be easily rewritten as the desired expression (3.6).

A host of partition identities tumble directly from (1.3) and (2.4) on comparing various ways of combining powers of  $\arcsin$ .

**COROLLARY 3.3.** *Given  $m, n, \{M_i\}_{i=1}^m \geq 1$ , and  $\{N_i\}_{i=1}^n \geq 0$ , let*

$$\begin{aligned} T &:= 2M_1 + \cdots + 2M_m + (2N_1 + 1) + \cdots + (2N_n + 1), \\ W &:= 2^{-n} (2M_1)! \cdots (2M_m)! (2N_1 + 1)! \cdots (2N_n + 1)!. \end{aligned} \quad (3.8)$$

*Then, for any  $s \geq n$ , where  $s$  and  $n$  have the same parity,*

$$\begin{aligned} W \sum & \frac{\binom{2k_1}{k_1} \cdots \binom{2k_n}{k_n} H_{M_1}(j_1) \cdots H_{M_m}(j_m) G_{N_1}(k_1) \cdots G_{N_n}(k_n) 4^{2n}}{\binom{2j_1}{j_1} \cdots \binom{2j_m}{j_m} (j_1 \cdots j_m)^2 (2k_1 - 1) \cdots (2k_n - 1) 4^{2k_1 + \cdots + 2k_n}} \\ &= \begin{cases} 4T! \frac{H_{T/2}(s/2)}{\binom{s}{s/2} s^2}, & n \text{ even,} \\ \frac{T!}{2} \left( \frac{s-1}{s-1} \right) \frac{G_{(T-1)/2}((s-1)/2)}{s 4^{s-1}}, & n \text{ odd,} \end{cases} \end{aligned} \quad (3.9)$$

*where the sum is taken over all partitions of*

$$\frac{s+n}{2} = j_1 + \cdots + j_m + k_1 + \cdots + k_n, \quad (3.10)$$

*where  $j_1, j_2, \dots, j_m \geq 1$  and  $k_1, k_2, \dots, k_n \geq 1$ .*

An interesting special case occurs when we specify  $n = 0$ ,  $m = 2$ ,  $M_1 = M_2 = 1$ , and  $s = 2k$ :

$$\sum_{j=1}^{k-1} \frac{1}{\binom{2j}{j} \binom{2k-2j}{k-j} j^2 (k-j)^2} = \frac{6}{\binom{2k}{k} k^2} \sum_{j=1}^{k-1} \frac{1}{j^2}, \quad (3.11)$$

hence

$$\lim_{k \rightarrow \infty} \sum_{j=1}^{k-1} \frac{\binom{2k}{k}}{\binom{2j}{j} \binom{2k-2j}{k-j}} \frac{k^2}{j^2 (k-j)^2} = \pi^2. \quad (3.12)$$

#### 4. Powers of arcsin via iterated integrals

Since  $\arcsin'(x) = 1/\sqrt{1-x^2}$ , we may also express any power of  $\arcsin(x)$  as an iterated integral. Specifically,

$$\frac{\arcsin^n x}{n!} = \int_0^x \frac{dy_1}{\sqrt{1-y_1^2}} \int_0^{y_1} \frac{dy_2}{\sqrt{1-y_2^2}} \int_0^{y_2} \frac{dy_3}{\sqrt{1-y_3^2}} \cdots \int_0^{y_{n-1}} \frac{dy_n}{\sqrt{1-y_n^2}}. \quad (4.1)$$

To convert the multiple integral into a multiple sum, note that the binomial theorem gives

$$\frac{1}{\sqrt{1-x^2}} = \sum_{k=0}^{\infty} \frac{1}{4^k} \binom{2k}{k} x^{2k}, \quad (4.2)$$

which may be repeatedly used in (4.1) to yield

$$\begin{aligned} \frac{\arcsin^n x}{n!} &= \sum_{k_1, k_2, \dots, k_n=0}^{\infty} \frac{\binom{2k_1}{k_1} \binom{2k_2}{k_2} \cdots \binom{2k_n}{k_n} x^{2w_n+n}}{4^{w_n} (2k_1+1)(2k_1+2k_2+2) \cdots (2w_n+n)} \\ &= x^n \sum_{k=0}^{\infty} \left\{ \sum_{k_1=0}^k \frac{\binom{2k_1}{k_1}}{4^{k_1} (2k_1+1)} \sum_{k_2=0}^{k-k_1} \frac{\binom{2k_2}{k_2}}{4^{k_2} (2k_1+2k_2+2)} \cdots \sum_{k_n=0}^{k+k_{n-1}-w_n} \frac{\binom{2k_n}{k_n}}{4^{k_n} (2w_n+n)} \right\} x^k \\ &= x^n \sum_{k=0}^{\infty} \left\{ \sum_{k_1=0}^k \frac{\binom{2k_1}{k_1}}{(2k_1+1)} \sum_{k_2=k_1}^k \frac{\binom{2k_2-2k_1}{k_2-k_1}}{(2k_2+2)} \cdots \sum_{k_n=k_{n-1}}^k \frac{\binom{2k_n-2k_{n-1}}{k_n-k_{n-1}}}{(2k_n+n)} \frac{1}{4^{k_n}} \right\} x^k, \end{aligned} \quad (4.3)$$

where  $w_n := k_1 + k_2 + \cdots + k_n$ .

Though this process also writes the coefficients in terms of nested sums, these terms are not nearly as simple as  $H_N$  and  $G_N$ .

#### 5. Related series manipulations

After having discovered formulae such as (1.1) and (1.3), the analyst's natural inclination is to "mine" them for striking examples. We rescale (1.1) and (1.3) to obtain

$$\begin{aligned} \arcsin^{2N}(x) &= (2N)! \sum_{k=1}^{\infty} \frac{H_N(k) 4^k}{\binom{2k}{k} k^2} x^{2k}, \\ \arcsin^{2N+1}(x) &= (2N+1)! \sum_{k=0}^{\infty} \frac{G_N(k) \binom{2k}{k}}{(2k+1) 4^k} x^{2k+1}. \end{aligned} \quad (5.1)$$

These series, or their derivatives, may be evaluated at values such as  $x = \pi/2, \pi/3, \pi/4, \pi/6, i/2$ , and  $i$  to obtain many formulae along the lines of those found by Lehmer and

others as described in [4, pages 384–386]. A few examples are

$$\begin{aligned}
 \sum_{k=1}^{\infty} \frac{H_N(k)4^k}{\binom{2k}{k}k^2} &= \frac{1}{(2N)!} \left(\frac{\pi}{2}\right)^{2N}, & \sum_{k=1}^{\infty} \frac{H_N(k)3^k}{\binom{2k}{k}k^2} &= \frac{1}{(2N)!} \left(\frac{\pi}{3}\right)^{2N}, \\
 \sum_{k=1}^{\infty} \frac{H_N(k)2^k}{\binom{2k}{k}k^2} &= \frac{1}{(2N)!} \left(\frac{\pi}{4}\right)^{2N}, & \sum_{k=1}^{\infty} \frac{H_N(k)}{\binom{2k}{k}k^2} &= \frac{1}{(2N)!} \left(\frac{\pi}{6}\right)^{2N}, \\
 \sum_{k=1}^{\infty} \frac{H_N(k)(-1)^k}{\binom{2k}{k}k^2} &= \frac{(-1)^N}{(2N)!} \left(\log \frac{\sqrt{5}-1}{2}\right)^{2N}, \\
 \sum_{k=1}^{\infty} \frac{H_N(k)(-4)^k}{\binom{2k}{k}k^2} &= \frac{(-1)^N}{(2N)!} (\log(1+\sqrt{2}))^{2N}.
 \end{aligned} \tag{5.2}$$

Integrating (1.1) and (1.3) is, naturally, much more challenging. Replacing  $x$  by  $ix$  in (1.1) and (1.3) provides the Maclaurin series for positive integer powers of the form  $\log^N(x + \sqrt{x^2 + 1})$ .

These expressions may then be integrated (or differentiated). In particular, see [4, Exercise 17, page 189], we have

$$\frac{1}{4} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3 \binom{2n}{n}} = \int_0^{1/2} \frac{\log^2(x + \sqrt{x^2 + 1})}{x} dx = \frac{\zeta(3)}{10}. \tag{5.3}$$

Likewise, with more work, see [11], we obtain

$$\begin{aligned}
 \int_0^{1/2} \frac{\ln^4(x + \sqrt{x^2 + 1})}{x} dx &= -\frac{3}{2} \text{Li}_5(g^2) + 3\text{Li}_4(g^2) \ln(g) + \frac{3}{2} \zeta(5) \\
 &\quad - \frac{12}{5} \zeta(3) \ln^2(g) - \frac{4}{15} \pi^2 \ln^3(g) + \frac{4}{5} \ln^5(g),
 \end{aligned} \tag{5.4}$$

where  $g = (\sqrt{5} - 1)/2$  is the golden ratio and where  $\text{Li}_n(z) = \sum_{k=1}^{\infty} z^k/k^n$  is the *polylogarithm* of order  $n$ .

This process may be generalized as follows. Define

$$\mathcal{L}_n := \frac{2^n}{n!} \int_0^{1/2} \frac{\ln^n(x + \sqrt{x^2 + 1})}{x} dx. \tag{5.5}$$

We leave it to the reader to determine the explicit series expression for  $\mathcal{L}_n$ , for  $n$  is even and for  $n$  is odd.

Extensive experimentation with *Maple* coupled with pattern lookups with Sloane's *online encyclopedia* [12] produced

$$\mathcal{L}_n = \zeta(n+1) - \frac{n(-\log g^2)^{n+1}}{2(n+1)!} - \sum_{j=2}^{n+1} \frac{(-\log g^2)^{n+1-j}}{(n+1-j)!} \text{Li}_j(g^2). \tag{5.6}$$

Multiplying by  $x^n$ , and summing over  $n \geq 1$ , shows that this is equivalent to the next generating function, written in terms of  $\psi$ , the *digamma* function:

$$\begin{aligned} -x \int_0^{-\log g} (e^{2xy} - 1) \coth(y) dy &= \frac{1}{2} + x(\gamma + \psi(1-x)) - \frac{1/2 + x \log g}{e^{2x \log g}} \\ &\quad + e^{-2x \log g} \sum_{k=2}^{\infty} \text{Li}_k(g^2) x^k. \end{aligned} \tag{5.7}$$

These observations can be made rigorous with the next result.

**THEOREM 5.1.** *For  $n \geq 1$  and  $|x| \leq 1$ ,*

$$\begin{aligned} \frac{1}{n!} \int_0^x \frac{\arcsin^n(y)}{y} dy &= - \sum_{k=2}^{n+1} \text{Li}_k((\sqrt{1-x^2} + ix)^2) \frac{(-2i)^{1-k} \arcsin(x)^{n+1-k}}{(n+1-k)!} \\ &\quad - \frac{i \arcsin^{n+1}(x)}{(n+1)!} + \frac{\arcsin^n(x)}{n!} \log(2x^2 - 2ix\sqrt{1-x^2}) + \left(\frac{i}{2}\right)^n \zeta(n+1). \end{aligned} \tag{5.8}$$

*Proof.* The derivatives of each side match, and the equation holds for  $x = 0$ . Alternatively, this is a reworking of [13, Formula (7.48), page 199].  $\square$

In hindsight, this equation should not come as much of a surprise since Ramanujan's entries immediately preceding (2.1) give similar formulae using the Clausen functions. One can substitute (1.1) or (1.3) to obtain further formulae. This allows for extensions to the complex plane. For example, we have the following corollary.

**COROLLARY 5.2.** *For  $n \geq 1$ ,  $|x| \leq 1$ , and  $\text{Re}(x) \geq 0$ ,*

$$\begin{aligned} \frac{(-1)^n}{2} \sum_{k=1}^{\infty} \frac{H_n(k)(-4)^k}{\binom{2k}{k} k^3} x^{2k} &= - \sum_{k=2}^{2n+1} \text{Li}_k((x - \sqrt{1+x^2})^2) \frac{2^{1-k} \text{arcsinh}(x)^{2n+1-k}}{(2n+1-k)!} \\ &\quad + \frac{\text{arcsinh}^{2n+1}(x)}{(2n+1)!} + \frac{\text{arcsinh}^{2n}(x)}{2n!} \log(2x\sqrt{1+x^2} - 2x^2) \\ &\quad + \frac{\zeta(2n+1)}{4^n}. \end{aligned} \tag{5.9}$$

*Example 5.3.* Substituting  $x = i$  gives

$$\begin{aligned} \frac{1}{2} \sum_{k=1}^{\infty} \frac{H_n(k)4^k}{\binom{2k}{k} k^3} &= \sum_{k=1}^n (1 - 2^{-2k}) \zeta(2k+1) \left(-\frac{1}{4}\right)^k \frac{(\pi/2)^{2n-2k}}{(2n-2k)!} \\ &\quad + \frac{(\pi/2)^{2n}}{(2n)!} \log 2 + \left(-\frac{1}{4}\right)^n \zeta(2n+1), \end{aligned} \tag{5.10}$$

with the special case  $n = 1$  yielding

$$\sum_{k=1}^{\infty} \frac{4^k}{\binom{2k}{k} k^3} = \pi^2 \log 2 - \frac{7}{2} \zeta(3). \quad (5.11)$$

## 6. Conclusion

We hope this examination of (1.1) and (1.3) encourages readers to similarly explore what transpires for  $\arctan$  and other functions. Armed with a good computer algebra system and an internet connection, one can quite fearlessly undertake this task.

## Acknowledgment

Research is supported by NSERC and by the Canada Research Chair Programme.

## References

- [1] J. M. Borwein and D. Bailey, *Mathematics by Experiment*, A K Peters, Natick, Mass, USA, 2004.
- [2] J. M. Borwein, D. Bailey, and R. Girgensohn, *Experimentation in Mathematics*, A K Peters, Natick, Mass, USA, 2004.
- [3] E. W. Weisstein, “Inverse Sine,” from MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/InverseSine.html>.
- [4] J. M. Borwein and P. B. Borwein, *Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity*, Canadian Mathematical Society Series of Monographs and Advanced Texts, John Wiley & Sons, New York, NY, USA, 1987.
- [5] J. Edwards, *Differential Calculus*, MacMillan, London, UK, 2nd edition, 1982.
- [6] B. C. Berndt, *Ramanujan’s Notebooks—Part I*, Springer, New York, NY, USA, 1985.
- [7] A. G. Konheim, J. W. Wrench Jr., and M. S. Klamkin, “A Well-Known Series,” *American Mathematical Monthly*, vol. 69, no. 10, p. 1011, December 1962.
- [8] E. R. Hansen, *A Table of Series and Products*, Prentice-Hall, Englewood Cliffs, NJ, USA, 1975.
- [9] M. Yu. Kalmykov and A. Sheplyakov, “Isjk—a C++ library for arbitrary-precision numeric evaluation of the generalized log-sine functions,” *Computer Physics Communications*, vol. 172, no. 1, pp. 45–59, 2005.
- [10] D. Borwein, J. M. Borwein, and R. E. Crandall, “Effective laguerre asymptotics,” submitted to *SIAM Journal on Numerical Analysis*.
- [11] J. M. Borwein, D. J. Broadhurst, and J. Kamnitzer, “Central binomial sums, multiple Clausen values, and zeta values,” *Experimental Mathematics*, vol. 10, no. 1, pp. 25–34, 2001.
- [12] N. J. Sloane, “Online Encyclopedia of Integer Sequences,” (<http://www.research.att.com/~njas/sequences/>).
- [13] L. Lewin, *Polylogarithms and Associated Functions*, North-Holland, New York, NY, USA, 1981.

Jonathan M. Borwein: Faculty of Computer Science, Dalhousie University, Halifax, NS, Canada B3H2W5  
*Email address:* jborwein@cs.dal.ca

Marc Chamberland: Department of Mathematics and Statistics, Grinnell College, Grinnell, IA 50112, USA  
*Email address:* chamberl@math.grinnell.edu

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)