

*Research Article*  
**On Generalized Periodic-Like Rings**

Howard E. Bell and Adil Yaqub

Received 21 November 2006; Accepted 27 February 2007

Recommended by Francois Goichot

Let  $R$  be a ring with center  $Z$ , Jacobson radical  $J$ , and set  $N$  of all nilpotent elements. Call  $R$  generalized periodic-like if for all  $x \in R \setminus (N \cup J \cup Z)$  there exist positive integers  $m, n$  of opposite parity for which  $x^m - x^n \in N \cap Z$ . We identify some basic properties of such rings and prove some results on commutativity.

Copyright © 2007 H. E. Bell and A. Yaqub. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Let  $R$  be a ring; and let  $N = N(R)$ ,  $Z = Z(R)$  and  $J = J(R)$  denote, respectively, the set of nilpotent elements, the center, and the Jacobson radical. As usual, we call  $R$  periodic if for each  $x \in R$ , there exist distinct positive integers  $m, n$  such that  $x^m = x^n$ . In [1] we defined  $R$  to be generalized periodic (g-p) if for each  $x \in R \setminus (N \cup Z)$ ,

there exist positive integers  $m, n$  of opposite parity such that  $x^m - x^n \in N \cap Z$ . (\*)

We now define  $R$  to be generalized periodic-like (g-p-l) if (\*) holds for each  $x \in R \setminus (N \cup J \cup Z)$ . Clearly, the class of g-p-l rings contains all commutative rings, all nil rings, all Jacobson radical rings, all g-p rings, and some (but not all) periodic rings. It is our purpose to exhibit some general properties of g-p-l rings and to study commutativity of such rings.

## 2. Preliminary results

To simplify our discussion, we denote by  $((m, n))$  the ordered pair of integers  $m, n$  of opposite parity. The rest of our notation and terminology is standard. For elements  $x, y \in R$ , the symbol  $[x, y]$  denotes the commutator  $xy - yx$ ; for subsets  $X, Y \subseteq R$ ,  $[X, Y]$  denotes

the set  $\{[x, y] \mid x \in X, y \in Y\}$ ; and  $C(R)$  denotes the commutator ideal of  $R$ . An element  $x \in R$  is called regular if it is not a zero divisor; it is called periodic if there exist distinct positive integers  $m, n$  for which  $x^m = x^n$ ; and it is called potent if there exists an integer  $n > 1$  for which  $x^n = x$ . The set of all potent elements of  $R$  is denoted by  $P$  or  $P(R)$ , and the prime radical by  $\mathfrak{P}(R)$ . Finally,  $R$  is called reduced if  $N(R) = \{0\}$ .

LEMMA 2.1. *Let  $R$  be an arbitrary g-p-l ring.*

- (i) *Every epimorphic image of  $R$  is a g-p-l ring.*
- (ii)  $N \subseteq J$ .
- (iii) *If  $[N, J] = \{0\}$ , then  $N$  is an ideal.*
- (iv)  $C(R) \subseteq J$ .
- (v) *If  $e$  is an idempotent, the additive order of which is not a power of 2, then  $e \in Z$ .*

*Proof.* (i) is clear, once we recall that if  $\sigma : R \rightarrow S$  is an epimorphism, then  $\sigma(J(R)) \subseteq J(S)$ .

(ii) Let  $S = R/J(R)$ . Then by (i),  $S$  is a g-p-l ring; and since  $J(S) = \{0\}$ ,  $S$  is a g-p ring. It follows from [1, Theorem 1] that  $N(S)$  is an ideal of  $S$ , hence  $N(S) \subseteq J(S) = \{0\}$  and therefore  $N(R) \subseteq J(R)$ .

(iii) Since  $N \subseteq J$ ,  $N$  is commutative and hence  $(N, +)$  is an additive subgroup. Let  $a \in N$  and  $x \in R$ . Then  $ax \in J$ , so  $[a, ax] = 0$ , that is,  $a^2x = axa$ . It follows that  $(ax)^2 = a^2x^2$  and that  $(ax)^n = a^n x^n$  for all positive integers  $n$ . Therefore,  $ax \in N$ .

(iv) As in (ii),  $R/J(R)$  is a g-p ring; hence, by [1, Lemma 2],  $C(R/J(R)) = \{0\}$ . Therefore,  $C(R) \subseteq J(R)$ .

(v) If  $e \notin Z$ , then  $-e \notin J \cup Z$  and there exists  $((m, n))$  such that  $(-e)^m - (-e)^n \in N \cap Z$ . Since  $m, n$  are of opposite parity, we get  $2e \in N$ , so that  $2^k e = 0$  for some  $k$ .  $\square$

LEMMA 2.2. *Let  $R$  be an arbitrary g-p-l ring, and let  $x \in R$ . Then either  $x \in J \cup Z$ , or there exist a positive integer  $q$  and an idempotent  $e$  such that  $x^q = x^q e$ .*

*Proof.* If  $x \notin J \cup Z$ , there exists  $((m, n))$  such that  $x^m - x^n \in N \cap Z$ . Therefore, there exist a positive integer  $q$  and  $g(t) \in \mathbb{Z}[t]$  such that  $x^q = x^{q+1}g(x)$ . It is now easy to verify that  $e = (xg(x))^q$  is an idempotent with  $x^q = x^q e$ .  $\square$

LEMMA 2.3. *Let  $R$  be a g-p-l ring and  $\sigma$  an epimorphism from  $R$  to  $S$ . Then  $N(S) \subseteq \sigma(J(R)) \cup Z(S)$ .*

*Proof.* Let  $s \in N(S)$  with  $s^k = 0$  and let  $d \in R$  such that  $\sigma(d) = s$ . If  $d \in J(R) \cup Z(R)$ , then obviously  $s \in \sigma(J(R)) \cup Z(S)$ ; hence we may suppose that there exists  $((m, n))$  with  $n > m$  such that  $d^m - d^n \in N(R) \cap Z(R)$ . It is easy to show that  $d - d^h \in N$ , where  $h = n - m + 1$ ; thus

$$d - d^{k+1}d^{k(h-2)} = d - d^h + d^{h-1}(d - d^h) + \dots + (d^{h-1})^{k-1}(d - d^h) \quad (2.1)$$

is a sum of commuting nilpotent elements, hence it is in  $N(R)$  and therefore in  $J(R)$ . Consequently,  $s - s^{k+1}s^{k(h-2)} \in \sigma(J(R))$ ; and since  $s^{k+1} = 0$ ,  $s \in \sigma(J(R))$ .  $\square$

We finish this section by stating two known results on periodic elements.

LEMMA 2.4. Let  $R$  be an arbitrary ring, and let  $N^* = \{x \in R \mid x^2 = 0\}$ .

- (i) [2, Lemma 1] If  $x \in R$  is periodic, then  $x \in P + N$ .
- (ii) [3, Theorem 2] If  $N^*$  is commutative and  $N$  is multiplicatively closed, then  $PN \subseteq N$ .

### 3. Commutativity results

THEOREM 3.1. If  $R$  is a g-p-l ring with  $J \subseteq Z$ , then  $R$  is commutative.

*Proof.* Suppose  $x \notin Z$ . Then by Lemma 2.1(ii), we have  $((m, n))$  with  $n > m$  such that  $x^m - x^n \in N \cap Z$ . Consequently  $x^{n-m+1} - x \in N$ ; and since  $N \subseteq Z$ , commutativity of  $R$  follows by a well-known theorem of Herstein [4].  $\square$

THEOREM 3.2. If  $R$  is any g-p-l ring with 1, then  $R$  is commutative.

*Proof.* We show that if  $R$  is g-p-l with 1, then  $J \subseteq Z$ . Suppose that  $x \in J \setminus Z$ . Then  $-1 + x \notin J \cup Z$ , so there exists  $((m, n))$  such that  $(-1 + x)^m - (-1 + x)^n \in N \cap Z$ ; and we may assume that  $m$  is even and  $n$  is odd. Since  $N \subseteq J$ , it follows that  $2 \in J$ ; thus for every integer  $m, 2m \in J$ , and hence  $2m + 1$  is invertible.

Now consider  $((m_1, n_1))$  such that  $(1 + x)^{m_1} - (1 + x)^{n_1} \in N \cap Z$ . Then  $(m_1 - n_1)x + x^2p(x) \in N \cap Z$  for some  $p(t) \in \mathbb{Z}[t]$ ; and since  $m_1 - n_1$  is central and invertible, we get  $x + x^2w$  in  $N \cap Z$  for some  $w$  in  $R$  with  $[x, w] = 0$ . Thus, we have a positive integer  $q$  and an element  $y$  in  $R$  such that  $[x, y] = 0$  and  $x^q = x^{q+1}y$ . It follows that  $e = (xy)^q$  is an idempotent such that  $x^q = x^q e$ ; and since  $J$  contains no nonzero idempotents,  $x$  is in  $N$ .

Let  $\alpha$  be the smallest positive integer for which  $x^k \in Z$  for all  $k \geq \alpha$ , and note that, since  $x \notin Z$ ,  $\alpha \geq 2$ . But  $1 + x^{\alpha-1} \notin J \cup Z$ , so there exists  $((m_2, n_2))$  such that  $(1 + x^{\alpha-1})^{m_2} - (1 + x^{\alpha-1})^{n_2} \in N \cap Z$ ; hence  $(m_2 - n_2)x^{\alpha-1} \in Z$ . But since  $m_2 - n_2$  is invertible and central, we conclude that  $x^{\alpha-1} \in Z$ —a contradiction.  $\square$

THEOREM 3.3. If  $R$  is a reduced g-p-l ring with  $R \neq J$ , then  $R$  is commutative.

*Proof.* If  $R = J \cup Z$ , then  $R = Z$  and we are finished. Otherwise, if  $x \in R \setminus (J \cup Z)$ , there exists  $((m, n))$  such that  $x^m - x^n \in N \cap Z = \{0\}$ ; hence  $x$  is periodic, and by Lemma 2.4(i),  $x \in P$ . Thus,  $R = P \cup J \cup Z$ ; and to complete the proof we need only to show that  $P \subseteq Z$ .

Let  $y \in P$ , and let  $k > 1$  be such that  $y^k = y$ . Then  $e = y^{k-1}$  is an idempotent for which  $y = ye$ , and  $e \in Z$  since  $N = \{0\}$ . Now  $eR$  is an ideal of  $R$ , so that  $J(eR) = eR \cap J(R)$ ; hence  $eR$  is a g-p-l ring with 1, which is commutative by Theorem 3.2. Therefore,  $[ey, ew] = 0$  for all  $w \in R$ ; and since  $ey = y$  and  $e \in Z$ , we conclude that  $[y, w] = 0$  for all  $w \in R$ , that is,  $y \in Z$ .  $\square$

THEOREM 3.4. If  $R$  is a g-p-l ring in which  $J$  is commutative and all idempotents are central, then  $R$  is commutative.

*Proof.* We may express  $R$  as a subdirect product of subdirectly irreducible rings, each of which is an epimorphic image of  $R$ . Let  $R_\alpha$  be such a subdirectly irreducible ring, and let  $\sigma : R \rightarrow R_\alpha$  be an epimorphism. Let  $x_\alpha \in R_\alpha$  and let  $x \in R$  such that  $\sigma(x) = x_\alpha$ . By Lemma 2.2,  $x \in J(R) \cup Z(R)$  or there exist an idempotent  $e \in R$  and a positive integer  $q$  such that  $x^q = x^q e$ . Thus, either  $x_\alpha \in \sigma(J(R)) \cup \sigma(Z(R))$  or  $x_\alpha^q = x_\alpha^q e_\alpha$ , where  $e_\alpha = \sigma(e)$  is a central idempotent of  $R_\alpha$ . But  $R_\alpha$  is subdirectly irreducible, hence if  $R_\alpha$  has a nonzero central idempotent, then  $R_\alpha$  has 1 and is commutative by Theorem 3.2.

To complete the proof, we need only consider the case that for each  $x_\alpha \in R_\alpha$ ,  $x_\alpha \in \sigma(J(R)) \cup Z(R_\alpha) \cup N(R_\alpha)$ . Now by Lemma 2.3,  $N(R_\alpha) \subseteq \sigma(J(R)) \cup Z(R_\alpha)$ ; hence  $R_\alpha = \sigma(J(R)) \cup Z(R_\alpha)$ , which is clearly commutative. Therefore,  $R$  is commutative.  $\square$

Theorem 3.4 has two corollaries, the first of which is immediate when we recall Lemma 2.1(v).

**COROLLARY 3.5.** *If  $R$  is a 2-torsion-free g-p-l ring with  $J$  commutative, then  $R$  is commutative.*

**COROLLARY 3.6.** *Let  $R$  be a g-p-l ring containing a regular central element  $c$ . If  $J$  is commutative, then  $R$  is commutative.*

*Proof.* It suffices to show that  $N \subseteq Z$  since this condition implies that idempotents are central. Consider first the case  $c \in J$ . Then  $cJ \subseteq J^2$ , which is central since  $J$  is commutative. Since  $c$  is regular and central, it is immediate that  $J \subseteq Z$ , so certainly  $N \subseteq Z$ .

Now assume that  $c \notin J$ , and suppose that  $a \in N \setminus Z$ . Then  $c + a \notin J \cup Z$ , and there exists  $((m, n))$  such that  $(c + a)^m - (c + a)^n \in N \cap Z$ . It follows that  $c^m - c^n$  is a sum of commuting nilpotent elements, hence  $c^m - c^n \in N$  and there exists  $q$  such that  $c^q = c^{q+1}p(c)$  for some  $p(t) \in \mathbb{Z}[t]$ . As before, we get an idempotent  $e$  such that  $c^q = c^q e$  and  $[c, e] = 0$ . Now  $e$  cannot be a zero divisor, since that would force  $c$  to be a zero divisor; therefore,  $R$  has a regular idempotent, that is,  $R$  has 1. We have contradicted Theorem 3.2, so  $N \subseteq Z$  as claimed.  $\square$

#### 4. Nil-commutator-ideal theorems

**THEOREM 4.1.** *Let  $R$  be a g-p-l ring. If  $R \neq J$  and  $N$  is an ideal, then  $C(R)$  is nil.*

*Proof.* We may assume  $R \neq J \cup Z$ , since otherwise  $R$  is commutative. Let  $\bar{R} = R/N$ , and let the element  $x + N$  of  $\bar{R}$  be denoted by  $\bar{x}$ . We need to show that  $\bar{R}$  is commutative—a conclusion that follows from Theorem 3.3 once we show that  $J(\bar{R}) \neq \bar{R}$ .

Suppose that  $J(\bar{R}) = \bar{R}$ , and let  $x \in R \setminus (J \cup Z)$ . By Lemma 2.2, there exists a positive integer  $q$  and an idempotent  $e \in R$  such that  $x^q = x^q e$ ; and it follows that  $\bar{e}$  is an idempotent of  $\bar{R}$  such that  $\bar{x}^q = \bar{x}^q \bar{e}$ . But  $\bar{R} = J(\bar{R})$  contains no nonzero idempotents, so that  $\bar{x}^q = 0 = \bar{x}$  and hence  $x \in N(R)$ . This contradicts the fact that  $x \notin J \cup Z$ , hence  $\bar{R} \neq J(\bar{R})$  as required.  $\square$

**THEOREM 4.2.** *If  $R$  is a g-p-l ring and  $J$  is commutative, then  $C(R)$  is nil.*

*Proof.* If  $R = J$ , then  $R$  is commutative. If  $R \neq J$ ,  $N$  is an ideal by Lemma 2.1(iii) and  $C(R)$  is nil by Theorem 4.1.  $\square$

In fact, we can improve this result as follows.

**THEOREM 4.3.** *Let  $R$  be a g-p-l ring with  $R \neq J$ . If  $N$  is commutative, then  $C(R)$  is nil.*

This result follows from Theorem 4.1, once we prove our final theorem.

**THEOREM 4.4.** *Let  $R$  be a g-p-l ring with  $R \neq J$ . If  $N$  is commutative, then  $N$  is an ideal.*

*Proof.* Again we may assume that  $R \neq J \cup Z$ . Since  $N$  is commutative,  $N$  is an additive subgroup of  $R$ . To show that  $RN \subseteq N$ , it is convenient to work with the ring  $\bar{R} = R/\mathfrak{P}(R)$ . As in the proof of Theorem 4.1, we have  $J(\bar{R}) \neq \bar{R}$ ; and if  $\bar{R} = Z(\bar{R})$ , then  $C(R) \subseteq \mathfrak{P}(R) \subseteq N$ . Therefore, we assume that  $\bar{R} \neq J(\bar{R}) \cup Z(\bar{R})$ . We note that if  $x + N = \bar{x} \in N(\bar{R})$ , then  $x \in N(R)$ ; consequently,  $N(\bar{R})$  is commutative and hence is an additive subgroup of  $\bar{R}$ .

Now  $\bar{R}$  is semiprime and therefore  $N(\bar{R}) \cap Z(\bar{R}) = \{0\}$ . It follows that if  $\bar{x} \in \bar{R} \setminus (J(\bar{R}) \cup Z(\bar{R}))$ , there exists  $((m, n))$  such that  $\bar{x}^m = \bar{x}^n$ , that is,  $\bar{x}$  is periodic. Thus  $\bar{x} \in P(\bar{R}) + N(\bar{R})$  by Lemma 2.4(i); and by commutativity of  $N(\bar{R})$  and Lemma 2.4(ii) we get  $\bar{x}N(\bar{R}) \subseteq N(\bar{R})$ . Moreover, if  $\bar{y} \in Z(\bar{R})$ ,  $\bar{y}N(\bar{R}) \subseteq N(\bar{R})$ . Now let  $\bar{y} \in J(\bar{R}) \setminus Z(\bar{R})$ , and let  $\bar{x} \in \bar{R} \setminus (J(\bar{R}) \cup Z(\bar{R}))$ . Then  $\bar{x} + \bar{y} \notin J(\bar{R})$ , hence it is in  $\bar{R} \setminus (J(\bar{R}) \cup Z(\bar{R}))$  or in  $Z(\bar{R})$ ; and in either case  $(\bar{x} + \bar{y})N(\bar{R})$  and  $\bar{x}N(\bar{R})$  are in  $N(\bar{R})$ , so that  $\bar{y}N(\bar{R}) \subseteq N(\bar{R})$ . We have shown that  $N(\bar{R})$  is an ideal of  $\bar{R}$ ; therefore, if  $x \in R$  and  $a \in N(R)$ ,  $\bar{x}a \in N(\bar{R})$  and hence  $xa \in N(R)$ . Thus,  $N(R)$  is an ideal of  $R$ .  $\square$

*Remark 4.5.* There exist noncommutative g–p–l rings with  $J$  commutative. An accessible example is

$$\left\{ \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix} \mid a, b \in GF(2) \right\}. \quad (4.1)$$

### Acknowledgment

The first author was supported by the Natural Sciences and Engineering Research Council of Canada, Grant 3961.

### References

- [1] H. E. Bell and A. Yaqub, “Generalized periodic rings,” *International Journal of Mathematics and Mathematical Sciences*, vol. 19, no. 1, pp. 87–92, 1996.
- [2] H. E. Bell, “A commutativity study for periodic rings,” *Pacific Journal of Mathematics*, vol. 70, no. 1, pp. 29–36, 1977.
- [3] H. E. Bell and H. Tominaga, “On periodic rings and related rings,” *Mathematical Journal of Okayama University*, vol. 28, pp. 101–103, 1986.
- [4] I. N. Herstein, “A generalization of a theorem of Jacobson III,” *American Journal of Mathematics*, vol. 75, no. 1, pp. 105–111, 1953.

Howard E. Bell: Department of Mathematics, Brock University, St. Catharines, Ontario, Canada L2S 3A1  
*Email address:* hbell@brocku.ca

Adil Yaqub: Department of Mathematics, University of California, Santa Barbara, CA 93106, USA  
*Email address:* yaqub@math.ucsb.edu

## Special Issue on Intelligent Computational Methods for Financial Engineering

### Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Lean Yu**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [yulean@amss.ac.cn](mailto:yulean@amss.ac.cn)

**Shouyang Wang**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; [sywang@amss.ac.cn](mailto:sywang@amss.ac.cn)

**K. K. Lai**, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [mskklai@cityu.edu.hk](mailto:mskklai@cityu.edu.hk)