

Research Article

Certain Coefficient Bounds for p -Valent Functions

C. Ramachandran, S. Sivasubramanian, and H. Silverman

Received 1 October 2006; Revised 18 November 2006; Accepted 18 January 2007

Recommended by Brigitte Forster-Heinlein

In the present paper, the authors obtain sharp bounds for certain subclasses of p -valent functions. The results are extended to functions defined by convolution.

Copyright © 2007 Hindawi Publishing Corporation. All rights reserved.

1. Introduction

Let \mathcal{A}_p denote the class of all analytic functions $f(z)$ of the form

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n \quad (1.1)$$

defined on the *open* unit disk

$$\Delta = \{z : z \in \mathbb{C} : |z| < 1\}, \quad (1.2)$$

and let $\mathcal{A}_1 := \mathcal{A}$. For $f(z)$ given by (1.1) and $g(z)$ given by

$$g(z) = z^p + \sum_{n=p+1}^{\infty} b_n z^n, \quad (1.3)$$

their convolution (or Hadamard product), denoted by $(f * g)$, is defined as

$$(f * g)(z) = z^p + \sum_{n=p+1}^{\infty} a_n b_n z^n. \quad (1.4)$$

With a view to recalling the principle of subordination between analytic functions, let the functions f and g be analytic in Δ . Then we say that the function f is *subordinate* to g if

there exists a Schwarz function $\omega(z)$, analytic in Δ with

$$\omega(0) = 0, \quad |\omega(z)| < 1 \quad (z \in \Delta), \quad (1.5)$$

such that

$$f(z) = g(\omega(z)) \quad (z \in \Delta). \quad (1.6)$$

We denote this subordination by

$$f \prec g \quad \text{or} \quad f(z) \prec g(z) \quad (z \in \Delta). \quad (1.7)$$

In particular, if the function g is univalent in Δ , the above subordination is equivalent to

$$f(0) = g(0), \quad f(\Delta) \subset g(\Delta). \quad (1.8)$$

Let $\phi(z)$ be an analytic function with positive real part on Δ with $\phi(0) = 1$, $\phi'(0) > 0$ which maps the open unit disk Δ onto a region starlike with respect to 1 and is symmetric with respect to the real axis. Ali et al. [1] defined and studied the class $S_{b,p}^*(\phi)$ to be the class of functions in $f \in \mathcal{A}_p$ for which

$$1 + \frac{1}{b} \left(\frac{1}{p} \frac{zf'(z)}{f(z)} - 1 \right) \prec \phi(z) \quad (z \in \Delta, b \in \mathbb{C} \setminus \{0\}), \quad (1.9)$$

and the class $C_{b,p}(\phi)$ of all functions in $f \in \mathcal{A}_p$ for which

$$1 - \frac{1}{b} + \frac{1}{bp} \left(1 + \frac{zf''(z)}{f'(z)} \right) \prec \phi(z) \quad (z \in \Delta, b \in \mathbb{C} \setminus \{0\}). \quad (1.10)$$

Ali et al. [1] also defined and studied the class $R_{b,p}(\phi)$ to be the class of all functions $f \in \mathcal{A}_p$ for which

$$1 + \frac{1}{b} \left(\frac{f'(z)}{pz^{p-1}} - 1 \right) \prec \phi(z) \quad (z \in \Delta, b \in \mathbb{C} \setminus \{0\}). \quad (1.11)$$

Note that $S_{1,1}^*(\phi) = S^*(\phi)$ and $C_{1,1}(\phi) = C(\phi)$, the classes introduced and studied by Ma and Minda [2]. The familiar class $S^*(\gamma)$ of *starlike functions* of order γ and the class $C(\gamma)$ of *convex functions* of order γ , $0 \leq \gamma < 1$ are the special case of $S_{1,1}^*(\phi)$ and $C_{1,1}(\phi)$, respectively, when $\phi(z) = (1 + (1 - 2\gamma)z)/(1 - z)$.

Owa [3] introduced and studied the class $H_p(A, B, \alpha, \beta)$ of all functions $f \in \mathcal{A}_p$ satisfying

$$(1 - \beta) \left(\frac{f(z)}{z^p} \right)^\alpha + \beta \frac{zf'(z)}{pf(z)} \left(\frac{f(z)}{z^p} \right)^\alpha \prec \frac{1 + Az}{1 + Bz}, \quad (1.12)$$

where $z \in \Delta$, $-1 \leq B < A \leq 1$, $0 \leq \beta \leq 1$, $\alpha \geq 0$. We note that $H_1(A, B, \alpha, \beta)$ is a subclass of Bazilevič functions [4].

Motivated by the classes $H_p(A, B, \alpha, \beta)$ and $R_{b,p}(\phi)$ studied, respectively, by Owa [3] and Ali et al. [1], we now define a class of functions which extends the classes $S_{b,p}^*(\phi)$, $H_p(A, B, \alpha, \beta)$, and $R_{b,p}(\phi)$ in the following.

Definition 1.1. Let $\phi(z)$ be a univalent starlike function with respect to 1 which maps the open unit disk Δ onto a region in the right half-plane and is symmetric with respect to the real axis, $\phi(0) = 1$ and $\phi'(0) > 0$. A function $f \in \mathcal{A}_p$ is in the class $R_{p,b,\alpha,\beta}(\phi)$ if

$$1 + \frac{1}{b} \left\{ (1 - \beta) \left(\frac{f(z)}{z^p} \right)^\alpha + \beta \frac{zf'(z)}{pf(z)} \left(\frac{f(z)}{z^p} \right)^\alpha - 1 \right\} \prec \phi(z) \quad (0 \leq \beta \leq 1, \alpha \geq 0). \quad (1.13)$$

Also, $R_{p,b,\alpha,\beta,g}(\phi)$ is the class of all functions $f \in \mathcal{A}_p$ for which $f * g \in R_{p,b,\alpha,\beta}(\phi)$, where g is a fixed function with positive coefficients.

The class $R_{p,b,\alpha,\beta}(\phi)$ reduces to the following earlier classes.

- (1) $R_{p,b,0,1}(\phi) \equiv S_{b,p}^*(\phi)$ introduced and studied by Ali et al. [1].
- (2) $R_{p,b,1,1}(\phi) \equiv R_{b,p}(\phi)$ introduced and studied by Ali et al. [1].
- (3) $R_{1,1,\alpha,1}(\phi) \equiv B^\alpha(\phi)$ introduced and studied by Ravichandran et al. [5].
- (4) For $\phi(z) = (1 + Az)/(1 + Bz)$, the class $R_{p,1,\alpha,\beta}(\phi)$ reduces to $H_p(A, B, \alpha, \beta)$ introduced and studied by Owa [3].
- (5) For $\phi(z) = (1 + (1 - 2\gamma)z)/(1 - z)$, the class $R_{p,1,\alpha,0}(\phi)$ reduces to

$$\begin{aligned} H_p(1 - 2\gamma, -1, \alpha, 0) &\equiv \mathcal{B}_p(\gamma, \alpha) \\ &= \left\{ f \in \mathcal{A}_p : \operatorname{Re} \left(\frac{f(z)}{z^p} \right)^\alpha > \gamma, 0 \leq \gamma < 1, z \in \Delta \right\}. \end{aligned} \quad (1.14)$$

- (6) For $\phi(z) = (1 + (1 - 2\gamma)z)/(1 - z)$, the class $R_{p,1,\alpha,1}(\phi)$ reduces to

$$\begin{aligned} H_p(1 - 2\gamma, -1, \alpha, 1) &\equiv \mathcal{C}_p(\gamma, \alpha) \\ &= \left\{ f \in \mathcal{A}_p : \operatorname{Re} \left(\frac{f'(z)(f(z))^{\alpha-1}}{pz^{p-1}} \right) > \gamma, 0 \leq \gamma < 1, z \in \Delta \right\}. \end{aligned} \quad (1.15)$$

- (7) $R_{1,1,0,1}(\phi) \equiv S^*(\phi)$ [2].

Very recently, Ali et al. [1] obtained the sharp coefficient inequality for functions in the class $S_{b,p}^*(\phi)$ and many other subclasses \mathcal{A}_p .

In the present paper, we prove a sharp coefficient inequality in Theorem 2.1 for the more general class $R_{p,1,\alpha,\beta}(\phi)$. Also we give applications of our results to certain functions defined through Hadamard product. The results obtained in this paper generalize the results obtained by Ali et al. [1], Ma and Minda [2], Ravichandran et al. [5], and Srivastava and Mishra [6].

Let Ω be the class of analytic functions of the form

$$w(z) = w_1 z + w_2 z^2 + \dots \quad (1.16)$$

in the open unit disk Δ satisfying $|w(z)| < 1$.

To prove our main result, we need the following.

LEMMA 1.2 [1]. *If $w \in \Omega$, then*

$$|w_2 - tw_1^2| \leq \begin{cases} -t & \text{if } t < -1, \\ 1 & \text{if } -1 \leq t \leq 1, \\ t & \text{if } t > 1. \end{cases} \quad (1.17)$$

When $t < -1$ or $t > 1$, the equality holds if and only if $w(z) = z$ or one of its rotations. If $-1 < t < 1$, then equality holds if and only if $w(z) = z^2$ or one of its rotations. Equality holds for $t = -1$ if and only if

$$w(z) = z \frac{\lambda + z}{1 + \lambda z} \quad (0 \leq \lambda \leq 1) \quad (1.18)$$

or one of its rotations, while for $t = 1$, the equality holds if and only if

$$w(z) = -z \frac{\lambda + z}{1 + \lambda z} \quad (0 \leq \lambda \leq 1) \quad (1.19)$$

or one of its rotations.

Although the above upper bound is sharp, it can be improved as follows when $-1 < t < 1$:

$$\begin{aligned} |w_2 - tw_1^2| + (t+1)|w_1|^2 &\leq 1 \quad (-1 < t \leq 0), \\ |w_2 - tw_1^2| + (1-t)|w_1|^2 &\leq 1 \quad (0 < t < 1). \end{aligned} \quad (1.20)$$

LEMMA 1.3 [7]. If $w \in \Omega$, then for any complex number t ,

$$|w_2 - tw_1^2| \leq \max \{1; |t|\}. \quad (1.21)$$

The result is sharp for the functions $w(z) = z$ or $w(z) = z^2$.

LEMMA 1.4 [8]. If $w \in \Omega$, then for any real numbers q_1 and q_2 , the following sharp estimate holds:

$$|w_3 + q_1 w_1 w_2 + q_2 w_1^3| \leq H(q_1, q_2), \quad (1.22)$$

where

$$H(q_1, q_2) = \begin{cases} 1 & \text{for } (q_1, q_2) \in D_1 \cup D_2, \\ |q_2| & \text{for } (q_1, q_2) \in \bigcup_{k=3}^7 D_k, \\ \frac{2}{3}(|q_1| + 1) \left(\frac{|q_1| + 1}{3(|q_1| + 1 + q_2)} \right)^{1/2} & \text{for } (q_1, q_2) \in D_8 \cup D_9, \\ \frac{q_2}{3} \left(\frac{q_1^2 - 4}{q_1^2 - 4q_2} \right) \left(\frac{q_1^2 - 4}{3(q_2 - 1)} \right)^{1/2} & \text{for } (q_1, q_2) \in D_{10} \cup D_{11} \setminus \{\pm 2, 1\}, \\ \frac{2}{3}(|q_1| - 1) \left(\frac{|q_1| - 1}{3(|q_1| - 1 - q_2)} \right)^{1/2} & \text{for } (q_1, q_2) \in D_{12}. \end{cases} \quad (1.23)$$

The extremal functions, up to rotations, are of the form

$$\begin{aligned}
 w(z) &= z^3, & w(z) &= z, & w(z) = w_0(z) &= \frac{(z[(1-\lambda)\varepsilon_2 + \lambda\varepsilon_1] - \varepsilon_1\varepsilon_2 z)}{1 - [(1-\lambda)\varepsilon_1 + \lambda\varepsilon_2]z}, \\
 w(z) &= w_1(z) = \frac{z(t_1 - z)}{1 - t_1 z}, & w(z) = w_2(z) &= \frac{z(t_2 + z)}{1 + t_2 z}, \\
 |\varepsilon_1| &= |\varepsilon_2| = 1, & \varepsilon_1 &= t_0 - e^{-i\theta_0/2}(a \mp b), & \varepsilon_2 &= -e^{-i\theta_0/2}(ia \pm b), \\
 a &= t_0 \cos \frac{\theta_0}{2}, & b &= \sqrt{1 - t_0^2 \sin^2 \frac{\theta_0}{2}}, & \lambda &= \frac{b \pm a}{2b}, \\
 t_0 &= \left[\frac{2q_2(q_1^2 + 2) - 3q_1^2}{3(q_2 - 1)(q_1^2 - 4q_2)} \right]^{1/2}, & t_1 &= \left(\frac{|q_1| + 1}{3(|q_1| + 1 + q_2)} \right)^{1/2}, \\
 t_2 &= \left(\frac{|q_1| - 1}{3(|q_1| - 1 - q_2)} \right)^{1/2}, & \cos \frac{\theta_0}{2} &= \frac{q_1}{2} \left[\frac{q_2(q_1^2 + 8) - 2(q_1^2 + 2)}{2q_2(q_1^2 + 2) - 3q_1^2} \right]. \tag{1.24}
 \end{aligned}$$

The sets D_k , $k = 1, 2, \dots, 12$, are defined as follows:

$$\begin{aligned}
 D_1 &= \left\{ (q_1, q_2) : |q_1| \leq \frac{1}{2}, |q_2| \leq 1 \right\}, \\
 D_2 &= \left\{ (q_1, q_2) : \frac{1}{2} \leq |q_1| \leq 2, \frac{4}{27}(|q_1| + 1)^3 - (|q_1| + 1) \leq q_2 \leq 1 \right\}, \\
 D_3 &= \left\{ (q_1, q_2) : |q_1| \leq \frac{1}{2}, q_2 \leq -1 \right\}, \\
 D_4 &= \left\{ (q_1, q_2) : |q_1| \geq \frac{1}{2}, q_2 \leq -\frac{2}{3}(|q_1| + 1) \right\}, \\
 D_5 &= \left\{ (q_1, q_2) : |q_1| \leq 2, q_2 \geq 1 \right\}, \\
 D_6 &= \left\{ (q_1, q_2) : 2 \leq |q_1| \leq 4, q_2 \geq \frac{1}{12}(q_1^2 + 8) \right\}, \\
 D_7 &= \left\{ (q_1, q_2) : |q_1| \geq 4, q_2 \geq \frac{2}{3}(|q_1| - 1) \right\}, \\
 D_8 &= \left\{ (q_1, q_2) : \frac{1}{2} \leq |q_1| \leq 2, -\frac{2}{3}(|q_1| + 1) \leq q_2 \leq \frac{4}{27}(|q_1| + 1)^3 - (|q_1| + 1) \right\}, \\
 D_9 &= \left\{ (q_1, q_2) : |q_1| \geq 2, -\frac{2}{3}(|q_1| + 1) \leq q_2 \leq \frac{2|q_1|(|q_1| + 1)}{q_1^2 + 2|q_1| + 4} \right\}, \\
 D_{10} &= \left\{ (q_1, q_2) : 2 \leq |q_1| \leq 4, \frac{2|q_1|(|q_1| + 1)}{q_1^2 + 2|q_1| + 4} \leq q_2 \leq \frac{1}{12}(q_1^2 + 8) \right\}, \\
 D_{11} &= \left\{ (q_1, q_2) : |q_1| \geq 4, \frac{2|q_1|(|q_1| + 1)}{q_1^2 + 2|q_1| + 4} \leq q_2 \leq \frac{2|q_1|(|q_1| - 1)}{q_1^2 - 2|q_1| + 4} \right\}, \\
 D_{12} &= \left\{ (q_1, q_2) : |q_1| \geq 4, \frac{2|q_1|(|q_1| - 1)}{q_1^2 - 2|q_1| + 4} \leq q_2 \leq \frac{2}{3}(|q_1| - 1) \right\}. \tag{1.25}
 \end{aligned}$$

2. Coefficient bounds

By making use of Lemmas 1.2–1.4, we prove the following.

THEOREM 2.1. *Let $\phi(z) = 1 + B_1z + B_2z^2 + B_3z^3 + \dots$, where B_n 's are real with $B_1 > 0$ and $B_2 \geq 0$. Let $0 < \beta \leq 1$, $\alpha \geq 0$, $0 \leq \mu \leq 1$, and*

$$\begin{aligned}\sigma_1 &:= \frac{(\alpha p + \beta)^2}{2pB_1^2(\alpha p + 2\beta)} \left\{ 2(B_2 - B_1) - pB_1^2 \frac{(\alpha - 1)(\alpha p + 2\beta)}{(\alpha + \beta)^2} \right\}, \\ \sigma_2 &:= \frac{(\alpha p + \beta)^2}{2pB_1^2(\alpha p + 2\beta)} \left\{ 2(B_2 + B_1) - pB_1^2 \frac{(\alpha - 1)(\alpha p + 2\beta)}{(\alpha + \beta)^2} \right\}, \\ \sigma_3 &:= \frac{(\alpha p + \beta)^2}{2pB_1^2(\alpha p + 2\beta)} \left\{ 2B_2 - pB_1^2 \frac{(\alpha - 1)(\alpha p + 2\beta)}{(\alpha + \beta)^2} \right\}, \\ \Lambda(p, \alpha, \beta, \mu) &:= \frac{(\alpha p + 2\beta)(2\mu + \alpha - 1)}{2(\alpha p + \beta)^2}.\end{aligned}\tag{2.1}$$

If $f(z)$ given by (1.1) belongs to $R_{p,1,\alpha,\beta}(\phi)$, then

$$|a_{p+2} - \mu a_{p+1}^2| \leq \begin{cases} \frac{p}{\alpha p + 2\beta} \{B_2 - pB_1^2 \Lambda(p, \alpha, \beta, \mu)\} & \text{if } \mu < \sigma_1, \\ \frac{pB_1}{\alpha p + 2\beta} & \text{if } \sigma_1 \leq \mu \leq \sigma_2, \\ -\frac{p}{\alpha p + 2\beta} \{B_2 - pB_1^2 \Lambda(p, \alpha, \beta, \mu)\} & \text{if } \mu > \sigma_2. \end{cases}\tag{2.2}$$

Further, if $\sigma_1 \leq \mu \leq \sigma_3$, then

$$|a_{p+2} - \mu a_{p+1}^2| + \frac{1}{2pB_1} \left\{ 2 \left(1 - \frac{B_2}{B_1} \right) \frac{(\alpha p + \beta)^2}{\alpha p + 2\beta} + (2\mu + \alpha - 1)pB_1 \right\} |a_{p+1}|^2 \leq \frac{pB_1}{\alpha p + 2\beta}.\tag{2.3}$$

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$|a_{p+2} - \mu a_{p+1}^2| + \frac{1}{2pB_1} \left\{ 2 \left(1 + \frac{B_2}{B_1} \right) \frac{(\alpha p + \beta)^2}{\alpha p + 2\beta} - (2\mu + \alpha - 1)pB_1 \right\} |a_{p+1}|^2 \leq \frac{pB_1}{\alpha p + 2\beta}.\tag{2.4}$$

For any complex number μ ,

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{pB_1}{\alpha p + 2\beta} \max \left\{ 1, \left| \frac{pB_1}{2} \Lambda(p, \alpha, \beta, \mu) - \frac{B_2}{B_1} \right| \right\}.\tag{2.5}$$

Further,

$$|a_{p+3}| \leq \frac{pB_1}{\alpha p + 3\beta} H(q_1, q_2),\tag{2.6}$$

where $H(q_1, q_2)$ is as defined in Lemma 1.4,

$$\begin{aligned} q_1 &:= 2 \frac{B_2}{B_1} + p B_1 \frac{(1-\alpha)(\alpha p + 3\beta)}{(\alpha p + \beta)(\alpha p + 2\beta)}, \\ q_2 &:= \frac{B_3}{B_1} + p^2 B_1^2 \frac{(\alpha-1)(2\alpha-1)(\alpha p + 3\beta)}{6(\alpha p + \beta)^3} + p B_2 \frac{(1-\alpha)(\alpha p + 3\beta)}{(\alpha p + \beta)(\alpha p + 2\beta)}. \end{aligned} \quad (2.7)$$

These results are sharp.

Proof. If $f(z) \in R_{p,1,\alpha,\beta}(\phi)$, then there is a Schwarz function

$$w(z) = w_1 z + w_2 z^2 + \dots \in \Omega \quad (2.8)$$

such that

$$(1-\beta) \left(\frac{f(z)}{z^p} \right)^\alpha + \beta \frac{zf'(z)}{pf(z)} \left(\frac{f(z)}{z^p} \right)^\alpha = \phi(w(z)). \quad (2.9)$$

Since

$$\begin{aligned} &(1-\beta) \left(\frac{f(z)}{z^p} \right)^\alpha + \beta \frac{zf'(z)}{pf(z)} \left(\frac{f(z)}{z^p} \right)^\alpha \\ &= \begin{cases} 1 + \frac{1}{p}(\alpha p + \beta)a_{p+1}z + \frac{1}{2p}(\alpha p + 2\beta)\{2a_{p+2} + (\alpha - 1)a_{p+1}^2\}z^2 \\ + \frac{\alpha p + 3\beta}{p}\left\{a_{p+3} + (\alpha - 1)a_{p+1}a_{p+2} + \frac{(\alpha - 1)(\alpha - 2)}{6}a_{p+1}^3\right\}z^3 + \dots, \end{cases} \end{aligned} \quad (2.10)$$

from (2.9), we have

$$\begin{aligned} a_{p+1} &= \frac{pB_1 w_1}{\alpha p + \beta}, \\ a_{p+2} &= \frac{pB_1}{\alpha p + 2\beta} \left\{ w_2 - w_1^2 \left\{ pB_1 \left(\frac{\alpha-1}{2} \right) \left(\frac{\alpha p + 2\beta}{(\alpha p + \beta)^2} \right) - \frac{B_2}{B_1} \right\} \right\}, \\ a_{p+3} &= \frac{pB_1}{\alpha p + 3\beta} \{w_3 + q_1 w_1 w_2 + q_3 w_1^3\}, \end{aligned} \quad (2.11)$$

where q_1 and q_2 as defined in (2.7). Therefore, we have

$$a_{p+2} - \mu a_{p+1}^2 = \frac{pB_1}{\alpha p + 2\beta} \{w_2 - \nu w_1^2\}, \quad (2.12)$$

where

$$\nu := pB_1 \Lambda(p, \alpha, \beta, \mu) - \frac{B_2}{B_1}. \quad (2.13)$$

The results (2.2)–(2.5) are established by an application of Lemma 1.2, inequality (2.5) by Lemma 1.3, and (2.6) follows from Lemma 1.4. To show that the bounds in (2.2)–(2.5)

are sharp, we define the functions $K_{\phi n}$ ($n = 2, 3, \dots$) by

$$(1 - \beta) \left(\frac{K_{\phi n}(z)}{z^p} \right)^\alpha + \beta \frac{z K'_{\phi n}(z)}{p f(z)} \left(\frac{K_{\phi n}(z)}{z^p} \right)^\alpha = \phi(z^{n-1}), \quad K_{\phi n}(0) = 0 = [K_{\phi n}]'(0) - 1 \quad (2.14)$$

and the functions F_λ and G_λ ($0 \leq \lambda \leq 1$) by

$$\begin{aligned} (1 - \beta) \left(\frac{F_\lambda(z)}{z^p} \right)^\alpha + \beta \frac{z F_\lambda(z)}{p f(z)} \left(\frac{F_\lambda(z)}{z^p} \right)^\alpha &= \phi \left(\frac{z(z + \lambda)}{1 + \lambda z} \right), \quad F_\lambda(0) = 0 = F'_\lambda(0) - 1, \\ (1 - \beta) \left(\frac{G_\lambda(z)}{z^p} \right)^\alpha + \beta \frac{z G_\lambda(z)}{p f(z)} \left(\frac{G_\lambda(z)}{z^p} \right)^\alpha &= \phi \left(-\frac{z(z + \lambda)}{1 + \lambda z} \right), \quad G_\lambda(0) = 0 = G'_\lambda(0) - 1. \end{aligned} \quad (2.15)$$

Clearly, the functions $K_{\phi n}, F_\lambda, G_\lambda \in R_{p,1,\alpha,\beta}(\phi)$. Also we write $K_\phi := K_{\phi 2}$. If $\mu < \sigma_1$ or $\mu > \sigma_2$, then the equality holds if and only if f is K_ϕ or one of its rotations. When $\sigma_1 < \mu < \sigma_2$, then the equality holds if and only if f is $K_{\phi 3}$ or one of its rotations. If $\mu = \sigma_1$, then the equality holds if and only if f is F_λ or one of its rotations. If $\mu = \sigma_2$, then the equality holds if and only if f is G_λ or one of its rotations. \square

Remark 2.2. For $\alpha = 0$ and $\beta = 1$, results (2.2)–(2.6) coincide with the results obtained for the class $S_p^*(\phi)$ by Ali et al. [1].

Remark 2.3. For $\alpha = 0$, $p = 1$ and $\beta = 1$, results (2.2)–(2.6) coincide with the results obtained for the class $S^*(\phi)$ by Ma and Minda [2].

Remark 2.4. For $p = 1$ and $\beta = 1$, results (2.2)–(2.6) coincide with the results obtained for the Bazilevic class $B^\alpha(\phi)$ by Ravichandran et al. [5].

3. Applications to functions defined by convolution

We define $R_{p,b,\alpha,\beta,g}(\phi)$ to be the class of all functions $f \in \mathcal{A}_p$ for which $f * g \in R_{p,b,\alpha,\beta}(\phi)$, where g is a fixed function with positive coefficients and the class $R_{p,b,\alpha,\beta}(\phi)$ is as defined in Definition 1.1. In Theorem 2.1, we obtained the coefficient estimate for the class $R_{p,1,\alpha,\beta}(\phi)$. Now, we obtain the coefficient estimate for the class $R_{p,1,\alpha,\beta,g}(\phi)$.

THEOREM 3.1. Let $\phi(z) = 1 + B_1 z + B_2 z^2 + B_3 z^3 + \dots$, where B_n 's are real with $B_1 > 0$ and $B_2 \geq 0$. Let $0 < \beta \leq 1$, $\alpha \geq 0$, $0 \leq \mu \leq 1$, and

$$\begin{aligned} \sigma_1 &:= \frac{g_{p+1}^2}{g_{p+2}} \frac{(\alpha p + \beta)^2}{2p B_1^2 (\alpha p + 2\beta)} \left\{ 2(B_2 - B_1) - p B_1^2 \frac{(\alpha - 1)(\alpha p + 2\beta)}{(\alpha + \beta)^2} \right\}, \\ \sigma_2 &:= \frac{g_{p+1}^2}{g_{p+2}} \frac{(\alpha p + \beta)^2}{2p B_1^2 (\alpha p + 2\beta)} \left\{ 2(B_2 + B_1) - p B_1^2 \frac{(\alpha - 1)(\alpha p + 2\beta)}{(\alpha + \beta)^2} \right\}, \\ \sigma_3 &:= \frac{g_{p+1}^2}{g_{p+2}} \frac{(\alpha p + \beta)^2}{2p B_1^2 (\alpha p + 2\beta)} \left\{ 2B_2 - p B_1^2 \frac{(\alpha - 1)(\alpha p + 2\beta)}{(\alpha + \beta)^2} \right\}, \\ \Lambda_1(p, \alpha, \beta, g, \mu) &:= \frac{(\alpha p + 2\beta)(2\mu((g_{p+2})/(g_{p+1}^2)) + \alpha - 1)}{2(\alpha p + \beta)^2}. \end{aligned} \quad (3.1)$$

If $f(z)$ given by (1.1) belongs to $R_{p,1,\alpha,\beta,g}(\phi)$, then

$$|a_{p+2} - \mu a_{p+1}^2| \leq \begin{cases} \frac{p}{(\alpha p + 2\beta)g_{p+2}} \{B_2 - pB_1^2 \Lambda_1(p, \alpha, \beta, g, \mu)\} & \text{if } \mu < \sigma_1, \\ \frac{pB_1}{(\alpha p + 2\beta)g_{p+2}} & \text{if } \sigma_1 \leq \mu \leq \sigma_2, \\ -\frac{p}{(\alpha p + 2\beta)g_{p+2}} \{B_2 - pB_1^2 \Lambda_1(p, \alpha, \beta, g, \mu)\} & \text{if } \mu > \sigma_2. \end{cases} \quad (3.2)$$

Further, if $\sigma_1 \leq \mu \leq \sigma_3$, then

$$\begin{aligned} |a_{p+2} - \mu a_{p+1}^2| &+ \frac{g_{p+1}^2}{g_{p+2}} \frac{1}{2pB_1} \left\{ 2 \left(1 - \frac{B_2}{B_1}\right) \frac{(\alpha p + \beta)^2}{\alpha p + 2\beta} + (2\mu + \alpha - 1)pB_1 \right\} |a_{p+1}|^2 \\ &\leq \frac{pB_1}{(\alpha p + 2\beta)g_{p+2}}. \end{aligned} \quad (3.3)$$

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$\begin{aligned} |a_{p+2} - \mu a_{p+1}^2| &+ \frac{g_{p+1}^2}{g_{p+2}} \frac{1}{2pB_1} \left\{ 2 \left(1 + \frac{B_2}{B_1}\right) \frac{(\alpha p + \beta)^2}{\alpha p + 2\beta} - (2\mu + \alpha - 1)pB_1 \right\} |a_{p+1}|^2 \\ &\leq \frac{pB_1}{(\alpha p + 2\beta)g_{p+2}}. \end{aligned} \quad (3.4)$$

For any complex number μ ,

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{pB_1}{(\alpha p + 2\beta)g_{p+2}} \max \left\{ 1, \left| \frac{pB_1}{2} \Lambda_1(p, \alpha, \beta, g, \mu) - \frac{B_2}{B_1} \right| \right\}. \quad (3.5)$$

Further,

$$|a_{p+3}| \leq \frac{pB_1}{(\alpha p + 3\beta)g_{p+3}} H(q_1, q_2), \quad (3.6)$$

where $H(q_1, q_2)$ is as defined in Lemma 1.4,

$$\begin{aligned} q_1 &:= 2 \frac{B_2}{B_1} + pB_1 \frac{(1 - \alpha)(\alpha p + 3\beta)}{(\alpha p + \beta)(\alpha p + 2\beta)}, \\ q_2 &:= \frac{B_3}{B_1} + p^2 B_1^2 \frac{(\alpha - 1)(2\alpha - 1)(\alpha p + 3\beta)}{6(\alpha p + \beta)^3} + pB_2 \frac{(1 - \alpha)(\alpha p + 3\beta)}{(\alpha p + \beta)(\alpha p + 2\beta)}. \end{aligned} \quad (3.7)$$

These results are sharp.

Proof. If $f(z) \in R_{p,1,\alpha,\beta,g}(\phi)$, then there is a Schwarz function

$$w(z) = w_1 z + w_2 z^2 + \dots \in \Omega \quad (3.8)$$

such that

$$(1-\beta) \left(\frac{(f*g)(z)}{z^p} \right)^\alpha + \beta \frac{z(f*g)'(z)}{p(f*g)(z)} \left(\frac{(f*g)(z)}{z^p} \right)^\alpha = \phi(w(z)). \quad (3.9)$$

Hence

$$\begin{aligned} & (1-\beta) \left(\frac{(f*g)(z)}{z^p} \right)^\alpha + \beta \frac{z(f*g)'(z)}{p(f*g)(z)} \left(\frac{(f*g)(z)}{z^p} \right)^\alpha \\ &= \begin{cases} 1 + \frac{1}{p}(\alpha p + \beta) a_{p+1} g_{p+1} z + \frac{1}{2p}(\alpha p + 2\beta) \{2a_{p+2}g_{p+2} + (\alpha - 1)a_{p+1}^2 g_{p+1}^2\} z^2 \\ + \frac{\alpha p + 3\beta}{p} \left\{ a_{p+3}g_{p+3} + (\alpha - 1)a_{p+1}g_{p+1}a_{p+2}g_{p+2} + \frac{(\alpha - 1)(\alpha - 2)}{6}a_{p+1}^3 g_{p+1}^3 \right\} z^3 + \dots \end{cases} \end{aligned} \quad (3.10)$$

The remaining proof of the theorem is similar to the proof of Theorem 2.1 and hence omitted. \square

Remark 3.2. For $\alpha = 1$ and $\beta = 1$, results (3.2)–(3.4) coincide with the results obtained for the class $R_{b,p}(\phi)$ by Ali et al. [1].

Remark 3.3. For $p = 1$, $\alpha = 0$, and $\beta = 1$, results (3.5) coincide with the result for the class $S_b^*(\phi)$ obtained by Ravichandran et al. [9].

Remark 3.4. For $p = 1$, $\alpha = 1$, $\beta = 1$, and $\phi(z) = (1 + Az)/(1 + Bz)$, $-1 \leq B < A \leq 1$, inequality (3.5) coincides with the result obtained by Dixit and Pal [10].

Remark 3.5. For $p = 1$, $\alpha = 0$, and $\beta = 1$,

$$\begin{aligned} g_2 &:= \frac{\Gamma(3)\Gamma(2-\lambda)}{\Gamma(3-\lambda)} = \frac{2}{2-\lambda}, \\ g_3 &:= \frac{\Gamma(4)\Gamma(2-\lambda)}{\Gamma(4-\lambda)} = \frac{6}{(2-\lambda)(3-\lambda)}, \\ B_1 &= \frac{8}{\pi^2}, \quad B_2 = \frac{16}{3\pi^2}, \end{aligned} \quad (3.11)$$

in inequalities (3.2)–(3.4), we get the result obtained by Srivastava and Mishra [6].

THEOREM 3.6. *Let $\phi(z)$ be as in Theorem 2.1. If $f(z)$ given by (1.1) belongs to $R_{p,b,\alpha,\beta,g}(\phi)$, then for any complex number μ , with $B_1 > 0$, $B_2 \geq 0$, $0 < \beta \leq 1$, $\alpha \geq 0$,*

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{p|b|B_1}{(\alpha p + 2\beta)g_{p+2}} \max \left\{ 1, \left| b p B_1 \Lambda_2(p, b, \alpha, \mu, g) + \frac{B_2}{B_1} \right| \right\}, \quad (3.12)$$

where

$$\Lambda_2(p, b, \alpha, \beta, \mu, g) := \frac{(\alpha p + 2\beta)(2\mu((g_{p+2})/(g_{p+1}^2)) + \alpha - 1)}{2(\alpha p + \beta)^2}. \quad (3.13)$$

Proof. The proof is similar to the proof of Theorem 2.1 and hence omitted. \square

Remark 3.7. For $p = 1$, $\beta = 1$, and $\alpha = 0$, the result in (3.12) coincides with the results obtained by Ravichandran et al. [9].

References

- [1] R. M. Ali, V. Ravichandran, and N. Seenivasagan, “Coefficient bounds for p -valent functions,” to appear in *Applied Mathematics and Computation*.
- [2] W. C. Ma and D. Minda, “A unified treatment of some special classes of univalent functions,” in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, Z. Li, F. Ren, L. Yang, and S. Zhang, Eds., Conf. Proc. Lecture Notes Anal., I, pp. 157–169, International Press, Cambridge, Mass, USA, 1994.
- [3] S. Owa, “Properties of certain integral operators,” *Southeast Asian Bulletin of Mathematics*, vol. 24, no. 3, pp. 411–419, 2000.
- [4] I. E. Bazilevič, “On a case of integrability in quadratures of the Loewner-Kufarev equation,” *Matematicheskiĭ Sbornik. Novaya Seriya*, vol. 37(79), pp. 471–476, 1955.
- [5] V. Ravichandran, A. Gangadharan, and M. Darus, “Fekete-Szegö inequality for certain class of Bazilevic functions,” *Far East Journal of Mathematical Sciences*, vol. 15, no. 2, pp. 171–180, 2004.
- [6] H. M. Srivastava and A. K. Mishra, “Applications of fractional calculus to parabolic starlike and uniformly convex functions,” *Computers & Mathematics with Applications*, vol. 39, no. 3-4, pp. 57–69, 2000.
- [7] F. R. Keogh and E. P. Merkes, “A coefficient inequality for certain classes of analytic functions,” *Proceedings of the American Mathematical Society*, vol. 20, no. 1, pp. 8–12, 1969.
- [8] D. V. Prokhorov and J. Szynal, “Inverse coefficients for (α, β) -convex functions,” *Annales Universitatis Mariae Curie-Skłodowska. Sectio A*, vol. 35, pp. 125–143 (1984), 1981.
- [9] V. Ravichandran, Y. Polatoglu, M. Bolcal, and A. Sen, “Certain subclasses of starlike and convex functions of complex order,” *Hacettepe Journal of Mathematics and Statistics*, vol. 34, pp. 9–15, 2005.
- [10] K. K. Dixit and S. K. Pal, “On a class of univalent functions related to complex order,” *Indian Journal of Pure and Applied Mathematics*, vol. 26, no. 9, pp. 889–896, 1995.

C. Ramachandran: Department of Mathematics, Easwari Engineering College, Ramapuram, Chennai, Tamil Nadu 600 089, India
Email address: crjsp2004@yahoo.com

S. Sivasubramanian: Department of Mathematics, Easwari Engineering College, Ramapuram, Chennai, Tamil Nadu 600 089, India
Email address: sivasaisastha@rediffmail.com

H. Silverman: Department of Mathematics, College of Charleston, Charleston, SC 29424, USA
Email address: silvermanh@cofc.edu

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk