

*Research Article*

## Sufficient Conditions for Janowski Starlikeness

Rosihan M. Ali, V. Ravichandran, and N. Seenivasagan

Received 10 March 2007; Revised 15 June 2007; Accepted 4 July 2007

Recommended by Teodor Bulboaca

Let  $A, B, D, E \in [-1, 1]$  and let  $p(z)$  be an analytic function defined on the open unit disk,  $p(0) = 1$ . Conditions on  $A, B, D$ , and  $E$  are determined so that  $1 + \beta z p'(z)$  being subordinated to  $(1 + Dz)/(1 + Ez)$  implies that  $p(z)$  is subordinated to  $(1 + Az)/(1 + Bz)$ . Similar results are obtained by considering the expressions  $1 + \beta(zp'(z)/p(z))$  and  $1 + \beta(zp'(z)/p^2(z))$ . These results are then applied to obtain sufficient conditions for analytic functions to be Janowski starlike.

Copyright © 2007 Rosihan M. Ali et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### 1. Introduction

Let  $\mathcal{A}$  be the class of all analytic functions  $f(z)$  defined in the open unit disk  $U := \{z \in \mathbb{C} : |z| < 1\}$  and normalized by the conditions  $f(0) = 0 = f'(0) - 1$ . Let  $S^*[A, B]$  denote the class of functions  $f \in \mathcal{A}$  satisfying the subordination

$$\frac{zf'(z)}{f(z)} \prec \frac{1+Az}{1+Bz}, \quad (-1 \leq B < A \leq 1). \quad (1.1)$$

Functions in  $S^*[A, B]$  are called the *Janowski starlike functions* ([1, 2]). Certain well-known subclasses of starlike functions are special cases of the class  $S^*[A, B]$  for suitable choices of the parameters  $A$  and  $B$ . For example, when  $0 \leq \alpha < 1$ ,  $S^*[1 - 2\alpha, -1] =: S_\alpha^*$  is the familiar class of starlike functions of order  $\alpha$  and  $S^*[1 - \alpha, 0] = \{f \in \mathcal{A} : |zf'(z)/f(z) - 1| < 1 - \alpha \quad (z \in U)\} =: S^*(\alpha)$ . For  $0 < \alpha \leq 1$ , let  $S^*[\alpha, -\alpha] = \{f \in \mathcal{A} : |zf'(z)/f(z) - 1| < \alpha|zf'(z)/f(z) + 1| \quad (z \in U)\} =: S^*[\alpha]$ .

Silverman [3], Obradović and Tuneski [4], and many others (see [5–9]) have studied properties of functions defined in terms of the quotient  $(1 + zf''(z)/f'(z))/(zf'(z)/f(z))$ . In fact, Silverman [3] has obtained the order of starlikeness for the functions in the class

$G_b$  defined by

$$G_b := \left\{ f \in \mathcal{A} : \left| \frac{1 + zf''(z)/f'(z)}{zf'(z)/f(z)} - 1 \right| < b, 0 < b \leq 1, z \in U \right\} \quad (1.2)$$

and Obradović and Tuneski [4] improved the result of Silverman [3] by showing  $G_b \subset S^*[0, -b] \subset S^*(2/(1 + \sqrt{1 + 8b}))$ . Later, Tuneski [10] obtained conditions for the inclusion  $G_b \subset S^*[A, B]$  to hold. If we let  $zf'(z)/f(z) =: p(z)$ , then  $G_b \subset S^*[A, B]$  becomes

$$1 + \frac{zp'(z)}{p(z)^2} \prec 1 + bz \implies p(z) \prec \frac{1 + Az}{1 + Bz}. \quad (1.3)$$

Let  $f \in \mathcal{A}$  and  $0 \leq \alpha < 1$ . Frasin and Darus [11] have shown that

$$\frac{(zf(z))''}{f'(z)} - \frac{2zf'(z)}{f(z)} \prec \frac{(1 - \alpha)z}{2 - \alpha} \implies \left| \frac{z^2 f'(z)}{f^2(z)} - 1 \right| < 1 - \alpha. \quad (1.4)$$

Again by writing  $z^2 f'(z)/(f(z))^2$  as  $p(z)$ , we see that the above implication is special case of

$$1 + \beta \frac{zp'(z)}{p(z)} \prec \frac{1 + Dz}{1 + Ez} \implies p(z) \prec \frac{1 + Az}{1 + Bz}. \quad (1.5)$$

Another special case of the above implications was considered by Ponnusamy and Raja Sekaran [12].

Nunokawa et al. [13] have shown that if  $p(z)$  is analytic in  $U$ ,  $p(0) = 1$  and  $1 + zp'(z) \prec 1 + z$ , then  $p(z) \prec 1 + z$ . Using this, they have obtained a criterion for a normalized analytic function to be univalent. In this paper, we extend the result by replacing the subordinate function  $1 + z$  by a function of the form  $(1 + Dz)/(1 + Ez)$ . In fact, we determine conditions on  $A, B, D, E \in [-1, 1]$  so that

$$1 + \beta zp'(z) \prec \frac{1 + Dz}{1 + Ez} \implies p(z) \prec \frac{1 + Az}{1 + Bz}. \quad (1.6)$$

Similar results are obtained by considering the expressions  $1 + \beta(zp'(z)/p^2(z))$ ,  $1 + \beta(zp'(z)/p(z))$ . These results are then applied to obtain sufficient conditions for analytic functions to be Janowski starlike.

## 2. Differential subordination

LEMMA 2.1. *Let  $-1 \leq B < A \leq 1$ ,  $-1 \leq E < D \leq 1$ , and  $\beta \neq 0$ . Assume that*

$$(A - B)|\beta| \geq (D - E)(1 + B^2) + |2B(D - E) - E\beta(A - B)|. \quad (2.1)$$

If  $p(z)$  is analytic in  $U$  with  $p(0) = 1$  and

$$1 + \beta z p'(z) \prec \frac{1 + Dz}{1 + Ez}, \quad (2.2)$$

then

$$p(z) \prec \frac{1 + Az}{1 + Bz}. \quad (2.3)$$

*Proof.* Define the function  $P(z)$  by

$$P(z) := 1 + \beta z p'(z), \quad (2.4)$$

and the function  $w(z)$  by

$$w(z) := \frac{p(z) - 1}{A - Bp(z)}, \quad (2.5)$$

or equivalently by

$$p(z) = \frac{1 + Aw(z)}{1 + Bw(z)}. \quad (2.6)$$

Then  $w(z)$  is meromorphic in  $U$  and  $w(0) = 0$ . We need to show that  $|w(z)| < 1$  in  $U$ . By a computation, we get

$$P(z) = \frac{(1 + Bw(z))^2 + (A - B)\beta zw'(z)}{(1 + Bw(z))^2}. \quad (2.7)$$

Therefore

$$\frac{P(z) - 1}{D - EP(z)} = \frac{(A - B)\beta zw'(z)}{(D - E)(1 + Bw(z))^2 - E(A - B)\beta zw'(z)}. \quad (2.8)$$

Assume that there exists a point  $z_0 \in U$  such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1. \quad (2.9)$$

Then by [14, Lemma 1.3, page 28], there exists  $k \geq 1$  such that  $z_0 w'(z_0) = kw(z_0)$ . Let  $w(z_0) = e^{i\theta}$ . For this  $z_0$ , we have

$$\begin{aligned} \left| \frac{P(z_0) - 1}{D - EP(z_0)} \right| &= \frac{(A - B)k|\beta|}{[I^2 + (H - J)^2 + 4HJt^2 + 4I(H + J)t]^{1/2}} \\ &\geq \frac{(A - B)k|\beta|}{\max_{-1 \leq t \leq 1} \left\{ [I^2 + (H - J)^2 + 4HJt^2 + 4I(H + J)t]^{1/2} \right\}}, \end{aligned} \quad (2.10)$$

where  $I := 2B(D - E) - k\beta E(A - B)$ ,  $J := (D - E)B^2$ ,  $H := (D - E)$ , and  $t := \cos \theta$ . A computation shows that

$$\left| \frac{P(z_0) - 1}{D - EP(z_0)} \right| \geq \frac{(A - B)|\beta|k}{H + |I| + J}. \quad (2.11)$$

Yet another calculation shows that the function  $\psi(k) := (A - B)|\beta|k/(H + |I| + J)$  is an increasing function of  $k$ . Since  $k \geq 1$ , we have  $\psi(k) \geq \psi(1)$  and therefore

$$\left| \frac{P(z_0) - 1}{D - EP(z_0)} \right| \geq \frac{(A - B)|\beta|}{(D - E)(1 + B^2) + |2B(D - E) - E\beta(A - B)|}, \quad (2.12)$$

which by (2.1) is greater than or equal to 1. This contradicts  $P(z) \prec (1 + Dz)/(1 + Ez)$  and completes the proof.  $\square$

*Remark 2.2.* When  $\beta = 1$ ,  $E = 0 = B$ , and  $D = 1 = A$ , Lemma 2.1 reduces to [13, Lemma 1, page 1035]. Further if  $p(z) = z^2 f'(z)/f(z)^2$ , Lemma 2.1 reduces to [13, Theorem 1, page 1036].

By taking  $p(z) = zf'(z)/f(z)$  in Lemma 2.1, we have the following result.

**THEOREM 2.3.** *Let the conditions of Lemma 2.1 hold. If  $f \in \mathcal{A}$  satisfies*

$$1 + \beta \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \prec \frac{1 + Dz}{1 + Ez}, \quad (2.13)$$

*then  $f \in S^*[A, B]$ .*

By taking  $\beta = 1$ ,  $A = \alpha = -B$ , and  $D = -E = \delta$  ( $0 < \alpha, \delta \leq 1$ ) in Theorem 2.3, we have the following result.

**COROLLARY 2.4.** *Let  $0 < \alpha \leq 1$  and  $\delta = \alpha/(1 + \alpha)^2$ . If  $f \in \mathcal{A}$  satisfies*

$$\left| \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right| < \delta \left| 2 + \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right|, \quad (2.14)$$

*then  $f(z) \in \mathcal{S}^*[\alpha]$ .*

By taking  $\beta = 1$ ,  $A = 1 - 2\alpha$ ,  $B = -1$ ,  $D = (1 - \alpha)/2$ , and  $E = 0$  ( $0 \leq \alpha < 1$ ) in Theorem 2.3, we have the following result.

**COROLLARY 2.5.** *If  $f \in \mathcal{A}$  satisfies*

$$\left| \frac{zf'(z)}{f(z)} \left( 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right| < \frac{1 - \alpha}{3} \quad (0 \leq \alpha < 1), \quad (2.15)$$

*then  $f(z) \in \mathcal{S}_\alpha^*$ .*

By replacing  $p(z)$  by  $1/p(z)$ ,  $\beta = -1$ ,  $A$  by  $-B$ , and  $B$  by  $-A$  in Lemma 2.1, we have the following result.

LEMMA 2.6. Let  $-1 \leq B < A \leq 1$ ,  $-1 \leq E < D \leq 1$ . Assume that

$$(A - B) \geq (D - E)(1 + A^2) + |E(A - B) - 2A(D - E)|. \quad (2.16)$$

If  $p(z)$  is analytic in  $U$  with  $p(0) = 1$  and

$$1 + \frac{zp'(z)}{p^2(z)} \prec \frac{1 + Dz}{1 + Ez}, \quad (2.17)$$

then

$$p(z) \prec \frac{1 + Az}{1 + Bz}. \quad (2.18)$$

When  $p(z) = zf'(z)/f(z)$ , in Lemma 2.6, we have the following theorem.

THEOREM 2.7. Let  $-1 \leq B < A \leq 1$ ,  $-1 \leq E < D \leq 1$ . Assume that (2.16) holds. If  $f \in \mathcal{A}$  satisfies

$$\frac{1 + zf''(z)/f'(z)}{zf'(z)/f(z)} \prec \frac{1 + Dz}{1 + Ez}, \quad (2.19)$$

then  $f \in S^*[A, B]$ .

Example 2.8. If  $f \in G_{1-\alpha/(2-\alpha)^2}$  ( $0 \leq \alpha < 1$ ), then  $f \in S^*(\alpha)$ . If  $f \in \mathcal{A}$  satisfies

$$\left| 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right| < \beta \left| 1 + \frac{zf''(z)}{f'(z)} + \frac{zf'(z)}{f(z)} \right| \quad \left( \beta = \frac{\alpha}{1 + 3\alpha + \alpha^2}, 0 < \alpha \leq 1 \right), \quad (2.20)$$

then  $f \in S^*[\alpha]$ . Similarly if (2.20) holds with  $\beta = (1 - \alpha)/[1 + (1 - 2\alpha)^2 + |5\alpha - 3|]$  ( $0 \leq \alpha < 1$ ), then  $f \in S_\alpha^*$ .

Remark 2.9. When  $E = 0$  and  $D = b$  ( $0 < b \leq 1$ ), Corollary 2.5 reduces to [10, Corollary 2.6, page 203]. When  $A = 0 = E$  and  $D = -B = b$  ( $0 < b \leq 1$ ), Corollary 2.5 reduces to [4, Theorem 1, page 61]. When  $A = 0 = E$  and  $D = -B = 1$ , Corollary 2.5 reduces to [3, Corollary 1, page 76].

LEMMA 2.10. Let  $-1 \leq B < A \leq 1$ ,  $-1 \leq E < D \leq 1$ ,  $AB \geq 0$ , and  $\beta \neq 0$ . Assume that

$$|\beta|(A - B) \geq (D - E)(1 + AB) + |(D - E)(A + B) - E\beta(A - B)|. \quad (2.21)$$

Let  $p(z)$  be analytic in  $U$  with  $p(0) = 1$  and

$$1 + \beta \frac{zp'(z)}{p(z)} \prec \frac{1 + Dz}{1 + Ez}, \quad (2.22)$$

then

$$p(z) \prec \frac{1 + Az}{1 + Bz}. \quad (2.23)$$

*Proof.* The proof is similar to the proof of Lemma 2.1.  $\square$

*Remark 2.11.* When  $E\beta \leq 0$ ,  $AB \leq 0$ , Lemma 2.10 is valid provided the following conditions hold:

$$(1 - A\beta)^2 \{2E\beta(A + B)(D - E) - (A - B)[(D - E)^2 + (E\beta)^2]\} \geq 4\beta^2(A - B)AB \quad (2.24)$$

instead of (2.21).

*Remark 2.12.* When  $\beta = -1$ ,  $A = \lambda = E$ , and  $D = B = 0$  ( $|\lambda| \leq 1$ ), Lemma 2.10 reduces to [12, Theorem 1(iii), page 195].

*Example 2.13.* By taking  $\beta = 1$ ,  $B = 0$ ,  $D = A/(1 + A)$ , and  $E = 0$  in Lemma 2.10, we have the following result. Let  $0 < A \leq 1$ . Let  $p(z)$  be analytic in  $U$  with  $p(0) = 1$ . If  $|zp'(z)/p(z)| < A/(1 + A)$ , then  $p(z) \prec 1 + Az$ . When  $p(z) = zf'(z)/f(z)$ ,  $A = 1 - \alpha$ , we have the following result.

If  $f(z) \in \mathcal{A}$  satisfies

$$\left| 1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right| < \frac{1 - \alpha}{2 - \alpha} \quad (0 \leq \alpha < 1), \quad (2.25)$$

then  $f(z) \in \mathcal{G}^*(\alpha)$ .

By taking  $p(z) = z^2 f'(z)/f^2(z)$  in Lemma 2.10, we have the following result.

**THEOREM 2.14.** *Let the conditions of Lemma 2.10 hold. If  $f \in \mathcal{A}$  satisfies*

$$1 + \beta \left( \frac{(zf(z))''}{f'(z)} - \frac{2zf'(z)}{f(z)} \right) \prec \frac{1 + Dz}{1 + Ez}, \quad (2.26)$$

*then*

$$\frac{z^2 f'(z)}{f^2(z)} \prec \frac{1 + Az}{1 + Bz}. \quad (2.27)$$

*Remark 2.15.* When  $\beta = 1$ ,  $A = \alpha$ ,  $B = 0$ ,  $E = 0$ , and  $D = (1 - \alpha)/(2 - \alpha)$  ( $0 \leq \alpha < 1$ ), Theorem 2.14 reduces to [11, Theorem 2.4, page 307].

### Acknowledgment

The authors gratefully acknowledge the support from the research Grant IRPA 09-02-05-00020 EAR.

### References

- [1] W. Janowski, "Some extremal problems for certain families of analytic functions. I," *Annales Polonici Mathematici*, vol. 28, pp. 297–326, 1973.
- [2] Y. Polatoglu and M. Bolcal, "The radius of convexity for the class of Janowski convex functions of complex order," *Matematichki Vesnik*, vol. 54, no. 1-2, pp. 9–12, 2002.
- [3] H. Silverman, "Convex and starlike criteria," *International Journal of Mathematics and Mathematical Sciences*, vol. 22, no. 1, pp. 75–79, 1999.

- [4] M. Obradović and N. Tuneski, “On the starlike criteria defined by Silverman,” *Zeszyty Naukowe Politechniki Rzeszowskiej. Matematyka*, no. 24, pp. 59–64, 2000.
- [5] M. Nunokawa, S. Owa, H. Saitoh, A. Ikeda, and N. Koike, “Some results for strongly starlike functions,” *Journal of Mathematical Analysis and Applications*, vol. 212, no. 1, pp. 98–106, 1997.
- [6] M. Nunokawa, S. Owa, H. Saitoh, and N. Takahashi, “On a strongly starlikeness criteria,” *Bulletin of the Institute of Mathematics. Academia Sinica*, vol. 31, no. 3, pp. 195–199, 2003.
- [7] M. Obradović and S. Owa, “On some criterions for starlikeness of order  $\alpha$ ,” *Rendiconti di Matematica e delle sue Applicazioni. Serie VII*, vol. 8, no. 2, pp. 283–289, 1988.
- [8] V. Ravichandran and M. Darus, “On a criteria for starlikeness,” *International Mathematical Journal*, vol. 4, no. 2, pp. 119–125, 2003.
- [9] V. Ravichandran, M. Darus, and N. Seenivasagan, “On a criteria for strong starlikeness,” *The Australian Journal of Mathematical Analysis and Applications*, vol. 2, no. 1, pp. 1–12, 2005, article 6.
- [10] N. Tuneski, “On the quotient of the representations of convexity and starlikeness,” *Mathematische Nachrichten*, vol. 248–249, no. 1, pp. 200–203, 2003.
- [11] B. A. Frasin and M. Darus, “On certain analytic univalent functions,” *International Journal of Mathematics and Mathematical Sciences*, vol. 25, no. 5, pp. 305–310, 2001.
- [12] S. Ponnusamy and S. Rajasekaran, “New sufficient conditions for starlike and univalent functions,” *Soochow Journal of Mathematics*, vol. 21, no. 2, pp. 193–201, 1995.
- [13] M. Nunokawa, M. Obradović, and S. Owa, “One criterion for univalency,” *Proceedings of the American Mathematical Society*, vol. 106, no. 4, pp. 1035–1037, 1989.
- [14] S. Ruscheweyh, *Convolutions in Geometric Function Theory*, vol. 83 of *Seminar on Higher Mathematics*, Presses de l’Université de Montréal, Montreal, Quebec, Canada, 1982.

Rosihan M. Ali: School of Mathematical Sciences, Universiti Sains Malaysia,  
11800 USM Penang, Malaysia  
Email address: rosihan@cs.usm.my

V. Ravichandran: School of Mathematical Sciences, Universiti Sains Malaysia,  
11800 USM Penang, Malaysia  
Email address: vravi@cs.usm.my

N. Seenivasagan: School of Mathematical Sciences, Universiti Sains Malaysia,  
11800 USM Penang, Malaysia  
Email address: vasagan2000@yahoo.co.in

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)