

ON MODULES OF CONTINUOUS LINEAR MAPPINGS

DINAMÉRICO P. POMBO JR.

Instituto de Matemática
Universidade Federal Fluminense
Rua São Paulo, s/nº
24020-005 Niterói, RJ, BRASIL

(Received April 4, 1996)

ABSTRACT. Modules of continuous linear mappings with values in topological modules of continuous mappings are studied

KEY WORDS AND PHRASES: Topological modules, continuous linear mappings, continuous mappings.

1991 AMS SUBJECT CLASSIFICATION CODE: 46H25

INTRODUCTION

Grothendieck [1] has proved that, under certain conditions, spaces of continuous linear mappings on locally convex spaces are naturally isomorphic to spaces of continuous mappings on locally compact topological spaces, either as vector spaces or even as locally convex spaces. Non-linear versions of his results have been discussed in [2].

In the present note, we use Grothendieck's argument to show that modules of continuous linear mappings with values in topological modules of continuous mappings may be identified with modules of continuous mappings with values in topological modules of continuous linear forms.

Throughout this note, A denotes a commutative topological ring with an identity element and all A -modules under consideration are unitary. T represents an arbitrary topological space, $\mathcal{C}(T; S)$ the A -module of all continuous mappings from T into the topological A -module S and $\mathcal{C}(T) := \mathcal{C}(T; A)$. Given two topological A -modules R and S , $\mathcal{L}(R; S)$ represents the A -module of all continuous A -linear mappings from R into S and $R' := \mathcal{L}(R; A)$. Moreover, R'_s denotes the A -module R' endowed with the A -module topology τ_s of pointwise convergence.

PROPOSITION 1. Let E, F be topological A -modules and $u \in \mathcal{L}(E; F)$. Then ${}^t u \in \mathcal{L}(F'_s; E'_s)$ and ${}^t u$ transforms equicontinuous subsets of F' into equicontinuous subsets of E' , where ${}^t u$ is the adjoint of u .

PROOF. Straightforward.

Consider $\mathcal{C}(T)$ endowed with the topology of compact convergence. By Theorem 15.4 (1) of [5] and Proposition (a) of [3], $\mathcal{C}(T)$ is a topological A -module. (The proposition just mentioned will also ensure that other function spaces which will appear in the sequel are topological A -modules.)

For each $t \in T$, let $\delta(t) : \mathcal{C}(T) \rightarrow A$ be given by $\delta(t)(f) = f(t)$ for $f \in \mathcal{C}(T)$. Then $\delta(t) \in (\mathcal{C}(T))'$.

PROPOSITION 2. The mapping

$$\delta : t \in T \mapsto \delta(t) \in (\mathcal{C}(T))'$$

is continuous when $(\mathcal{C}(T))'$ is endowed with τ_s , and transforms compact subsets of T into equicontinuous subsets of $(\mathcal{C}(T))'$.

PROOF. Let $t_0 \in T$, $f \in \mathcal{C}(T)$ and W a neighborhood of zero in A . By the continuity of f at t_0 , there exists a neighborhood V of t_0 in T such that $f(t) - f(t_0) \in W$ for all $t \in V$, that is, $(\delta(t) - \delta(t_0))(f) \in W$ for all $t \in V$. This proves the continuity of δ at t_0 . Moreover, if K is a compact subset of T , then

$$\delta(K)(\{f \in \mathcal{C}(T); f(K) \subset W\}) \subset W.$$

This shows that $\delta(K)$ is equicontinuous, thereby concluding the proof.

Let E be a topological A -module. For each $u \in \mathcal{L}(E; \mathcal{C}(T))$, define $\Psi(u) = {}^t u \circ \delta$. By Propositions 1 and 2, $\Psi(u) \in \mathcal{C}(T; E'_s)$ and $\Psi(u)$ transforms compact subsets of T into equicontinuous subsets of E' . Let Ψ be the A -linear mapping

$$u \in \mathcal{L}(E; \mathcal{C}(T)) \mapsto \Psi(u) \in \mathcal{C}(T; E'_s),$$

and let H be the submodule of $\mathcal{C}(T; E'_s)$ formed by the continuous mappings $h : T \rightarrow E'_s$ such that $h(K)$ is an equicontinuous subset of E' for every compact subset K of T .

THEOREM. Ψ is an A -module isomorphism between $\mathcal{L}(E; \mathcal{C}(T))$ and H .

PROOF. We have just observed that $\text{Im}(\Psi) \subset H$.

We claim that Ψ is injective. Indeed, take a $u \in \mathcal{L}(E; \mathcal{C}(T))$ such that $\Psi(u) = 0$, and fix an $x \in E$. Then, for all $t \in T$,

$$[({}^t u \circ \delta)(t)](x) = [\delta(t) \circ u](x) = u(x)(t) = 0.$$

Therefore $u(x) = 0$, and so $u = 0$.

Now, let us verify that $H \subset \text{Im}(\Psi)$. Indeed, if $h \in H$, define $u(x)(t) = h(t)(x)$ for $x \in E$, $t \in T$. Then $u(x) \in \mathcal{C}(T)$ since $u(x) = \Phi \circ h$, where Φ is the τ_s -continuous A -linear mapping $\varphi \in E' \mapsto \varphi(x) \in A$. Moreover, it is easily seen that u is an A -linear mapping from E into $\mathcal{C}(T)$. In order to prove the continuity of u , let K be a compact subset of T and W a neighborhood of zero in A . By the equicontinuity of $h(K)$, there exists a neighborhood U of zero in E such that $h(K)(U) \subset W$. Therefore

$$u(U) \subset \{f \in \mathcal{C}(T); f(K) \subset W\},$$

which proves the continuity of u . Finally, it is clear that $\Psi(u) = h$, which concludes the proof of the theorem.

Let E be a topological A -module. Let \mathcal{M} be a family of bounded subsets of E such that for every $B_1, B_2 \in \mathcal{M}$ there is a $B_3 \in \mathcal{M}$ with $B_1 \cup B_2 \subset B_3$, and let $\tau_{\mathcal{M}}$ be the A -module topology on E' of \mathcal{M} -convergence. By Theorem 15.2 (1), (4) of [5], the set G of all mappings $g : T \rightarrow E'$ such that $g(K)$ is $\tau_{\mathcal{M}}$ -bounded in E' for every compact subset K of T is an A -module. By Theorem 25.5 of [5], $H \subset G$, and hence H is a submodule of G . Consider E' endowed with $\tau_{\mathcal{M}}$. Then G , endowed with the topology of compact convergence, is a topological A -module. Consider on H the A -module topology induced by that of G .

COROLLARY 1. Ψ is a topological A -module isomorphism between $\mathcal{L}_{\mathcal{M}}(E; \mathcal{C}(T))$ and H , where $\mathcal{L}_{\mathcal{M}}(E; \mathcal{C}(T))$ denotes $\mathcal{L}(E; \mathcal{C}(T))$ endowed with the A -module topology of \mathcal{M} -convergence.

PROOF. It suffices to observe that, if $B \in \mathcal{M}$, K is a compact subset of T and W is a neighborhood of zero in A , then $u \in \{v \in \mathcal{L}(E; \mathcal{C}(T)); v(B)(K) \subset W\}$ (a basic neighborhood of zero in $\mathcal{L}_{\mathcal{M}}(E; \mathcal{C}(T))$) if and only if $\Psi(u) \in \{h \in H; h(K)(B) \subset W\}$ (a basic neighborhood of zero in H).

If E is a barrelled topological A -module ([4], Definition 2.1), Theorem 15.4 (1) of [5] and Theorem 3.1 of [4] imply that $H = \mathcal{C}(T; E'_s)$. As a consequence, Corollary 1 yields:

COROLLARY 2. If E is a barrelled topological A -module, then $\mathcal{L}_{\mathcal{M}}(E; \mathcal{C}(T))$ and $\mathcal{C}(T; E'_s)$ are isomorphic as topological A -modules.

REFERENCES

- [1] GROTHENDIECK, A., *Espaces Vectoriels Topologiques*, 3rd ed., Publ. Soc. Mat. São Paulo (1964).
- [2] POMBO, D.P., Jr., On adjoints of non-linear mappings, *Bull. Austral. Math. Soc.* **33** (1986), 227-236.
- [3] POMBO, D.P., Jr., On the completion of certain topological modules, *Math. Japonica* **37** (1992), 333-336.
- [4] POMBO, D.P., Jr., On barrelled topological modules, *Internat. J. Math. & Math. Sci.* **19** (1996), 45-52.
- [5] WARNER, S., *Topological Fields*, Notas de Matemática No. 126, North-Holland Publishing Company (1989).

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru