

## QUASI-HOMOGENEOUS ASSOCIATIVE FUNCTIONS

BRUCE R. EBANKS

Department of Mathematics  
Marshall University  
Huntington, WV 25755, U. S. A.

(Received July 15, 1996 and in revised version March 18, 1997)

**ABSTRACT.** A triangular norm is a special kind of associative function on the closed unit interval  $[0,1]$ . Triangular norms (or  $t$ -norms) were introduced in the context of probabilistic metric space theory, and they have found applications also in other areas, such as fuzzy set theory. We determine the explicit forms of all  $t$ -norms which satisfy a generalized homogeneity property called quasi-homogeneity.

**KEY WORDS AND PHRASES:** Triangular norm ( $t$ -norm), homogeneous function, probabilistic metric space.

**1991 AMS SUBJECT CLASSIFICATION CODES:** 39B12, 54E70, 39B22.

### 0. INTRODUCTION

Let  $I = [0, 1]$  be the closed unit interval on the real line  $\mathbf{R}$ . A *triangular norm* (or  $t$ -norm) is a map  $T: I \times I \rightarrow I$  satisfying the following four hypotheses. First,  $T$  is associative:

$$T(x, T(y, z)) = T(T(x, y), z), \quad x, y, z \in I;$$

second,  $T$  is nondecreasing in each variable; third,  $T$  is commutative; and fourth, for all  $x \in I$ ,

$$T(x, 1) = x. \quad (0.1)$$

$T$ -norms were introduced by Menger [1] as a means of generalizing the triangle inequality to statistical (later, probabilistic) metric spaces. Associated with a family  $\{F_{pq}\}$  of probability distribution functions is a map  $T: I \times I \rightarrow I$  (later, a  $t$ -norm) such that

$$F_{pr}(x + y) \geq T[F_{pq}(x), F_{qr}(y)] \quad (0.2)$$

for all  $p, q, r$  in some space  $S$  and all nonnegative reals  $x$  and  $y$ . Interpreting  $F_{pq}(x)$  as the probability that the distance between  $p$  and  $q$  is less than  $x$ , inequality (0.2) means that the probability that the distance from  $p$  to  $r$  is less than  $x + y$  is at least as great as the  $T$ -value of the probabilities that the distance from  $p$  to  $q$  is less than  $x$  and the distance from  $q$  to  $r$  is less than  $y$ . For further information on the history, theory and applications of  $t$ -norms, see Schweizer and Sklar [2].

In light of the interpretation above, we examine now those  $t$ -norms satisfying an additional property. If the probabilities  $F_{pq}(x)$  and  $F_{qr}(y)$  on the right side of (0.2) are shrunk by a factor  $t \in I$ , then it may be reasonable to suppose that the probability  $F_{pr}(x + y)$  is also shrunk by a factor depending on  $t$ , possibly after some change of scale. More precisely, we shall study those  $t$ -norms  $T$  which satisfy

$$T(tx, ty) = \Pi^{-1}\{\varphi(t) \Pi[T(x, y)]\}, \quad (0.3)$$

for some map  $\varphi: I \rightarrow I$  and some continuous injection  $\Pi: I \rightarrow [0, \infty)$ . Such  $t$ -norms will be called *quasi-homogeneous*. If  $\Pi$  in (0.3) is the identity map, then  $T$  is called *homogeneous*, and the forms of such  $t$ -norms are known already (e.g. [3]). In that case, either  $T(x, y) = \text{Min}(x, y)$  with  $\varphi(t) = t$ , or  $T(x, y) = xy$  with  $\varphi(t) = t^2$ . We shall obtain that result as a corollary of the main result of the present paper, in which we determine the general solution of (0.3) for  $t$ -norms.

Before proceeding, let us observe some other properties of  $t$ -norms. It is easy to deduce the following from the definition of triangular norm:

$$T(x, 0) = T(0, x) = 0, \quad x \in I,$$

$$T(x, y) \leq \text{Min}(x, y), \quad x, y \in I$$

(i.e.  $\text{Min}$  is the “maximal”  $t$ -norm), so in particular

$$T(x, x) \leq x, \quad x \in I.$$

Note also that (0.3) forces  $\Pi$  to be strictly increasing. For, setting  $x = y = 0$  there, we have  $T(0, 0) = \Pi^{-1}\{\varphi(t) \Pi[T(0, 0)]\}$ . Since  $T(0, 0) = 0$ , this means  $\Pi(0) = \varphi(t) \Pi(0)$  for all  $t \in [0, 1]$ . Hence  $\Pi(0) = 0$ , since otherwise  $\varphi \equiv 1$ , which in (0.3) with  $t = 0$  would yield  $0 = T(x, y)$  for all  $x, y \in I$ , in violation of (0.1).

## 1. PRELIMINARIES

In the sequel, let  $\bar{\mathbf{R}}^+ = [0, \infty]$  be the extended nonnegative real half-line. The structure of continuous  $t$ -norms is known. (The continuity assumption can be weakened somewhat, but that is not relevant to the current discussion.)

**THEOREM 1.1.** (Schweizer and Sklar [2: Sections 5.3-5.5]) Let  $T: I \times I \rightarrow I$  be a continuous  $t$ -norm.

(i)  $T$  satisfies  $T(x, x) < x$  for all  $x \in (0, 1)$  if and only if  $T$  admits the representation

$$T(x, y) = f[g(x) + g(y)], \quad (1.1)$$

where  $g: I \rightarrow \bar{\mathbf{R}}^+$  is continuous, strictly decreasing with  $g(1) = 0$ , where  $f: \bar{\mathbf{R}}^+ \rightarrow I$  is onto, continuous, strictly decreasing on  $[0, g(0)]$  with  $f(u) = 0$  for  $u \geq g(0)$ , and where  $f \circ g$  is the identity map on  $I$  (i.e.  $g$  is a “quasi-inverse” of  $f$ ).

(ii) If  $T(x, x) = x$  for all  $x \in I$ , then  $T = \text{Min}$  on  $I \times I$ .

(iii) Otherwise, the semigroup  $(I, T)$  is an ordinal sum of semigroups  $\{(S_k, T_k)\}$ . Here each  $S_k$  is a proper closed (nontrivial) subinterval of  $I$ , each  $T_k$  admits a representation of the form (1.1) on  $S_k \times S_k$ , with  $g_k$  on  $S_k$  and  $f_k$  onto  $S_k$ , and  $T = \text{Min}$  on  $(I \times I) \setminus \bigcup_k (S_k \times S_k)$  [2].

**REMARK.** Any  $t$ -norm of the form given by case (i) of Theorem 1.1 is called *Archimedean*. If in addition  $g(0) = \infty$ , then  $f = g^{-1}$  and  $T$  is called *strict*.

We shall use Theorem 1.1 to find the forms of quasi-homogeneous  $t$ -norms. We shall also need the following known result.

**THEOREM 1.2.** (See e.g. Aczel and Dhombres [4: Chapter 15, Theorem 1]. The general solution of

$$g(tx) = a(t) g(x) + b(t), \quad (1.2)$$

among maps  $g, a, b: I \rightarrow \bar{\mathbf{R}}^+$ , is given by the following:

$$g(x) = \ell(x) + c, a(x) = 1, b(x) = \ell(x); \quad (1.3)$$

$$g(x) = c, a \text{ arbitrary}, b(x) = c[1 - a(x)]; \quad (1.4)$$

or

$$g(x) = cm(x) + d, a(x) = m(x), b(x) = d[1 - m(x)]; \quad (1.5)$$

for all  $x \in I$ , where  $c$  and  $d$  are arbitrary constants,  $\ell: I \rightarrow \bar{\mathbb{R}}^+$  is an arbitrary solution of the logarithmic functional equation

$$\ell(xy) = \ell(x) + \ell(y), \quad x, y \in I,$$

and  $m: I \rightarrow \bar{\mathbb{R}}^+$  is an arbitrary solution of the multiplicative functional equation

$$m(xy) = m(x)m(y), \quad x, y \in I.$$

## 2. DETERMINATION OF QUASI-HOMOGENEOUS T-NORMS

We prove first that quasi-homogeneity implies that  $T$  must be continuous and that only certain forms are possible for  $\varphi$ .

**LEMMA 2.1.** If a  $t$ -norm  $T$  is quasi-homogeneous in the sense of (0.3), then  $T$  is continuous and there exists a constant  $\alpha > 0$  such that

$$\varphi(t) = t^\alpha. \quad (2.1)$$

**PROOF:** Setting  $x = y = 1$  in (0.3), we find that

$$\Pi[T(t, t)] = \varphi(t) \Pi(1), \quad (2.2)$$

since  $T(1, 1) = 1$ . Since  $\Pi(1) \neq 0$ , this yields

$$\varphi(t) = \Pi(1)^{-1} \Pi[T(t, t)], \quad (2.3)$$

and shows immediately that  $\varphi$  is monotonic, with  $\varphi(1) = 1$ . It shows also that  $\varphi$  is multiplicative, since by (0.3) we have

$$\begin{aligned} \varphi(xy) &= \Pi(1)^{-1} \Pi[T(xy, xy)] \\ &= \Pi(1)^{-1} \varphi(x) \Pi[T(y, y)] \\ &= \varphi(x) \varphi(y). \end{aligned} \quad x, y \in I.$$

Moreover, (2.3) shows that  $\varphi(t) < 1$  for  $t < 1$ , for otherwise, since  $\Pi$  is strictly increasing we would have  $T(t, t) \geq 1$  for some  $t < 1$ , contradicting  $T \leq \text{Min}$ . Thus  $\varphi$  must be of the form (2.1) for some  $\alpha > 0$ .

Now (0.3) takes the form

$$T(tx, ty) = \Pi^{-1}\{t^\alpha \Pi[T(x, y)]\}. \quad (2.4)$$

For any given  $x, y \in I$ , if  $x \leq y$  we have

$$\Pi \circ T(x, y) = \Pi \circ T\left(y \cdot \frac{x}{y}, y \cdot 1\right) = y^\alpha \Pi \circ T\left(\frac{x}{y}, 1\right) = y^\alpha \Pi\left(\frac{x}{y}\right);$$

while if  $x > y$ , then

$$\Pi \circ T(x, y) = \Pi \circ T\left(x \cdot 1, x \cdot \frac{y}{x}\right) = x^\alpha \Pi \circ T\left(1, \frac{y}{x}\right) = x^\alpha \Pi\left(\frac{y}{x}\right).$$

In either case, putting  $m = \text{Min}(x, y)$  and  $M = \text{Max}(x, y)$ , we obtain

$$T(x, y) = \Pi^{-1}\left\{M^\alpha \Pi\left(\frac{m}{M}\right)\right\}.$$

Since  $\Pi$  is continuous, this shows that  $T$  is continuous and completes the proof of the lemma.

We observe in passing that the full force of the definition of  $t$ -norm was not used in Lemma 2.1. In fact, Lemma 2.1 is valid also whenever  $T: I \times I \rightarrow I$  is quasi-homogeneous, satisfies  $T(1, x) = T(x, 1) = x$ , and the diagonal map  $t \rightarrow T(t, t)$  is non-decreasing and satisfies  $T(t, t) < 1$  for all  $t < 1$ .

Now we may assume that  $T$  is continuous and use Theorem 1.1 to obtain additional information about the structure of  $T$ . The next step is to deal with the Archimedean case.

**THEOREM 2.2.**  $T: I \times I \rightarrow I$  is a quasi-homogeneous  $t$ -norm satisfying  $T(x, x) < x$  for all  $x \in (0, 1)$  if and only if either

$$T(x, y) = xy, \quad (2.5)$$

for all  $x, y \in I$ , with  $\varphi$  given by (2.1) and with  $\Pi(t) = \Pi(1) t^{\frac{\alpha}{2}}$  for some  $\alpha > 0$ , or

$$T(x, y) = \begin{cases} \left(x^\beta + y^\beta - 1\right)^{\frac{1}{\beta}}, & \forall x, y \in (0, 1] \\ 0, & \text{otherwise} \end{cases} \quad (2.6)$$

for some  $\beta < 0$ , with  $\varphi$  given by (2.1) for some  $\alpha > 0$  and with  $\Pi(t) = \Pi(1) \left[\frac{1}{2} \left(t^\beta + 1\right)\right]^{\frac{\alpha}{\beta}}$  if  $t > 0$  and  $\Pi(0) = 0$ .

**PROOF:** Suppose  $T$  is a quasi-homogeneous  $t$ -norm with  $T(x, x) < x$  on  $(0, 1)$ . By Lemma 2.1,  $T$  is continuous and  $\varphi$  has the form (2.1) for some  $\alpha > 0$ . Thus by Theorem 1.1(i),  $T$  has the form (1.1) where  $g$  is a quasi-inverse of  $f$ . Inserting the representation (1.1) into (2.4), we get

$$f[g(tx) + g(ty)] = \Pi^{-1}\{t^\alpha \Pi \circ f[g(x) + g(y)]\}. \quad (2.7)$$

First, we establish that  $g(0) = \infty$  (and hence  $f = g^{-1}$  and  $T$  is strict). Suppose, to the contrary, that  $g(0) < \infty$ . Since  $g$  is continuous, we can choose positive  $t$  so close to 0 that  $g(t) > \frac{1}{2}g(0)$ . Recalling that  $g(1) = 0$ ,  $f(0) = 1$  and  $f(u) = 0$  for  $u \geq g(0)$ , we deduce from (2.7) by putting  $x = y = 1$  that

$$0 = f[2g(t)] = \Pi^{-1}\{t^\alpha \Pi(1)\}$$

for all  $t$  sufficiently close to 0. But this implies  $\Pi(0) = 0 = \Pi(1)$ , contradicting the injectivity of  $\Pi$ . Thus  $g(0) = \infty$  and  $f = g^{-1}$ .

Now we can write (2.7) as

$$g(tx) + g(ty) = g \circ \Pi^{-1}\{t^\alpha \Pi \circ g^{-1}[g(x) + g(y)]\}, \quad (2.8)$$

where  $g$  is a strictly decreasing homeomorphism of  $I$  onto  $\bar{\mathbb{R}}^+$ . Fixing  $t$  temporarily, let  $u = g(x)$ ,  $v = g(y)$  and define

$$g_t(x) := g(tx), \quad k_t(x) := g \circ \Pi^{-1}[t^\alpha \Pi(x)],$$

for  $x \in I$ . Then (2.8) becomes

$$g_t \circ g^{-1}(u) + g_t \circ g^{-1}(v) = k_t \circ g^{-1}(u + v), \quad u, v \in \bar{\mathbb{R}}^+.$$

Since  $g_t$  and  $k_t$  are strictly monotonic by definition, the solution of this Pexider equation is

$$g_t \circ g^{-1}(u) = a_t u + b_t, \quad k_t \circ g^{-1}(u) = a_t u + 2b_t$$

for some “constants”  $a_t$  and  $b_t$  (depending on  $t$ ). Freeing  $t \in I$  and recalling the definitions of  $g_t$  and  $k_t$ , we have now

$$g(tx) = a(t) g(x) + b(t), \quad (2.9)$$

$$g \circ \Pi^{-1}[t^\alpha \Pi(x)] = a(t) g(x) + 2b(t), \quad (2.10)$$

valid for all  $t, x \in I$ .

The general solution of equation (2.9) is given in Theorem 1.2. We eliminate solution (1.4) here because  $g$  is strictly monotonic. We consider solutions (1.3) and (1.5) separately.

**Case 1.** Suppose the solution of (2.9) is of the form (1.3). Since  $g$  is strictly decreasing, so is the logarithmic function  $\ell$ . That is, there exists a constant  $b < 0$  for which

$$g(x) = b \log x + c, \quad a(t) = 1, \quad \text{and} \quad b(t) = \log t. \quad (2.11)$$

Substituting these into (2.10), we find that

$$t^\alpha \Pi(x) = \Pi(xt^2), \quad t, x \in I.$$

With  $x = 1$ , this yields  $\Pi(q) = q^{\alpha/2} \Pi(1)$  for all  $q \in I$ . Furthermore, inserting (2.11) into (1.1), with  $f = g^{-1}$ , we obtain (2.5) for  $T$ .

**Case 2.** Suppose the solution of (2.9) is of the form (1.5). Since  $g$  is strictly monotonic, the same is true of the multiplicative function  $m$ , so  $m(x) = x^\beta$ . Moreover, since  $g(0) = \infty$  and  $g(1) = 0$ , we must have

$$g(x) = c(x^\beta - 1), \quad a(t) = t^\beta, \quad \text{and} \quad b(t) = c(t^\beta - 1) \quad (2.12)$$

for some constants  $\beta < 0$  and  $c > 0$ . Inserting these into (2.10) and simplifying, we arrive at

$$\Pi^{-1}[t^\alpha \Pi(x)] = [(tx)^\beta + t^\beta - 1]^{\beta/2}.$$

Putting  $x = 1$  and  $q = (2t^\beta - 1)^{\beta/2}$ , this gives

$$\Pi(q) = \Pi(1) \left[ \frac{1}{2}(q^\beta + 1) \right]^{\alpha/\beta}.$$

Finally, (2.12) combined with (1.1) (and  $f = g^{-1}$ ) yields

$$T(x, y) = \left\{ 1 + \frac{1}{c} [c(x^\beta - 1) + c(y^\beta - 1)] \right\}^{1/\beta}$$

which simplifies to (2.6).

The converse is easily verified, and that completes the proof of Theorem 2.2.

Now we are ready to establish the main result.

**THEOREM 2.3.**  $T: I \times I \rightarrow I$  is a quasi-homogeneous  $t$ -norm if and only if  $T$  is given by (2.5), by (2.6) for some  $\beta < 0$ , or by  $T(x, y) = \text{Min}(x, y)$ . In the last case,  $\varphi$  is given by (2.1) for some  $\alpha > 0$ , and  $\Pi$  has the form  $\Pi(x) = \Pi(1) x^\alpha$ .

**REMARK.** If we define  $T_\beta$  to be the  $t$ -norm given by (2.6), then

$$\lim_{\beta \rightarrow 0} T_\beta(x, y) = xy, \text{ and } \lim_{\beta \rightarrow -\infty} T_\beta(x, y) = \text{Min}(x, y).$$

Defining  $T_0$  to be the product on  $I \times I$  and  $T_\infty$  to be  $\text{Min}$ , we can restate the conclusion of Theorem 2.3 as follows. The only quasi-homogeneous  $t$ -norms are the members of the family  $\{T_\beta\}$  for  $-\infty \leq \beta \leq 0$ .

**PROOF OF THEOREM 2.3:** Let  $T$  be a quasi-homogeneous  $t$ -norm. Then by Lemma 2.1,  $T$  is continuous and  $\varphi$  has the form (2.1) for some  $\alpha > 0$ . Now we apply Theorem 1.1.

If case (i) of Theorem 1.1 holds, then we find that (2.5) or (2.6) holds by Theorem 2.2.

In case (ii) of Theorem 1.1, we have  $T = \text{Min}$ , and equations (0.3) and (2.1) yield

$$\text{Min}(tx, ty) = \Pi^{-1}\{t^\alpha \Pi \circ \text{Min}(x, y)\}.$$

With  $x = y = 1$ , we obtain

$$\Pi(t) = t^\alpha \Pi(1), \quad t \in I.$$

Finally, we consider case (iii) of Theorem 1.1. Choose any  $x_0 \in (0, 1)$  such that  $T(x_0, x_0) = x_0$ , and let  $S_k$  be any subinterval of  $[x_0, 1]$  such that  $T_k$  admits representation (1.1) in the square  $S_k \times S_k$ . Then

$$T(x, y) = \text{Min}(x, y), \quad (x, y) \in D_1 \cup D_2,$$

where  $D_1$  and  $D_2$  are the rectangles

$$D_1 = \{(x, y) \mid 0 \leq y \leq x_0 \leq x \leq 1\}, \quad D_2 = \{(x, y) \mid 0 \leq x \leq x_0 \leq y \leq 1\}$$

in  $I \times I$ . We prove that in fact this case cannot occur.

Let us confine our attention to the lower triangle  $\{(x, y) \mid 0 < y \leq x \leq 1\}$ . Choose  $(x, y)$  in  $S_k \times S_k$  so that  $x_0 < y \leq x \leq 1$ . Then we choose  $t = \frac{x_0}{y}$ , so that  $(tx, ty)$  belongs to  $D_1$ . Now (0.3) and (2.1) yield

$$\Pi[T(x, y)] = \Pi(x_0) \left( \frac{y}{x_0} \right)^\alpha,$$

which implies that  $T(x, y)$  is independent of  $x$  for  $x \geq y > x_0$ . Specifically,

$$T(x, y) = \Pi^{-1} \left[ \left( \frac{y}{x_0} \right)^\alpha \Pi(x_0) \right] \quad (2.13)$$

when  $x \geq y > x_0$ , which is inconsistent with representation (1.1) for the restriction  $T_k$  of  $T$  to  $S_k \times S_k$ . Indeed, as in the proof of Theorem 2.2,  $T_k$  must be strict:

$$T_k(x, y) = g_k^{-1}[g_k(x) + g_k(y)], \quad x, y \in S_k.$$

This representation, with strictly monotonic  $g_k$ , shows that  $T_k(x, y)$  is injective in  $x$  for each  $y \in S_k$ . But the right hand side of (2.13) is independent of  $x$ . This contradiction shows that there can be no  $S_k$  in the interval  $[x_0, 1]$ .

A similar argument shows that there can be no  $S_k$  in the interval  $[0, x_0]$ . Thus there can be no proper ordinal sum. That is, case (iii) of Theorem 1.1 is incompatible with quasi-homogeneity. (In other words, if a quasi-homogeneous  $T$  satisfies  $T(x_0, x_0) = x_0$  for some  $x \in (0, 1)$ , then it satisfies  $T(x, x) = x$  for all  $x \in (0, 1)$ .)

The converse is easily verified, and this completes the proof of the theorem.

From Theorem 2.3, the structure theorem for homogeneous  $t$ -norms is easily extracted.

**COROLLARY 2.4.**  $T: I \times I \rightarrow I$  is a homogeneous  $t$ -norm if and only if either  $T(x, y) = xy$ , with  $\varphi(t) = t^2$ , or  $T(x, y) = \text{Min}(x, y)$ , with  $\varphi(t) = t$ .

**PROOF:** Suppose  $T$  is a homogeneous  $t$ -norm. We apply Theorem 2.3 in the special case in which  $\Pi$  in (0.3) is the identity map. When  $T$  is of the form (2.5), we have (cf. Theorem 2.2)  $\Pi(x) = \Pi(1)x^{\frac{\alpha}{2}}$ . This will give  $\Pi(x) = x$  only if  $\alpha = 2$ ; in (2.1) this yields  $\varphi(t) = t^2$ . When  $T$  is given by (2.6) for some  $\beta < 0$ , we note that  $\Pi(x) = \Pi(1) \left[ \frac{1}{2}(x^\beta + 1) \right]^{\frac{2}{\beta}}$ . Such  $\Pi$  can never be the identity map, so this case cannot arise. Finally, when  $T(x, y) = \text{Min}(x, y)$ , the accompanying  $\Pi$  is (cf. Theorem 2.3)  $\Pi(x) = \Pi(1)x^\alpha$ . This  $\Pi$  is the identity only if  $\alpha = 1$ , in which case (2.1) becomes  $\varphi(t) = t$ .

The simple converse completes the proof.

#### REFERENCES

- [1] MENGER, K. Statistical metrics, *Proc. Nat. Acad. Sci. USA* **28** (1942), 535-537.
- [2] SCHWEIZER, B. and SKLAR, A. *Probabilistic Metric Spaces*, North-Holland, New York, Amsterdam, Oxford, 1983.
- [3] SCHWEIZER, B. personal communication.
- [4] ACZEL, J. and DHOMBRES, J. *Functional Equations in Several Variables*, Cambridge U. Press, Cambridge, New York, Melbourne, 1989.

## Special Issue on Intelligent Computational Methods for Financial Engineering

### Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Lean Yu**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [yulean@amss.ac.cn](mailto:yulean@amss.ac.cn)

**Shouyang Wang**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; [sywang@amss.ac.cn](mailto:sywang@amss.ac.cn)

**K. K. Lai**, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [mskklai@cityu.edu.hk](mailto:mskklai@cityu.edu.hk)