

RESEARCH NOTES

CODIMENSION 2 FIBRATORS THAT ARE  
CLOSED UNDER FINITE PRODUCT

YOUNG HO IM

Department of Mathematics  
Pusan National University  
Pusan 609-735, KOREA

MEE KWANG KANG

Department of Mathematics  
Dongeui University  
Pusan 614-714, KOREA

KI MUN WOO

Department of Mathematics  
Pusan National University  
Pusan 609-735, KOREA

(Received March 14, 1996 and in revised form June 5, 1996)

**ABSTRACT.** In this paper, we show that if  $N^m$  is a closed manifold with hyperhopfian fundamental group,  $\pi_i(N) = 0$  for  $1 < i \leq n$  and  $S^n$  is a simply connected manifold, then  $N^m \times S^n$  satisfies the property that all proper, surjective maps from an orientable  $(n+2)$ -manifold  $M$  to a 2-manifold  $B$  for which each  $p^{-1}(b)$  is homotopy equivalent to  $N^m \times S^n$  necessarily are approximate fibrations.

**KEY WORDS AND PHRASES:** Approximate fibration, Hopfian group, hyperhopfian group, Hopfian manifold.

**1991 AMS SUBJECT CLASSIFICATION CODES:** 57N15; 55R65.

1. INTRODUCTION

In the study of proper maps between manifolds, the concepts of approximate fibrations play very important roles because it has nice and useful properties just as Hurewicz fibrations. As a generalization of Hurewicz fibrations and cell-like maps, the concept of an approximate fibration is initially introduced by Coram and Duvall [1].

A proper map  $p : M \rightarrow B$  between locally compact ANR's is called an *approximate fibration* if it has the following approximate homotopy lifting property for all spaces: given an open cover  $\epsilon$  of  $B$ , an arbitrary space  $X$ , and two maps  $g : X \rightarrow M$  and  $X \times I \rightarrow b$  such that  $p \circ g = F_0$ , there exists a map  $G : X \times I \rightarrow M$  such that  $G_0 = g$  and  $p \circ G$  is  $\epsilon$ -close to  $F$ .

If a proper map  $p : M \rightarrow B$  is an approximate fibration, then the point inverses are homotopy equivalent to each other. Naturally, the question arises as to under what conditions the converse of this fact holds. Many people have been interested in a setting which forces any proper map defined on an arbitrary manifold of a specified dimension to be an approximate fibration merely due to the fact that all point preimages are copies of some closed manifold  $N$ . In this paper, we mainly concentrate on such a closed  $n$ -manifold when the dimension of  $M$  is particularly  $n+2$ .

We assume all spaces are locally compact, metrizable ANR's, and all manifolds are finite dimensional, orientable, connected and boundaryless. A manifold  $M$  is said to be *closed* if  $M$  is compact, connected and boundaryless. A closed  $n$ -manifold  $N$  is called a *codimension 2 fibrator* if, whenever  $p : M \rightarrow B$  is a proper map from an arbitrary  $(n+2)$ -manifold  $M$  to a 2-manifold  $B$  such that each point preimage is homotopy equivalent to  $N$ ,  $p : M \rightarrow B$  is an approximate fibration.

The *degree* of a map  $R : N \rightarrow N$ , where  $N$  is a closed manifold, is the nonnegative integer  $d$  such that the induced endomorphism of  $H_n(N; \mathbb{Z}) \cong \mathbb{Z}$  amounts to multiplication by  $d$ , up to sign. Note that

a degree one map  $R : N \rightarrow N$  induces homology isomorphisms  $R_* : H_*(N) \rightarrow H_*(N)$  for all integers  $i \geq 0$ . The *continuity set*  $C$  of  $p : M \rightarrow B$  consists of those points  $c \in B$  such that, under any retraction  $R : p^{-1}U \rightarrow p^{-1}c$  defined over a neighborhood  $U \subset B$  of  $c$ ,  $c$  has another neighborhood  $V_c \subset U$  such that  $R|p^{-1}b : p^{-1}b \rightarrow p^{-1}c$  is a degree one map for all  $b \in V_c$ . The continuity set  $C$  is the maximal open subset of  $B$  over which the  $n$ th cohomology sheaf of the map  $p$  is locally constant. In [2], several characterizations for an approximate fibration were described. As a matter of fact, the essential point whether or not a proper map is an approximate fibration depends on the fact that any retraction  $R : p^{-1}U \rightarrow p^{-1}b$  restricts to homotopy equivalences  $p^{-1}c \rightarrow p^{-1}b$  for all  $c$  sufficiently close to  $b$  [2]. The following terms efficiently aid to convert a homology equivalence into a homotopy equivalence.

A closed manifold  $N$  is called *Hopfian* if every degree one map  $N \rightarrow N$  which induces a  $\pi_1$ -automorphism is a homotopy equivalence. A group  $H$  is *Hopfian* if every epimorphism  $\Psi : H \rightarrow H$  is necessarily an isomorphism, while a finitely presented group  $H$  is *hyperhopfian* if every endomorphism  $\Psi : H \rightarrow H$  with  $\Psi(H)$  normal and  $H/\Psi(H)$  cyclic is an automorphism. A hyperhopfian group implies a Hopfian one but the converse does not hold.

Daverman ([3],[4]) has shown that each of a simply connected closed manifold and an aspherical closed manifold with hyperhopfian fundamental group is a codimension 2 fibrator. The problem whether the class of codimension 2 fibrators is closed under finite product is not yet settled. A particular class of all closed surfaces with negative Euler characteristics is closed, which is claimed by Im ([5]). Also we extended this result to the extent that any product of an  $n$ -sphere  $S^n$  ( $n > 1$ ) and a finite product of closed surfaces of genus at least 2 is a codimension 2 fibrator ([6]).

The purpose of this paper is to extend the above results so that any finite product of a simply connected closed manifold  $S^n$  and a closed manifold  $N^m$  with hyperhopfian fundamental group and  $\pi_i(N) = 0$  for  $1 < i \leq n$  is a codimension 2 fibrator.

## 2. PRELIMINARIES

The investigation about codimension 2 fibrators compared with other codimensions is easier to approach due to the following result.

**LEMMA 2.1** [6]. If  $G$  is an upper semicontinuous decomposition of an  $(n+2)$ -manifold  $M$  into closed  $n$ -manifolds, then the decomposition space  $B (= M/G)$  is a 2-manifold and  $D = B \setminus C$  is locally finite in  $B$ , where  $C$  represents the continuity set of the decomposition map  $p : M \rightarrow B$ .

Because of the local finiteness of  $D$ , we can localize the problem to that of an open disk  $B$ , proved that  $p$  is an approximate fibration over the continuity set  $C$ , so that  $p : M \rightarrow B$  is an approximate fibration over  $B - b$  for some  $b \in B$ . In such a case, the homotopy exact sequence for  $p$  over  $B \setminus b$  can be reduced to a short exact sequence as follows:

$$0 \simeq \pi_2(B - b) \rightarrow \pi_1(p^{-1}(c)) \rightarrow \pi_1(M - p^{-1}(b)) \rightarrow \pi_1(B - b) \simeq \mathbb{Z} \rightarrow 0,$$

where  $c$  is any point of  $B \setminus b$ . Hence, if each preimage of  $p$  has the same homotopy type as  $N$ , the hyperhopfian property of  $\pi_1(N)$  makes the continuity set  $C$  the whole set  $B$ . And so, the hyperhopfian property is very valuable on discussing codimension 2 fibrators.

We state the established results about codimension 2 fibrators.

**THEOREM 2.2** [4]. All closed, Hopfian manifolds with hyperhopfian fundamental group is a codimension 2 fibrator.

**THEOREM 2.3** [3]. Every simply connected closed manifold is a codimension 2 fibrator.

The idea of the proof is to show that the continuity set of the decomposition map  $p : M^{n+2} \rightarrow M/G$  is the entire set  $M/G$ .

A closed surface with negative Euler characteristic has a hyperhopfian fundamental group [4]. Therefore, from Theorem 2.2 and 2.3, the following corollary follows

**COROLLARY 2.4.** Every closed surface  $F^2$  with nonzero Euler characteristic is a codimension 2 fibrator.

**THEOREM 2.5** [5]. Any finite product  $N = F_1 \times F_2 \times \cdots \times F_m$  of closed surfaces  $F_i^2$  ( $i = 1, \dots, m$ ) of genus at least 2 is a codimension 2 fibrator.

Since an  $n$ -sphere  $S^n$  is a simply connected closed manifold, each of the fundamental group and the first homology group of  $S^n \times F_1 \times \cdots \times F_m$  is isomorphic to each of those of  $F_1 \times \cdots \times F_m$ . Taking advantage of the method in the proof of Theorem 2.5 and the fact that  $S^n \times F_1 \times \cdots \times F_m$  is a Hopfian manifold, we can obtain the next consequence.

**THEOREM 2.6.** A finite product  $N = S^n \times F_1 \times \cdots \times F_m$  of  $n$ -sphere  $S^n$  ( $n > 1$ ) and closed orientable surfaces  $F_i^2$  ( $i = 1, \dots, m$ ) with negative Euler characteristics is a codimension 2 fibrator.

### 3. MAIN RESULTS

**PROPOSITION 3.1.** Let  $S$  be a simply connected closed  $n$ -manifold and  $N$  be a closed  $m$ -manifold with  $\pi_i(N) = 0$  for  $1 < i \leq n$ . If  $R : S \times N \rightarrow S \times N$  is a degree one map, then so is  $pr \circ R \circ i : S \rightarrow S$ , where  $pr : S \times N \rightarrow S$  is the first projection and  $i : S \rightarrow S \times N$  is an inclusion map.

**PROOF.** Assume that  $R : S \times N \rightarrow S \times N$  is a degree one map. By taking a universal covering space  $(\tilde{N}, \theta)$  of  $N$ ,  $(S \times \tilde{N}, id \times \theta)$  is a universal covering space of  $S \times N$ .

Let  $\tilde{i} : S \rightarrow S \times \tilde{N}$  be a continuous map for which  $(id \times \theta) \circ \tilde{i} = i$  and  $\tilde{R} : S \times \tilde{N} \rightarrow S \times \tilde{N}$  be a continuous map such that  $R \circ (id \times \theta) = (id \times \theta) \circ \tilde{R}$  by the lifting property. Consider the following commutative diagram

$$\begin{array}{ccccccc} S & \xrightarrow{\tilde{i}} & S \times \tilde{N} & \xrightarrow{\tilde{R}} & S \times \tilde{N} & \xrightarrow{q} & S \\ \downarrow id & & \downarrow id \times \theta & & \downarrow id \times \theta & & \downarrow id \\ S & \xrightarrow{i} & S \times \tilde{N} & \xrightarrow{R} & S \times \tilde{N} & \xrightarrow{pr} & S \end{array}$$

where  $q$  is the projection from  $S \times \tilde{N}$  onto  $S$ . Because of  $\pi_i(N) = 0$  for  $1 < i \leq n$ , we obtain that  $\pi_i(\tilde{N}) = 0$  for  $1 \leq i \leq n$ , and then  $H_i(\tilde{N}) = 0$  for  $1 \leq i \leq n$ . According to the Künneth Theorem,  $H_n(S \times \tilde{N}) \cong H_n(S)$  and  $H_n(S \times N)$  is isomorphic to the direct sum of  $\bigoplus_{i=0}^n H_{n-i}(S) \otimes H_i(N)$  and  $\bigoplus_{i=0}^{n-1} H_{n-i-1}(S) * H_i(N)$ . Then it easily checked that  $i_* \circ q_* = (id \times \theta)_*$  as homomorphisms from  $H_n(S \times N)$  to itself, and by the diagram chasing,  $R_*(H_n(S)) \subset H_n(S)$  holds when we restrict  $R_*$  to  $H_n(S) \subset H_n(S \times N)$ .

Rewrite  $H_n(S \times N)$  in a form  $\{\text{torsion-free}\} \oplus \{\text{torsion}\}$ . Since  $R_*$  is an isomorphism, the restriction of  $R_*$  to  $\{\text{torsion-free}\}$  of  $H_n(S \times N)$  is an isomorphism and induces an invertible  $k \times k$  matrix of the following form

$$K = \begin{pmatrix} K_{11} & K_{12} & \dots & K_{1k} \\ K_{21} & K_{22} & \dots & K_{2k} \\ \dots & \dots & \dots & \dots \\ K_{k1} & K_{k2} & \dots & K_{kk} \end{pmatrix}.$$

Here  $K_{11}$  is the matrix corresponding to the map  $R_*|H_n(S) : H_n(S) \rightarrow H_n(S)$  and  $K_{ij}$  is the matrix induced by the homomorphism from the  $i$ -th direct summand to the  $j$ -th direct summand of torsion-free part of  $H_n(S \times N)$ .

Since  $R_*(H_n(S)) \subset H_n(S)$ , the restriction  $R_*| \{\text{torsion-free}\}$  of  $R_*$  doesn't send first factor  $H_n(S)$  to any direct summand except itself and thus  $K_{1j}$  is zero for each  $j = 2, \dots, k$ . Hence, the isomorphism  $R_*| \{\text{torsion-free}\}$  induces  $\det K = \pm 1$  and  $\det K_{11} = \pm 1$ , so that  $pr \circ R \circ i$  is a degree one map.

**COROLLARY 3.2.** Let  $S$  be a simply connected closed  $n$ -manifold and  $N$  be a closed aspherical  $m$ -manifold. If  $R : S \times N \rightarrow S \times N$  is a degree one map, then so is  $pr \circ R \circ i : S \rightarrow S$ , where  $pr : S \times N \rightarrow S$  is the first projection and  $i : S \rightarrow S \times N$  is an inclusion map.

**COROLLARY 3.3.** Let  $S$  be a simply connected closed  $n$ -manifold and  $N$  be a finite product of closed orientable surfaces with nonpositive Euler characteristics. Then for every degree one map  $R : S \times N \rightarrow S \times N$ ,  $pr \circ R \circ i : S \rightarrow S$  is a degree one map.

**PROOF.** Every closed orientable surface except 2-sphere is aspherical and their finite product is a closed aspherical manifold.

**REMARK.** In the above corollary, a degree one map  $R : S_1 \times S_2 \rightarrow S_1 \times S_2$  on a product of closed simply connected manifolds  $S_1$  and  $S_2$  cannot guarantee that the degree of the restriction  $R|S_1$  is one, so that the condition nonpositive Euler characteristic cannot be omitted.

The following theorem is the main result in this section.

**THEOREM 3.4.** Let  $S$  be a closed simply connected manifold, and  $N$  be a closed manifold with hyperhopfian fundamental group and  $\pi_i(N) = 0$  for  $1 < i \leq n$ . Then  $S \times N$  is a codimension 2 fibrator.

**PROOF.** Since the fundamental group of  $S \times N$  is isomorphic to the hyperhopfian group  $\pi_1(N)$ , it suffices to show that  $S \times N$  is a Hopfian manifold by means of Theorem 2.2. Assume that  $R : S \times N \rightarrow S \times N$  is a degree one map. Since a degree one map between compact orientable manifolds of the same dimension induces a  $\pi_1$ -epimorphism ([8]),  $R_\# : \pi_1(S \times N) \rightarrow \pi_1(S \times N)$  is an epimorphism and actually it is a  $\pi_1$ -isomorphism by the Hopfian property of  $\pi_1(N)$ .

To claim that  $R$  is a homotopy equivalence, let us consider  $R_\# : \pi_i(S \times N) \rightarrow \pi_i(S \times N)$  for  $i \geq 2$ . By Proposition 3.1, the degree of  $pr \circ R \circ i : S \rightarrow S$  is one and so  $pr_* \circ R_* \circ i_* : H_i(S) \rightarrow H_i(S)$  is an isomorphism for each  $i \geq 1$ . Since  $S$  is simply connected, we can apply the Whitehead theorem and obtain that  $(pr \circ R \circ i)_\# : \pi_i(S) \rightarrow \pi_i(S)$  is a  $\pi_i$ -isomorphism for  $i \geq 1$ . Implying the fact that  $\pi_i(S \times N) \simeq \pi_i(S) \times \pi_i(N) \simeq \pi_i(S)$  for each  $i \geq n$ ,  $R_\# : \pi_i(S \times N) \rightarrow \pi_i(S \times N)$  is an isomorphism for each  $i \geq n$ . Since  $R$  has the property  $R \simeq$ , we obtain that  $R$  is a homotopy equivalence. Therefore,  $S \times N$  is a Hopfian manifold.

**COROLLARY 3.5.** Let  $S$  be a closed simply connected manifold, and  $N$  be a closed aspherical manifold with hyperhopfian fundamental group. Then  $S \times N$  is a codimension 2 fibrator.

**COROLLARY 3.6.** Let  $S$  be a closed simply connected manifold, and  $F$  be a closed surface with nonzero Euler characteristic. Then  $S \times F$  is a codimension 2 fibrator.

**PROOF.** If the Euler characteristic of  $F$  is negative, its fundamental group is hyperhopfian [4]. Otherwise,  $F$  is homeomorphic to 2-sphere and then  $S \times F$  is simply connected. Hence, it is a codimension 2 fibrator.

**COROLLARY 3.7.** Let  $S$  be a closed simply connected manifold, and  $N$  be a finite product of closed surfaces with nonzero Euler characteristic. Then  $S \times N$  is a codimension 2 fibrator.

**PROOF.** If a closed surface  $F$  has a positive Euler characteristic, then  $F$  is a 2-sphere. Therefore, we can rewrite  $S \times N$  as  $S' \times N'$ , where  $S'$  is simply connected and  $N'$  is a product of closed surfaces with negative Euler characteristic. Since  $N'$  is a closed aspherical manifold with hyperhopfian fundamental group [4],  $S \times N$  is a codimension 2 fibrator.

**ACKNOWLEDGEMENT.** The present studies were supported by the Basic Science Research Institute Program, Ministry of Education, 1996, Project No. BSRI-96-1433.

## REFERENCES

- [1] CORAM, D. and DUVALL, P., Approximate fibrations, *Rocky Mountain J. Math.* **7** (1977), 275-288.
- [2] CORAM, D. and DUVALL, P., Approximate fibrations and a movability condition for maps, *Pacific J. Math.* **72** (1977), 41-56.
- [3] DAVERMAN, R.J., Submanifold decompositions that induce approximate fibrations, *Top. Appl.* **33** (1989), 173-184.
- [4] DAVERMAN, R.J., Hyperhopfian groups and approximate fibrations, *Compositio Math.* **86** (1993), 159-176.
- [5] IM, Y.H., Decompositions into codimension two submanifolds that induce approximate fibrations, *Topology Appl.* **56** (1994), 1-11.
- [6] IM, Y.H., KANG, M.K. and WOO, K.M., Product spaces that induce approximate fibrations, *J. Korean Math. Soc.* **33** (1996), 145-154.
- [7] DAVERMAN, R.J. and WALSH, J.J., Decompositions into codimension two manifolds, *Trans. Amer. Math. Soc.* **288** (1985), 273-291.
- [8] HEMPEL, J., *3-Manifolds*, Princeton University Press, Princeton, NJ, 1976.

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)