

FIXED POINTS OF ROTATIONS OF n -SPHERE

NAGABHUSHANA PRABHU

(Received 18 May 1992)

ABSTRACT. We show that every rotation of an even-dimensional sphere must have a fixed point.

Keywords and phrases. Fixed point, eigenvalue.

1991 Mathematics Subject Classification. 51F10, 51F25, 51M04, 15A18.

The curious “Hairy Ball Theorem” [1] states that *there are no continuous nonvanishing vector fields tangent to the $2k$ -dimensional sphere S^{2k}* . Hairy Ball Theorem, however, is false for S^{2k-1} (easy to verify), which shows that one can geometrically determine the parity of n in S^n .

Here is another geometric and simpler asymmetry between spheres of odd and even dimensions:

THEOREM 1. *Every rotation of S^{2n} has at least one fixed point.*

Once again, as an example below illustrates, one can construct rotations of S^{2n-1} that have no fixed point.

PROOF. Rotation in \mathbb{R}^k is a linear transformation that preserves distance from the origin. Thus, if A denotes the transformation matrix, then for every $x \in \mathbb{R}^k$,

$$x^T x = (Ax)^T Ax = x^T A^T Ax, \quad (1)$$

which implies that $A^T A = I$ or $A^{-1} = A^T$ (i.e., A is an orthogonal matrix). $A^{-1} = A^T$ implies that $\det(A) = \pm 1$. But rotation is a continuous transformation and hence one can find a continuous chain of matrices $M(t)$ such that $M(0) = I$ and $M(1) = A$ and each $M(t)$, $0 \leq t < 1$, represents a rotation. $f(t) = \det(M(t))$ is a continuous function of t with $f(0) = 1$. If $f(1) = -1$, by intermediate value theorem $f(t') = 0$ for $0 < t' < 1$, which contradicts the assumption that $M(t')$ represents a rotation and is therefore nonsingular. Hence, $\det(A) = +1$ (orthogonal matrices with negative determinant represent reflection). $S^{2n} \subset \mathbb{R}^{2n+1}$. Hence, if A represents a rotation in \mathbb{R}^{2n+1} , then A is an order $2n+1$ matrix. The characteristic polynomial $P(x) = \det(A - xI)$ is hence of degree $2n+1$. Complex roots of $P(x)$ (if any) occur in conjugate pairs. Hence, $P(x)$ has at least one real root. Further, since the determinant of A is the product of its eigenvalues, the product of the roots of $P(x)$ equals $+1$. The product of a pair of complex conjugates is always nonnegative and hence A must have an even number of negative eigenvalues (counting multiplicity). Since $P(x)$ has $2n+1$ roots in all (counting multiplicity), it has at least one positive eigenvalue, say λ ; the eigenvector y of λ

is real.

$$(Ay)^T Ay = \lambda^2 y^T y = y^T y, \quad (2)$$

which implies that $\lambda = +1$ and $Ay = y$. Hence, y is a fixed point of the rotation A . \square

Next, consider the following rotation of $S^{2n-1} \subset \mathbb{R}^{2n}$

$$B(\phi_1, \dots, \phi_n) = \begin{bmatrix} \cos \phi_1 & -\sin \phi_1 & & & \\ \sin \phi_1 & \cos \phi_1 & & & \\ & & \ddots & & \\ & & & \cos \phi_n & -\sin \phi_n \\ & & & \sin \phi_n & \cos \phi_n \end{bmatrix} \quad (3)$$

with $0 < \phi_1, \dots, \phi_n < \pi/2$. The eigenvalues of B are $e^{\pm i\phi_1}, \dots, e^{\pm i\phi_n}$, none of which is real for $0 < \phi_1, \dots, \phi_n < \pi/2$. Since $+1$ is not an eigenvalue of B , the rotation B cannot have any fixed points.

REFERENCES

[1] J. Milnor, *Analytic proofs of the “hairy ball theorem” and the Brouwer fixed-point theorem*, Amer. Math. Monthly **85** (1978), 521–524. MR 80m:55001. Zbl 386.55001.

PRABHU: PURDUE UNIVERSITY, GRISSOM HALL, WEST LAFAYETTE, IN 47907, USA

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk