

## SUBSEQUENCES AND CATEGORY

ROBERT R. KALLMAN

(Received 21 July 1998)

**ABSTRACT.** If a sequence of functions diverges almost everywhere, then the set of subsequences which diverge almost everywhere is a residual set of subsequences.

**Keywords and phrases.** Baire category, measure, measurable functions, sequences.

1991 Mathematics Subject Classification. 28A20, 54E52.

**1. Introduction.** In [1], Bilyeu, Lewis, and Kallman proved a general theorem about rearrangements of a series of Banach space valued functions. This theorem settled a question on rearrangements of Fourier series posed by Kac and Zygmund. Kallman [3] proved an analog of this theorem for subseries of a series of Banach space valued functions. The purpose of this paper is to complete the cycle of these ideas by proving an analogous theorem (Theorem 1.1) for subsequences of a sequence of Banach space valued functions. Theorem 1.1 does not seem to follow directly from results of [1] or [3]. Other than [1, 3], the only precedent for Theorem 1.1 seems to be a paper [7] on subsequences of a sequence of complex numbers.

Let  $S$  be the set of all sequences  $s = (s_1, s_2, \dots)$ , where  $1 \leq s_1 < s_2 < \dots$  is a strictly increasing sequence of positive integers.  $S$  is a closed subset of the countable product of the positive integers, and so  $S$  is a complete separable metric space. Given any sequence of objects  $a_1, a_2, \dots$ , one can identify the set of its subsequences both as a set and as a topological space with  $S$ . In this context, it is natural to identify a collection of subsequences with a subset of  $S$  and ask if it is first category, second category, or residual ([5] or [6]). Define an equivalence relation  $\sim$  on  $S$  as follows: if  $s, t \in S$ , then  $s \sim t$  if and only if  $s_n = t_n$  for all sufficiently large  $n$ . Intuitively this states that  $s \sim t$  if and only if  $s$  and  $t$  agree from some point on. It is simple to check that any nonempty subset of  $S$  which is saturated with respect to  $\sim$  is dense.

The main result of this paper is the following theorem, which is proved in Section 2.

**THEOREM 1.1.** *Let  $(X, \mu)$  be a regular locally compact  $\sigma$ -finite measure space,  $Z$  a separable Banach space, and  $f_n : X \rightarrow Z$  a sequence of Borel measurable functions. Suppose that the sequence  $f_n(x)$  diverges for  $\mu$ -a.e.,  $x \in X$ . Then  $[s \in S \mid f_{s_n}(x) \text{ diverges for } \mu\text{-a.e. } x \in X]$  is a residual set in  $S$ .*

Just as in [1, 3], this measure-category result has a category-category analog which is discussed in Section 3.

**2. Proof of Theorem 1.1.** The following special case of Theorem 1.1 will be proved first.

**LEMMA 2.1.** *Let  $K$  be a compact Hausdorff space,  $Z$  a Banach space, and  $f_n : K \rightarrow Z$  a sequence of continuous functions, and  $\delta > 0$ . Suppose that for every  $x \in K$  and positive integer  $N$ , there exists a pair of integers  $n = n(x, N)$  and  $m = m(x, N)$  so that  $N \leq n \leq m$  and  $\|f_m(x) - f_n(x)\| > \delta$ . Then  $[s \in S \mid f_{s_n}(x) \text{ diverges for every } x \in K]$  is a residual set in  $S$ .*

**PROOF.** If  $m, n$  is a pair of integers such that  $1 \leq n \leq m$  and  $s \in S$ , let  $g_{s,m,n} : K \rightarrow [0, +\infty)$  be defined by  $g_{s,m,n}(x) = \|f_{s_m}(x) - f_{s_n}(x)\|$ .  $g_{s,m,n}$  is continuous. Consider

$$A = \bigcap_{N \geq 1} \bigcup_{N \leq n_1 \leq m_1, \dots, N \leq n_p \leq m_p} \left[ s \in S \mid \cup_{1 \leq i \leq p} g_{s,m_i,n_i}^{-1}((\delta, +\infty)) = K \right]. \quad (2.1)$$

Fix  $1 \leq n \leq m$  and  $s \in S$ . Then  $V = [t \in S \mid t_m = s_m \text{ and } t_n = s_n]$  is an open neighborhood of  $s$  in  $S$ . Hence, if  $t \in V$ , then  $g_{t,m,n} = g_{s,m,n}$ . This in turn implies that  $A$  is a  $G_\delta$  subset of  $S$ . Furthermore,  $A$  is saturated with respect to the equivalence relation  $\sim$  and therefore is a dense  $G_\delta$  if it is nonempty.

$A$  is nonempty since  $t = (1, 2, 3, \dots)$  is in  $A$ . To see this, fix  $N \geq 1$ . For  $N \leq n \leq m$ , let  $U(m, n) = g_{t,m,n}^{-1}((\delta, +\infty))$ . Note that the collection  $\{U(m, n)\}_{N \leq n \leq m}$  is an open covering of  $K$  by hypothesis and so has a finite subcover, say  $U(m_1, n_1), \dots, U(m_p, n_p)$ . One easily concludes from this that  $t \in A$ .

Finally, note that the Cauchy criterion for convergence implies that if  $s \in A$ , then  $f_{s_n}(x)$  diverges for every  $x \in K$ . Hence,  $A \subseteq [s \in S \mid f_{s_n}(x) \text{ diverges for every } x \in K]$ . This proves Lemma 2.1.  $\square$

**PROOF OF Theorem 1.1.** We may assume that  $\mu$  is a probability measure since  $\mu$  is  $\sigma$ -finite. If  $q \geq 1$ , let

$$D_q = \bigcap_{N \geq 1} \bigcup_{N \leq n \leq m} \left[ x \in X \mid \|f_m(x) - f_n(x)\| > \frac{1}{q} \right]. \quad (2.2)$$

Each  $D_q$  is a Borel subset of  $X$ ,  $D_q \subseteq D_{q+1}$ , and the Cauchy criterion for convergence implies that  $\cup_{q \geq 1} D_q = [x \in X \mid f_n(x) \text{ diverges}]$ .  $\mu(\cup_{q \geq 1} D_q) = 1$  by assumption. Use a vector-valued version of Lusin's Theorem [2] to choose, for each  $q$ , a compact subset  $K_q$  of  $D_q$  so that each  $f_n \mid K_q$  is continuous and  $\mu(D_q - K_q) < 1/q$ .  $R_q = [s \in S \mid f_{s_n}(x) \text{ diverges for every } x \in K_q]$  is a residual subset of  $S$  by Lemma 2.1. Hence,  $R = \cap_{q \geq 1} R_q$  is a residual set in  $S$  and is contained in  $[s \in S \mid f_{s_n}(x) \text{ diverges for } \mu\text{-a.e., } x \in X]$  since  $\mu(\cup_{q \geq 1} K_q) = 1$ . This proves Theorem 1.1.  $\square$

**3. Sequences of functions with the Baire property.** Theorem 1.1 may be regarded as a measure-category result. The purpose of this section is to prove a category-category analog of Theorem 1.1 (cf. [1, Thm. 1.2] and [3, Thm. 3.1]).

Let  $X$  be a Polish space. A subset of  $X$  is said to have the Baire property if there exists an open set  $U$  in  $X$  so that  $A \triangle U$  is first category. The collection of all subsets of  $X$  with the Baire property is a  $\sigma$ -algebra which includes the analytic sets in  $X$ . Let  $Z$  be any other Polish space. A function  $f : X \rightarrow Z$  is said to have the Baire property if  $U$  open in  $Z$  implies that  $f^{-1}(U)$  has the Baire property in  $X$ . Any Borel function  $f : X \rightarrow Z$  is a function with the Baire property. See [4, 5] or [6] for a thorough discussion of this circle of ideas. The following theorem is then a category-category analog of Theorem 1.1.

**THEOREM 3.1.** *Let  $X$  be a Polish space,  $Z$  a separable Banach space, and  $f_n : X \rightarrow Z$  a sequence of functions with the Baire property. Suppose that  $[x \in X \mid f_n(x) \text{ diverges}]$  is a residual subset of  $X$ . Then  $[s \in S \mid f_{s_n}(x) \text{ diverges on a residual subset of } X]$  is a residual subset of  $S$ .*

The following proposition, of independent interest, is needed to prove Theorem 3.1.

**PROPOSITION 3.2.** *Let  $Z$  be a Banach space and let  $\{z_n\}_{n \geq 1}$  be a sequence in  $Z$ . Let  $A = [s \in S \mid z_{s_n} \text{ converges}]$ . Then either  $A = S$  or  $A$  is of first category in  $S$ .*

**PROOF.** For  $k \geq 1$  define

$$B_k = \bigcap_{N \geq 1} \bigcup_{N \leq n \leq m} \left[ s \in S \mid \|z_{s_m} - z_{s_n}\| > \frac{1}{k} \right]. \quad (3.1)$$

Note that  $B_k \subseteq B_{k+1}$ . Each set in square brackets is open in  $S$ . Hence, this formula shows that  $B_k$  is a  $G_\delta$ .  $B_k$  is dense if it is nonempty since it is saturated with respect to the equivalence relation  $\sim$ . Therefore,  $B_k$  is a residual set in  $S$  if it is nonempty since any dense  $G_\delta$  is residual.

The Cauchy criterion for convergence implies that  $A^c = \bigcup_{k \geq 1} B_k$ . Hence, either  $A = S$  or  $A^c$  is residual in  $S$ ; or either  $A = S$  or  $A$  is of first category in  $S$ . This proves Proposition 3.2.  $\square$

**PROOF OF Theorem 3.1.** Check that the mapping  $(x, s) \mapsto f_{s_n}(x)$ ,  $X \times S \rightarrow Z$ , is a function with the Baire property for every  $n \geq 1$ . Hence,

$$B = [(x, s) \mid f_{s_n}(x) \text{ diverges}] = \bigcup_{k \geq 1} \bigcap_{N \geq 1} \bigcup_{N \leq n \leq m} \left[ (x, s) \mid \|f_{s_m}(x) - f_{s_n}(x)\| > \frac{1}{k} \right] \quad (3.2)$$

is a subset of  $X \times S$  with the Baire property. For each  $x \in X$ , let  $B_x^c$  be the projection of  $B^c \cap ((x) \times S)$  onto  $S$ . The hypotheses of Theorem 3.1 plus Proposition 3.2 imply that each  $B_x^c$  is a first category subset of  $S$ , except for a first category set of  $x$ 's. But then  $B^c$  is itself a first category subset of  $X \times S$  [6, Thm. 15.4] and so  $B_S^c$ , the projection of  $B^c \cap (X \times (s))$  onto  $X$ , is a first category subset of  $X$ , except for a first category set of  $s$ 's (Theorem of Kuratowski-Ulam, [6, Thm. 15.1]). Hence,  $B_s$ , the projection of  $B \cap (X \times (s))$  onto  $X$ , is a residual subset of  $X$  for all except a first category set of  $s$ 's. This proves Theorem 3.1.  $\square$

## REFERENCES

- [1] R. G. Bilyeu, R. R. Kallman, and P. W. Lewis, *Rearrangements and category*, Pacific J. Math. **121** (1986), no. 1, 41–46. MR 87j:42019. Zbl 574.42007.
- [2] J. Diestel and J. J. Uhl, Jr., *Vector measures*, Mathematical Surveys, no. 15, American Mathematical Society, Providence, RI, 1977. MR 56 12216. Zbl 369.46039.
- [3] R. R. Kallman, *Subseries and category*, J. Math. Anal. Appl. **132** (1988), no. 1, 234–237. MR 89g:46074. Zbl 676.46004.
- [4] A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995. MR 96e:03057. Zbl 819.04002.
- [5] K. Kuratowski, *Topology. Vol. I.*, Academic Press, New York, London, 1966. MR 36#840. Zbl 158.40802.
- [6] J. C. Oxtoby, *Measure and category*, Graduate Texts in Mathematics, vol. 2, Springer-Verlag, New York, Berlin, 1980. MR 81j:28003. Zbl 435.28011.

[7] M. B. Rao, K. P. S. B. Rao, and B. V. Rao, *Remarks on subsequences, subseries and rearrangements*, Proc. Amer. Math. Soc. **67** (1977), no. 2, 293–296. MR 58 12068. Zbl 412.40001.

KALLMAN: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, P. O. BOX 305118, DENTON, TEXAS 76203-5118, USA

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)