

A NOTE ON THE COUNTABLE UNION OF PRIME SUBMODULES

M. R. POURNAKI and M. TOUSI

(Received 11 November 2000)

ABSTRACT. Let M be a finitely-generated module over a Noetherian ring R . Suppose \mathfrak{a} is an ideal of R and let $N = \mathfrak{a}M$ and $\mathfrak{b} = \text{Ann}(M/N)$. If $\mathfrak{b} \subseteq J(R)$, M is complete with respect to the \mathfrak{b} -adic topology, $\{P_i\}_{i \geq 1}$ is a countable family of prime submodules of M , and $x \in M$, then $x + N \subseteq \bigcup_{i \geq 1} P_i$ implies that $x + N \subseteq P_j$ for some $j \geq 1$. This extends a theorem of Sharp and Vámos concerning prime ideals to prime submodules.

2000 Mathematics Subject Classification. 13C99, 13J10, 13E05.

1. Introduction. Let R be a commutative ring with identity. One of the fundamental cornerstones of commutative ring theory is the “prime avoidance” theorem, which states that, if $\mathfrak{p}_1, \dots, \mathfrak{p}_n$ are prime ideals of R and \mathfrak{a} is an ideal of R such that $\mathfrak{a} \subseteq \bigcup_{i=1}^n \mathfrak{p}_i$, then $\mathfrak{a} \subseteq \mathfrak{p}_j$ for some $1 \leq j \leq n$. In [4], the authors proved and used an extension of the following.

THEOREM 1.1. *Let R be a Noetherian local ring having maximal ideal \mathfrak{m} . Let $\{\mathfrak{p}_i\}_{i \geq 1}$ be a countable family of prime ideals of R , \mathfrak{a} an ideal of R , and $x \in R$. If R is complete with respect to the \mathfrak{m} -adic topology, then $x + \mathfrak{a} \subseteq \bigcup_{i \geq 1} \mathfrak{p}_i$ implies that $x + \mathfrak{a} \subseteq \mathfrak{p}_j$ for some $j \geq 1$.*

Consider M as a unitary left R -module. A submodule P of M is called *prime submodule*, if $P \neq M$ and for each $r \in R$ and $m \in M$, $rm \in P$ implies that $r \in \text{Ann}(M/P)$ or $m \in P$. Of course, if we regard R as an R -module, then the concept of prime submodules is equivalent to the concept of prime ideals. Now it is natural to ask if the “prime avoidance” theorem is true for prime submodules. In fact the answer is affirmative as has been shown by Lu [2].

THEOREM 1.2. *Let M be an R -module and P_1, \dots, P_n be prime submodules of M . Let N be a submodule of M such that $N \subseteq \bigcup_{i=1}^n P_i$. Also assume that $\text{Ann}(M/P_j) \not\subseteq \text{Ann}(M/P_k)$ for $j \neq k$. Then $N \subseteq P_j$ for some $1 \leq j \leq n$.*

The main purpose of this paper is to generalize the above theorem to a countable family of prime submodules.

2. Main result. Let R be a Noetherian ring, \mathfrak{a} an ideal of R contained in the Jacobson radical of R , and M a finitely-generated R -module. The \mathfrak{a} -adic topology is metrizable and one suitable metric d may be given by requiring that, for $x, y \in M$ with x and y different, $d(x, y) = 1/2^t$, where t is the greatest integer i such that $x - y \in \mathfrak{a}^i M$.

Now, we apply the Baire's category theorem for deducing our main result. Note that the Baire's category theorem states that a complete metric space is not the union of a countable family of nowhere dense subsets, and a subset of a metric space M is nowhere dense in M if and only if its closure has no interior point (see [1, page 38]). In the sequel, we denote the Jacobson radical of R by $J(R)$.

MAIN THEOREM 2.1. *Let M be a finitely-generated module over a Noetherian ring R . Suppose \mathfrak{a} is an ideal of R and let $N = \mathfrak{a}M$ and $\mathfrak{b} = \text{Ann}(M/N)$. If $\mathfrak{b} \subseteq J(R)$, M is complete with respect to the \mathfrak{b} -adic topology, $\{P_i\}_{i \geq 1}$ is a countable family of prime submodules of M , and $x \in M$, then $x + N \subseteq \bigcup_{i \geq 1} P_i$ implies that $x + N \subseteq P_j$ for some $j \geq 1$.*

We will need the following well-known result which is essentially due to Krull.

LEMMA 2.2. *If M is a finitely-generated module over a Noetherian ring R and \mathfrak{b} an ideal of R such that $\mathfrak{b} \subseteq J(R)$, then each submodule of M with respect to the \mathfrak{b} -adic topology is closed (see [3, Theorem 8.10]).*

PROOF OF THE MAIN THEOREM. By Lemma 2.2, each submodule of M and in particular N is closed with respect to the \mathfrak{b} -adic topology. One easily obtains that $x + N$ is also a closed subset of M . So the completeness of M yields that $x + N$ is a complete metric space. But by hypothesis we have $x + N = \bigcup_{i \geq 1} ((x + N) \cap P_i)$ and therefore it turns out by Baire's category theorem that, there is some $j \geq 1$ for which the subset $(x + N) \cap P_j$ of $x + N$ is not nowhere dense. But $(x + N) \cap P_j$ is closed in $x + N$ and therefore the interior of $(x + N) \cap P_j$ in $x + N$ is not empty. Let c be an element of the interior of $(x + N) \cap P_j$. Thus there exists an open subset U of M such that $c \in (x + N) \cap U \subseteq (x + N) \cap P_j$. But $c \in U$ implies that, there is $k \in \mathbb{N}$ such that $c + \mathfrak{b}^k M \subseteq U$ and hence $c \in (x + N) \cap (c + \mathfrak{b}^k M) \subseteq (x + N) \cap P_j$. It is easily deduced that $N \cap \mathfrak{b}^k M \subseteq P_j$. But $\mathfrak{b}^k M \subseteq N$, so $\mathfrak{b}^k M \subseteq P_j$ and $\mathfrak{b}^k \subseteq \text{Ann}(M/P_j)$. Since P_j is a prime submodule of M , it follows that $\text{Ann}(M/P_j)$ is a prime ideal of R and so $\mathfrak{b} \subseteq \text{Ann}(M/P_j)$. Now we have $\mathfrak{a} \subseteq \mathfrak{b} \subseteq \text{Ann}(M/P_j)$, and hence $\mathfrak{a}M = N \subseteq P_j$. But $x - c \in N$, so $x + N \subseteq P_j$, as required. \square

REMARK 2.3. (i) Note that if M is a finitely-generated module over a Noetherian ring R , $\mathfrak{a}_1 \subseteq \mathfrak{a}_2$ ideals of R , and M complete with respect to the \mathfrak{a}_2 -adic topology, then it is also complete with respect to the \mathfrak{a}_1 -adic topology. This shows that, the main theorem is also true if we assume that M is complete in $J(R)$ -adic topology.

(ii) In the view of (i), it is clear that Theorem 1.1 is a particular case of the main theorem.

ACKNOWLEDGEMENT. This research was in part supported by a grant from Institute for Studies in Theoretical Physics and Mathematics (IPM).

REFERENCES

- [1] E. T. Copson, *Metric Spaces*, Cambridge Tracts in Mathematics and Mathematical Physics, no. 57, Cambridge University Press, London, 1968. [MR 37#877](#). [Zbl 177.25303](#).
- [2] C.-P. Lu, *Unions of prime submodules*, Houston J. Math. 23 (1997), no. 2, 203–213. [MR 2000c:13015](#). [Zbl 885.13004](#).

- [3] H. Matsumura, *Commutative Ring Theory*, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1989. [MR 90i:13001](#). [Zbl 666.13002](#).
- [4] R. Y. Sharp and P. Vámos, *Baire's category theorem and prime avoidance in complete local rings*, Arch. Math. (Basel) **44** (1985), no. 3, 243–248. [MR 86h:13004](#). [Zbl 546.13002](#).

M. R. POURNAKI: DEPARTMENT OF MATHEMATICAL SCIENCES, SHARIF UNIVERSITY OF TECHNOLOGY, P.O. BOX 11365-9415, TEHRAN, IRAN AND SCHOOL OF MATHEMATICS, INSTITUTE FOR STUDIES IN THEORETICAL PHYSICS AND MATHEMATICS, P.O. BOX 19395-5746, TEHRAN, IRAN

E-mail address: pournaki@karun.ipm.ac.ir

M. TOUSI: DEPARTMENT OF MATHEMATICS, SHAHID BEHESHTI UNIVERSITY, EVIN, TEHRAN 19834, IRAN AND SCHOOL OF MATHEMATICS, INSTITUTE FOR STUDIES IN THEORETICAL PHYSICS AND MATHEMATICS, P.O. BOX 19395-5746, TEHRAN, IRAN

E-mail address: mtousi@vax.ipm.ac.ir

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru