

NONEMPTY INTERSECTION THEOREMS MINIMAX THEOREMS WITH APPLICATIONS IN GENERALIZED INTERVAL SPACES

DA-CHENG WANG

(Received 7 August 2000)

ABSTRACT. We establish some new topological types of nonempty intersection theorems in more general topological spaces without linear structure. As applications, we utilize results to study the minimax problems, coincidence problems, and economy equilibrium problems in generalized interval spaces and some new results are obtained.

2000 Mathematics Subject Classification. 49K35, 54D20, 54D05, 54H25.

1. Introduction and preliminaries. Recently, some versions of Fan's KKM theorem and Fan's minimax theorem in topological spaces without linear structure have been considered by Cheng and Lin [2], Lin et al. [10, 11], and Tian [14]. The purpose of this paper is first to introduce and establish a class of more general topological space-generalized interval space, which has no linear structure and contains H -space, interval space as its special cases, and so as it contains strong interval space [2], convex space, contractible space, and topological vector space as its special cases. With the aid of this kind of topological frameworks some new topological types of nonempty intersection theorems are established in Section 2. As applications, we utilize these results to obtain some minimax theorems and coincidence theorems in generalized interval space. The results presented in this paper not only contain many recent results of Stacho [13], Kindler and Trost [8], Cheng and Lin [2], Brezis et al. [1], Tuy [7], Komornik [9], Geraghty and Lin [4], Sion [12], Wu [15], Hartung [6], and the famous Neumann's saddle theorem as its special cases, but also improve and extend the corresponding results of [1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14].

For the sake of convenience, we first give some definitions and notations.

Let X and Y be two Hausdorff topological spaces. We denote by $C(X, Y)$ the set of all continuous mappings from X to Y . Let LB be a subset of X , we denote by $\text{cl}B$ the closure of B . If M is a nonempty subset of X , we denote by $\mathcal{A}(M)$, $\mathcal{H}(M)$, and 2^X the families of all nonempty finite subsets, all nonempty connected subsets, and all subsets of M , respectively. We denote $\mathbb{R} = (-\infty, +\infty)$.

DEFINITION 1.1 (see [2]). Let E be a linear ordered space. We call E a completely dense linear space, if each subset of E has a least upper bound and for any $Z_1, Z_2 \in E$ with $Z_1 < Z_2$, there exists $z \in E$ such that $Z_1 < z < Z_2$.

DEFINITION 1.2. Let X be a topological space, X is called a generalized interval space, if there exists a mapping $\Gamma : X \times X \rightarrow \mathcal{H}(X)$ such that for any $(\mathcal{X}_1, \mathcal{X}_2) \in X \times X$,

$\Gamma(\mathcal{X}_1, \mathcal{X}_2) = \Gamma(\mathcal{X}_2, \mathcal{X}_1)$, and $\Gamma(\mathcal{X}_1, \mathcal{X}_2)$ is called a generalized interval associated with $\mathcal{X}_1, \mathcal{X}_2$.

DEFINITION 1.3. Let X be a generalized interval space, Y a topological space, and E a completely dense linear space. A subset B of X is called T -convex set, if for any $\mathcal{X}_1, \mathcal{X}_2 \in B$, we have $\Gamma(\mathcal{X}_1, \mathcal{X}_2) \subset B$. A mapping $f : X \rightarrow E$ is called T -quasi-concave, if for any $\mathcal{X}_1, \mathcal{X}_2 \in X$, there exist $\mathfrak{y}_1, \mathfrak{y}_2 \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$ such that $f(\mathfrak{y}_i) \geq f(\mathcal{X}_i)$, $i = 1, 2$, and for any $z \in E$, the set $\{x \in X : f(x) \geq z\}$ is T -convex. We call $F : X \rightarrow 2^Y$ a T -KKM mapping, if for any $\mathcal{X}_1, \mathcal{X}_2 \in X$ we have $F(x) \subset F(\mathcal{X}_1) \cup F(\mathcal{X}_2)$ for all $x \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$, and there exist $\mathfrak{y}_1, \mathfrak{y}_2 \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$ such that $F(\mathfrak{y}_i) \subset F(\mathcal{X}_i)$, $i = 1, 2$.

REMARK 1.4. For Definitions 1.2 and 1.3 it is easy to see that both interval space and H -space are the special cases of generalized interval space, and so as all the strong interval space [2], convex space, contractible space, and topological vector space are the special cases of generalized interval space. For H -space, the H -convex set is T -convex. For topological linear space, a convex subset is T -convex, a quasi-concave function is T -quasi-concave.

REMARK 1.5. It is obvious that a nonempty T -convex subset in a generalized interval space is also a generalized interval space.

DEFINITION 1.6 (see [14]). Let X and Y be two topological spaces. A mapping $F : X \rightarrow 2^Y$ is called transfer closed valued, if for each $x \in X$, $y \notin F(x)$ implies that there exists a point $\mathcal{X}^* \in X$ such that $y \notin \text{cl}F(\mathcal{X}^*)$.

LEMMA 1.7. Let X be a generalized interval space, E a completely dense linear ordered space, and $f : X \rightarrow E$ a mapping. Then f is T -quasi-concave if and only if for any $\mathcal{X}_1, \mathcal{X}_2 \in X$, there exist $\mathfrak{y}_1, \mathfrak{y}_2 \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$ such that $f(\mathfrak{y}_i) \geq f(\mathcal{X}_i)$, $i = 1, 2$, and $f(x) \geq \min\{f(\mathcal{X}_1), f(\mathcal{X}_2)\}$ for all $x \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$.

PROOF. The proof follows from Definition 1.3 immediately. \square

LEMMA 1.8. Let X be a generalized interval space, Y a topological spaces, and E a completely dense linear ordered space. Let $\alpha \in E$ and $\varphi : X \times X \rightarrow E$ be T -quasi-concave in X . If we define a mapping $F : X \rightarrow 2^Y$ by $F(x) = \{y \in Y : \varphi(x, y) \leq \alpha\}$ for all $x \in X$, then F is a T -KKM mapping.

PROOF. Since φ is T -quasi-concave in X , for any given $\mathcal{X}_1, \mathcal{X}_2 \in X$ and for any $x \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$ we have $\varphi(x, y) \geq \min\{\varphi(\mathcal{X}_1, y), \varphi(\mathcal{X}_2, y)\}$ for any $y \in Y$, and there exist $\mathcal{X}_1^*, \mathcal{X}_2^* \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$ such that $\varphi(\mathcal{X}_i^*, y) \geq \varphi(\mathcal{X}_i, y)$, $i = 1, 2$ for all $y \in Y$. Hence $F(\mathcal{X}_i^*) \subset F(\mathcal{X}_i)$, $i = 1, 2$, for all $x \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$, we have $F(x) \subset \bigcup_{i=1}^2 F(\mathcal{X}_i)$. This implies that F is a T -KKM mapping. Let X, Y be two topological spaces, $F : X \rightarrow 2^Y$. In the sequel we denote $\text{cl}\{F(x)\} = \text{cl}F(x)$. \square

2. Some nonempty intersection theorems

THEOREM 2.1. Let X be a generalized interval space, Y a topological space, and $F : X \rightarrow 2^Y$ an upper semi-continuous T -KKM mapping. Suppose that the following conditions are satisfied:

- (1) for any $x \in X$, $F(x)$ is a nonempty closed (open) set in Y ;

(2) for any $A \in \mathcal{A}(X)$, $\bigcap_{x \in A} F(x) \in \mathcal{K}(Y)$,
then

- (1) $\{F(x) : x \in X\}$ has the finite intersection property;
(2) if there exists an $A_0 \in \mathcal{A}(X)$ such that $\bigcap_{x \in A_0} \text{cl}F(x)$ is compact, then

$$\bigcap_{x \in A} \text{cl}F(x) \neq \emptyset. \quad (2.1)$$

PROOF. Letting $\Omega = \{F(x) : x \in X\}$, and by using induction we prove that Ω has the finite intersection property.

By condition (1), for any $F(x)$ in Ω , $x \in X$ is nonempty. Suppose that for any n elements in Ω , $n \geq 2$, their intersection is nonempty, next we prove that for any $n+1$ elements in Ω , their intersection is also nonempty. Suppose the contrary, then there exist $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_{n+1} \in X$ such that $\bigcap_{i=1}^{n+1} F(\mathcal{X}_i) = \emptyset$. Letting $H = \bigcap_{i=1}^{n+1} F(\mathcal{X}_i)$, then

$$\bigcap_{i=1}^2 H \cap F(\mathcal{X}_i) = \emptyset. \quad (2.2)$$

By condition (2) and the assumption of induction, for any $x \in X$

$$H \cap F(x) \text{ is nonempty and connected.} \quad (2.3)$$

Since F is a T -KKM mapping, $F(x) \subset \bigcup_{i=1}^2 F(\mathcal{X}_i)$ for all $x \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$ and there exist $\mathcal{X}_1^*, \mathcal{X}_2^* \in \Gamma(\mathcal{X}_1, \mathcal{X}_2)$ such that $F(\mathcal{X}_i^*) \subset F(\mathcal{X}_i)$, $i = 1, 2$. Hence,

$$H \cap F(x) \subset \bigcup_{i=1}^2 H \cap F(\mathcal{X}_i), \quad \forall x \in \Gamma(\mathcal{X}_1, \mathcal{X}_2), \quad (2.4)$$

letting $D_j = \{x \in \Gamma(\mathcal{X}_1, \mathcal{X}_2) : H \cap F(x) \subset H \cap F(\mathcal{X}_j)\}$, $j = 1, 2$, then $\mathcal{X}_i^* \in D_i$, $i = 1, 2$. It follows from (2.2), (2.3), (2.4), and condition (1) that $D_1 \cup D_2 = \Gamma(\mathcal{X}_1, \mathcal{X}_2)$. Since $\Gamma(\mathcal{X}_1, \mathcal{X}_2)$ is connected, we know that either $\text{cl}D_1 \cap D_2$ or $\text{cl}D_2 \cap D_1$ is nonempty. Without loss of generality, we can assume $\text{cl}D_2 \cap D_1 \neq \emptyset$. Hence there exists an $x_0 \in \text{cl}D_2 \cap D_1$. By (2.2), we have $(H \cap F(\mathcal{X}_0)) \cap (H \cap F(\mathcal{X}_2)) = \emptyset$, and it is easy to prove that there exists an open set U containing $F(\mathcal{X}_0)$ such that $(H \cap U) \cap (H \cap F(\mathcal{X}_2)) = \emptyset$. In fact, if F is open valued, it is sufficient to take $U = F(\mathcal{X}_0)$; if F is closed valued, taking $U = Y \setminus (H \cap F(\mathcal{X}_2))$, then U is an open set in Y and

$$(H \cap U) \cap (H \cap F(\mathcal{X}_2)) = \emptyset, \quad U \supset (H \cap F(\mathcal{X}_0)) \neq \emptyset. \quad (2.5)$$

Moreover, for any $y \notin U$, then $y \in H \cap F(\mathcal{X}_2)$ and $y \notin H \cap F(\mathcal{X}_0)$, and so $y \notin F(\mathcal{X}_0)$. Hence $F(\mathcal{X}_0) \subset U$. This means that U satisfies the desired condition.

By the upper semi-continuity of F , there exists an open neighborhood V of \mathcal{X}_0 such that for any $x \in V$, $F(x) \subset U$. Since $\mathcal{X}_0 \in \text{cl}D_2$, $V \cap D_2 \neq \emptyset$. Taking $\mathcal{X}^* \in V \cap D_2$, we have $H \cap F(\mathcal{X}^*) \subset H \cap F(\mathcal{X}_2)$, $H \cap F(\mathcal{X}^*) \subset H \cap U$. Hence we have $(H \cap F(\mathcal{X}_2)) \cap (H \cap U) \supset H \cap F(\mathcal{X}^*) \neq \emptyset$. This contradicts (2.5), therefore Ω has the finite intersection property.

In addition, if there exists an $A_0 \in \mathcal{A}(M)$ such that $\bigcap_{x \in A_0} \text{cl}F(x)$ is compact. Letting $K = \bigcap_{x \in A_0} \text{cl}F(x)$, it follows from the finite intersection property of Ω and A_0 being finite that $\{\text{cl}F(x) \cap K : x \in X\}$ is a family of closed set in the compact set K and has the finite intersection property, therefore $\bigcap_{x \in X} (\text{cl}F(x) \cap K) = \bigcap_{x \in X} \text{cl}F(x) \neq \emptyset$. This completes the proof. \square

THEOREM 2.2. *Let X be a generalized interval space, Y a topological space, $F : X \rightarrow 2^Y$ a mapping with nonempty values and satisfy the following conditions:*

- (1) F is transfer closed valued;
- (2) $\text{cl}F : X \rightarrow 2^Y$ is an upper semi-continuous T -KKM mapping;
- (3) for each $A \in \mathcal{A}(X)$, $\bigcap_{x \in X} \text{cl}F(x) \in \mathcal{K}(Y)$;
- (4) there exists an $A_0 \in \mathcal{A}(X)$ such that $\bigcap_{x \in A_0} \text{cl}F(x)$ is compact, then $\bigcap_{x \in X} F(x) \neq \emptyset$.

PROOF. First we prove $\bigcap_{x \in X} F(x) = \bigcap_{x \in X} \text{cl}F(x)$. In fact, since $\bigcap_{x \in X} F(x) \subset \bigcap_{x \in X} \text{cl}F(x)$, it is sufficient to prove that $\bigcap_{x \in X} F(x) \supset \bigcap_{x \in X} \text{cl}F(x)$. Suppose the contrary, then there exists $Y_0 \in \bigcap_{x \in X} \text{cl}F(x)$ such that $y_0 \notin \bigcap_{x \in X} F(x)$. Hence there exists an $x_0 \in X$ such that $y_0 \notin F(x_0)$. By condition (1), there exists $x' \in X$ such that $y_0 \notin \text{cl}F(x)$. This contradicts the choice y_0 . Hence the desired conclusion is proved. \square

It is obvious that $\text{cl}F : X \rightarrow 2^Y$ satisfies all the conditions in [Theorem 2.1](#). By [Theorem 2.1](#), we know that $\bigcap_{x \in X} F(x) = \bigcap_{x \in X} \text{cl}F(x) \neq \emptyset$.

THEOREM 2.3. *Let X be a generalized interval space, Y a topological space, $S \in C(X, Y)$ a given mapping, and $F : X \rightarrow 2^Y$ a mapping satisfying the following conditions:*

- (1) F is transfer closed valued;
- (2) $\text{cl}F : X \rightarrow 2^Y$ is an upper semi-continuous T -KKM mapping;
- (3) there exists a compact set L in Y and a subset K in X such that for any $A \in \mathcal{A}(X)$ there exists a compact T -convex subset $D \subset X$ with $K \cup A \subset D$ such that
 - (a) for any $x \in D$, $\text{cl}F(x) \cap s(D) \neq \emptyset$;
 - (b) for any finite set $\{x_1, x_2, \dots, x_n\} \subset D$, $\bigcap_{i=1}^n (\text{cl}F(x_i) \cap s(D)) \in \mathcal{K}(Y)$;
 - (c) $\bigcap_{x \in D} (\text{cl}F(x) \cap s(D)) \subset L$,

then $\bigcap_{x \in X} F(x) \neq \emptyset$.

PROOF. By condition (1), $\bigcap_{x \in X} F(x) = \bigcap_{x \in X} \text{cl}F(x)$. Hence in order to prove the conclusion of the theorem, it suffices to prove that $\bigcap_{x \in X} \text{cl}F(x) \neq \emptyset$. Since L is compact, for the purpose, it suffices to prove that $\{\text{cl}F(x) \cap L : x \in X\}$ has the finite intersection property. For any $A \in \mathcal{A}(X)$, let D be the set satisfying condition (3). Hence we have

$$\bigcap_{x \in A} (\text{cl}F(x) \cap L) \supset \bigcap_{x \in D} (\text{cl}F(x) \cap s(D)). \quad (2.6)$$

Letting $Y_0 = s(D)$, by the continuity of s and compactness of D , we know that Y_0 is a compact subset of Y .

Next we prove that $\bigcap_{x \in D} (\text{cl}F(x) \cap Y_0) \neq \emptyset$. In fact, letting $G(x) = \text{cl}F(x) \cap Y_0$, $x \in D$, by condition (3)(a), for any $x \in X$, $G(x)$ is nonempty compact. Since D is T -convex, D is a generalized interval space. It is easy to prove that $G : D \rightarrow 2^Y$ satisfies all conditions in [Theorem 2.1](#). By [Theorem 2.1](#), $\{G(x) : x \in X\}$ has the finite intersection property. Therefore, we have $\bigcap_{x \in D} G(x) \neq \emptyset$, that is, $\bigcap_{x \in D} (\text{cl}F(x) \cap Y_0) \neq \emptyset$. By (2.6), $\bigcap_{x \in A} (\text{cl}F(x) \cap L) \neq \emptyset$. This implies that $\{\text{cl}F(x) \cap L : x \in X\}$ has the finite intersection property. \square

THEOREM 2.4. *Let X be a generalized interval space, Y a compact topological space, and $G : X \rightarrow 2^Y$ a mapping with nonempty values and satisfy the following conditions:*

- (1) G is transfer closed valued;
- (2) $\text{cl}G : X \rightarrow 2^Y$ is a T -KKM mapping;
- (3) for any $A \in \mathcal{A}(X)$, $\bigcap_{x \in A} \text{cl}G(x) \in \mathcal{K}(Y)$;
- (4) $\text{cl}G$ is closed graph,

then $\bigcap_{x \in X} G(x) \neq \emptyset$.

PROOF. Since $\text{cl}G$ is a closed graph and Y is compact, by using Fan [3, Lemma 2] we know that $\text{cl}G$ is upper semi-continuous. The conclusion follows from [Theorem 2.2](#) immediately. \square

3. Applications to minimax problems. As applications, we use the results obtained in [Section 2](#) to give some minimax theorems in generalized interval spaces.

THEOREM 3.1. *Let X be a generalized interval space, Y a compact topological space, E a completely dense linear ordered space, and $\varphi : X \times X \rightarrow E$ a lower semi-continuous function satisfying the following conditions:*

- (1) $\varphi(x, y)$ is T -quasi-concave in x ;
- (2) for any $A \in \mathcal{A}(X)$ and for any $\alpha \in E$, $\bigcap_{x \in A} \{y \in Y : \varphi(x, y) \leq \alpha\} \in \mathcal{K}(Y)$,

then $\alpha_* = \sup_{x \in X} \inf_{y \in Y} \varphi(x, y) = \inf_{y \in Y} \sup_{x \in X} \varphi(x, y) = \alpha^*$.

PROOF. It is obvious that $\alpha_* \leq \alpha^*$.

Now we prove that $\alpha_* \geq \alpha^*$. Without loss of generality, we can assume that α_* is not the greatest element in E . For any $x \in X$ and for any given $\alpha > \alpha_*$, let

$$G(x, \alpha) = \{y \in Y : \varphi(x, y) \leq \alpha\}. \quad (3.1)$$

Next we prove that $\{G(x, \alpha) : x \in X, \alpha > \alpha_*\}$ has the finite intersection property. In fact, for any n points $x_1, \dots, x_n \in X$ and any n elements $\alpha_1, \dots, \alpha_n \in E$ with $\alpha_i > \alpha_*$, $i = 1, \dots, n$, since E is a linear ordered space, we can assume that $\alpha_n \geq \dots \geq \alpha_1 > \alpha_*$. By the density of E , there exists an $\bar{\alpha} \in E$ such that $\alpha_1 > \bar{\alpha} > \alpha_*$. Letting

$$F(x) = G(x, \bar{\alpha}), \quad x \in X, \quad (3.2)$$

by the lower semi-continuity of φ and the choice of $\bar{\alpha} \in E$, we know that $F : X \rightarrow 2^Y$ is nonempty closed valued. This implies that F satisfies condition (1) in [Theorem 2.1](#). From condition (2), F satisfies condition (2) in [Theorem 2.1](#). In view of condition (1) and [Lemma 1.7](#), we know that F is a T -KKM mapping. In addition, since $\varphi : X \times Y \rightarrow E$ is lower semi-continuous, F is closed graph. By Fan [3, Lemma 2], F is upper semi-continuous. Therefore F satisfies all the conditions in [Theorem 2.1](#). By [Theorem 2.1](#), $\{F(x) : x \in X\}$ has the finite intersection property, and so $\bigcap_{i=1}^n F(x_i) \neq \emptyset$. Since $\bigcap_{i=1}^n F(x_i) \subset \bigcap_{i=1}^n G(x_i, \alpha_i)$, $\{G(x, \alpha) : x \in X, \alpha > \alpha_*\}$ has the finite intersection property.

On the other hand, for any $x \in X$ and for any $\alpha > \alpha_*$, $G(x, \alpha)$ is closed. Since Y is compact, $\bigcap_{x \in X, \alpha > \alpha_*} G(x, \alpha) \neq \emptyset$. Hence, there exists $\bar{y} \in Y$ such that $\bar{y} \in G(x, \alpha)$ for all $x \in X$ and for all $\alpha > \alpha_*$, that is, $\varphi(x, \bar{y}) \leq \alpha$ for all $x \in X$ and for all $\alpha > \alpha_*$. Hence, we have $\inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq \alpha$, for all $\alpha > \alpha_*$.

Since E is a dense linear ordered space, we know that $\alpha^* = \inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq \alpha_*$.

Summing up the above arguments, we have $\alpha_* = \alpha^*$. This completes the proof. \square

REMARK 3.2. Theorem 3.1 not only contains Theorem 1 in Shacho [13] as its special case but also relaxes its continuous condition.

THEOREM 3.3. Let X be a generalized interval space, Y a topological space, E a completely dense linear ordered space, and $\varphi, \phi : X \times X \rightarrow E$ two functions satisfying the following conditions:

- (1) $\varphi(x, y) \leq \phi(x, y)$ for all $(x, y) \in X \times Y$;
- (2) $\varphi(x, y)$ is T -quasi-concave in x , and it is upper semi-continuous on any generalized interval of X ;
- (3) $\varphi(x, y)$ is lower semi-continuous in y and for any $A \in \mathcal{A}(X)$, the set $\bigcap_{x \in A} \{y \in Y : \varphi(x, y) < \alpha\}$ is connected for all $\alpha \in E$;
- (4) there exists an $A_0 \in \mathcal{A}(X)$ and a $z_0 > \sup_{x \in X} \inf_{y \in Y} \phi(x, y)$ such that $\bigcap_{x \in A_0} \{y \in Y : \varphi(x, y) \leq z_0\}$ is compact,

then $\inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq \sup_{x \in X} \inf_{y \in Y} \phi(x, y)$.

PROOF. Letting $z_* = \sup_{x \in X} \inf_{y \in Y} \phi(x, y)$, by the completeness of E , we know that z_* exists. From condition (4), z_* is not a largest element in E .

For any $x \in X$ and for any $z > z_*$, let $F(x, z) = \{y \in Y : \varphi(x, y) \leq z\}$. It follows from conditions (1) and (3) that $F(x, z)$ is a nonempty closed set. By condition (4), $\bigcap_{x \in A} F(x, z)$ is compact. Denote $\Omega = \{F(x, z) : x \in X, z \in Z\}$. Next we prove that Ω has the finite intersection property. In fact, we have proved that each element in Ω is nonempty. Suppose for any n elements in Ω their intersection is nonempty. Now we prove that any $n + 1$ elements in Ω their intersection is nonempty. Suppose the contrary, then there exist $\mathcal{X}_1, \dots, \mathcal{X}_n, \mathcal{X}_{n+1}$ in X and z_1, \dots, z_n, z_{n+1} in E with $z_i > z_*$, $i = 1, 2, \dots, n + 1$ such that $\bigcap_{i=1}^{n+1} F(x_i, z_i) = \emptyset$. Without loss of generality, we can assume that $z_* < z_1 \leq \dots \leq z_n \leq z_{n+1}$. By the density of E , there exists a $\bar{z} \in E$ such that $z_1 > \bar{z} > z_*$. Letting

$$T(x, z) = \{y \in Y : \varphi(x, y) < z\}, \quad \forall x \in X, \forall z > z_*;$$

$$T(x) = T(x, z_1), \quad \forall x \in X; H = \bigcap_{i=3}^{n+1} T(x_i). \tag{3.3}$$

By the assumption of the induction, for any $x \in X$,

$$H \cap T(x) \supset \left(\bigcap_{i=3}^{n+1} F(x_i; \bar{z}) \cap F(x; \bar{z}) \right) \neq \emptyset. \tag{3.4}$$

It follows from condition (2) and (3), for $x_1, x_2 \in X$ there exist $x'_1, x'_2 \in \Gamma(x_1, x_2)$ such that $T(x'_i) \subset T(x_i)$, $i = 1, 2$ and $T(x) \subset \bigcup_{i=1}^2 T(x_i)$ for all $X \in \Gamma(x_1, x_2)$. Hence we have

$$H \cap T(x) \subset \bigcup_{i=1}^2 (H \cap T(x_i)), \quad \forall x \in \Gamma(x_1, x_2). \tag{3.5}$$

By condition (3),

$$\bigcap_{i=1}^2 \text{cl}(H \cap T(x_i)) \subset \bigcap_{i=1}^{n+1} F(x_i, z_i) \neq \emptyset. \tag{3.6}$$

This shows that $H \cap T(x_1)$ and $H \cap T(x_2)$ are separated. Letting $I_i = \{x \in \Gamma(x_1, x_2) :$

$H \cap T(x) \subset H \cap T(x_i)$, $i = 1, 2$, from (3.5) and $T(x'_i) \subset T(x_i)$, $i = 1, 2$, we know that $x'_i \in I_i$, $i = 1, 2$ and $I_1 \cup I_2 = \Gamma(x_1, x_2)$. Since $\Gamma(x_1, x_2)$ is connected, either $I_1 \cap \text{cl}I_2$ or $\text{cl}I_1 \cap I_2$ is nonempty. Without loss of generality, we can assume that $I_1 \cup \text{cl}I_2 \neq \emptyset$. Taking $x_0 \in I_1 \cap \text{cl}I_2$, then there exists a net $\{x_y\}_{y \in \theta} \subset I_2$ such that $x_y \rightarrow x_0$ and $H \cap T(x_1) \supset H \cap T(x_0) \neq \emptyset$. Taking $y_0 \in H \cap T(x_0)$, we have

$$\varphi(x_0, y_0) < z_1. \tag{3.7}$$

It follows from (3.6) that $y_0 \notin H \cap T(x_2)$. Hence, $y_0 \notin H \cap T(x_y)$ for all $y \in \theta$, that is, $\varphi(x_y, y_0) \geq z_1$ for all $y \in \theta$.

By condition (2) and $x_y \rightarrow x_0$, we have $\varphi(x_0, y_0) \geq z_1$. This contradicts (3.7). Therefore Ω has finite intersection property.

By condition (4), $K = \bigcap_{x \in A_0} F(x, z_0)$ is compact and $\{F(x, z) \cap K : x \in X, z \in z_*\}$ has the finite intersection property, therefore we have $\bigcap_{x \in X, z \in z_*} F(x, z) = \bigcap_{x \in X, z \in z_*} (F(x, z) \cap K) \neq \emptyset$, and so $\inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq z_*$. This completes the proof. \square

REMARK 3.4. Theorem 3.3 improves and extends Theorem 1 in Lin and Quan [10].

COROLLARY 3.5. Let X be a generalized interval space, Y a topological space, E a completely dense linear ordered space, and $f : X \times X \rightarrow E$ a function satisfying the following conditions:

- (1) $f(x, y)$ is T -quasi-concave in x and it is upper semi-continuous on any generalized interval of X ;
- (2) $f(x, y)$ is lower semi-continuous in y and for any $A \in \mathcal{A}(X)$, the set

$$\bigcap_{x \in A} \{y \in Y : f(x, y) < \alpha\} \in \mathcal{K}(Y), \quad \forall \alpha \in E; \tag{3.8}$$

- (3) there exist an $A_0 \in \mathcal{A}(X)$ and a $z_0 > \sup_{x \in X} \inf_{y \in Y} f(x, y)$ such that

$$\bigcap_{x \in A_0} \{y \in Y : f(x, y) \leq z_0\} \in \mathcal{K}(Y), \tag{3.9}$$

then $\inf_{y \in Y} \sup_{x \in X} f(x, y) = \sup_{x \in X} \inf_{y \in Y} f(x, y)$.

PROOF. It is obvious that $\inf_{y \in Y} \sup_{x \in X} f(x, y) \geq \sup_{x \in X} \inf_{y \in Y} f(x, y)$. On the other hand, by Theorem 3.3 we know that $\inf_{y \in Y} \sup_{x \in X} f(x, y) \leq \sup_{x \in X} \inf_{y \in Y} f(x, y)$. This completes the proof. \square

REMARK 3.6. The corresponding results in Kindler and Trost [8], Tuy [7], Cheng and Lin [2], Brezis et al. [1], Komornik [9], Geraghty and Lin [4], Stacho [13], Sion [12], Wu [15], and Hartung [6] all are the special cases of Corollary 3.5.

COROLLARY 3.7. Let X be a compact generalized interval space, E a completely dense linear ordered space, and $\varphi, \phi : X \times X \rightarrow E$, $\theta \in E$. If the following conditions are satisfied:

- (1) for any $(x, y) \in X \times X$, $\varphi(x, y) \leq \phi(x, y)$;
- (2) $\varphi(x, y)$ is T -quasi-concave in x and it is upper semi-continuous on any generalized interval of X ; $\phi(x, y)$ is upper semi-continuous in x ;

- (3) $\varphi(x, y)$ is lower semi-continuous in y , and for any $A \in \mathcal{A}(X)$ and any $\alpha \in E$, the set

$$\bigcap_{x \in A} \{y \in X : f(x, y) < \alpha\} \in \mathcal{K}(X), \quad (3.10)$$

then

- (1) there exists $\bar{x} \in X$ such that $\phi(\bar{x}, \bar{x}) > \theta$, or
- (2) there exists $\bar{y} \in X$ such that $\varphi(x, \bar{y}) \leq \theta$ for all $x \in X$.

PROOF. Suppose that (1) is not true, then for any $x \in X$ we have $\phi(x, x) \leq \theta$. By condition (2), $\inf_{y \in Y} \phi(x, y)$ is upper semi-continuous in x . Moreover, by [Theorem 3.3](#) and the compactness of X there exists an $\bar{x} \in X$ such that

$$\inf_{y \in X} \sup_{x \in X} \varphi(x, y) \leq \sup_{x \in X} \inf_{y \in X} \phi(x, y) = \inf_{y \in X} \phi(\bar{x}, y) \leq \phi(\bar{x}, \bar{x}) \leq \theta. \quad (3.11)$$

Since X is compact, by condition (3) there exists a $\bar{y} \in X$ such that

$$\varphi(x, \bar{y}) \leq \sup_{x \in X} \varphi(x, \bar{y}) = \inf_{y \in X} \sup_{x \in X} \varphi(x, y), \quad \forall x \in X. \quad (3.12)$$

□

COROLLARY 3.8. Let X be a compact generalized interval space, Y a compact topological space, E a completely dense linear ordered space, and $f : X \times Y \rightarrow E$ a mapping satisfying the following conditions:

- (1) $x \mapsto f(x, y)$ is T -quasi-concave and upper semi-continuous;
 - (2) $y \mapsto f(x, y)$ is lower semi-continuous;
 - (3) for any $\alpha \in E$, any $A \in \mathcal{A}(X)$, $\bigcap_{x \in A} \{y \in Y : f(x, y) < \alpha\} \in \mathcal{K}(Y)$,
- then there exists a saddle point $(\bar{x}, \bar{y}) \in X \times Y$ such that $f(x, \bar{y}) \leq f(\bar{x}, \bar{y}) \leq f(\bar{x}, y)$ for all $x \in X$ and for all $y \in Y$.

PROOF. If $\sup_{x \in X} \inf_{y \in Y} f(x, y)$ is a greatest element in E , then it is obvious that

$$\inf_{y \in Y} \sup_{x \in X} f(x, y) = \sup_{x \in X} \inf_{y \in Y} f(x, y). \quad (3.13)$$

If $\sup_{x \in X} \inf_{y \in Y} f(x, y)$ is not a greatest element in E , then f satisfies all the conditions in [Corollary 3.5](#) obviously. By [Corollary 3.5](#), we also have

$$\inf_{y \in Y} \sup_{x \in X} f(x, y) = \sup_{x \in X} \inf_{y \in Y} f(x, y). \quad (3.14)$$

In view of the compactness of X and Y , there exist $\bar{x} \in X$ and $\bar{y} \in Y$ such that

$$\begin{aligned} \sup_{x \in X} \inf_{y \in Y} f(x, y) &= \inf_{y \in Y} f(\bar{x}, y) \leq f(\bar{x}, \bar{y}), \quad \forall y \in Y, \\ \inf_{y \in Y} \sup_{x \in X} f(x, y) &= \sup_{x \in X} f(x, \bar{y}) \geq f(x, \bar{y}), \quad \forall x \in X. \end{aligned} \quad (3.15)$$

Therefore, we have $f(\bar{x}, y) \geq f(x, \bar{y})$ for all $y \in Y$ and for all $x \in X$.

Taking $x = \bar{x}$ in the preceding inequality, we have $f(\bar{x}, y) \geq f(\bar{x}, \bar{y})$ for all $y \in Y$. Taking $y = \bar{y}$ in the preceding inequality, we have $f(\bar{x}, \bar{y}) \geq f(x, \bar{y})$ for all $x \in X$. Therefore, we have

$$f(x, \bar{y}) \leq f(\bar{x}, \bar{y}) \leq f(\bar{x}, y), \quad \forall y \in Y, \forall x \in X. \quad (3.16)$$

□

COROLLARY 3.9 (von Neumann). *Let C, F be two Hausdorff topological vector spaces, $X \subset C, Y \subset F$ two compact convex subsets, and $\phi : X \times Y \rightarrow \mathbb{R}$ a function satisfying the following conditions:*

- (1) *for any $x \in X, \phi(x, y)$ is an upper semi-continuous concave function in y ;*
 - (2) *for any $y \in Y, \phi(x, y)$ is a lower semi-continuous convex function in x ,*
- then there exists a saddle point $(\bar{x}, \bar{y}) \in X \times Y$ of ϕ .*

PROOF. For any $x_1, x_2 \in X$ and for all $y_1, y_2 \in Y$, letting

$$\Gamma(x_1, x_2) = \text{co}\{x_1, x_2\}, \quad \Gamma(y_1, y_2) = \text{co}\{y_1, y_2\}, \quad (3.17)$$

then ϕ satisfies all the conditions in [Corollary 3.8](#). Therefore, the conclusion of [Corollary 3.9](#) follows from [Corollary 3.8](#) immediately. \square

COROLLARY 3.10. *Let X be a compact generalized interval space, $f : X \times X \rightarrow \mathbb{R}$, and $h : X \rightarrow \mathbb{R}$ two functions satisfying the following conditions:*

- (1) *for any $x \in X, f(x, x) \leq 0$;*
- (2) *$x \mapsto f(x, y)$ is upper semi-continuous and $y \mapsto f(x, y)$ is lower semi-continuous; h is upper semi-continuous;*
- (3) *$f(x, y) + h(x)$ is T -quasi-concave in x ;*
- (4) *for any $A \in \mathcal{A}(X)$ and for any $\alpha \in \mathbb{R}$ the set $\bigcap_{x \in A} \{y \in Y : f(x, y) - h(x) < \alpha\} \in \mathcal{H}(X)$,*

then there exists a $\bar{y} \in X$ such that $f(x, \bar{y}) \leq h(\bar{y}) - h(x)$ for all $x \in X$.

PROOF. Letting $g(x, y) = f(x, y) - h(y) + h(x)$, it is easy to prove that g satisfies all the conditions in [Corollary 3.8](#). Hence, there exist $x^*, y^* \in X$ such that $g(x, y^*) \leq g(x^*, y^*)$ for all $x, y \in X$. Hence $g(x, y^*) \leq g(x^*, y^*) \leq 0$ for all $x \in X$, that is, $f(x, y^*) \leq h(y^*) - h(x)$ for all $x \in X$. \square

THEOREM 3.11. *Let X be a generalized interval space, Y a topological space, E a completely dense linear ordered space, and $\phi, \varphi : X \times X \rightarrow E$ two mappings satisfying the following conditions:*

- (1) *for any $(x, y) \in X \times Y, \varphi(x, y) \leq \phi(x, y)$;*
- (2) *$x \mapsto \varphi(x, y)$ is T -quasi-concave and upper semi-continuous, $y \mapsto \varphi(x, y)$ is lower semi-continuous;*
- (3) *$\phi(x, y)$ is upper semi-continuous in x ;*
- (4) *there exist a compact K of X and a nonempty subset Y_0 of Y satisfying the following condition: for any $B \in \mathcal{A}(Y)$ there exists a compact set M containing $Y_0 \cup B$ such that for any $A \in \mathcal{A}(X)$ and for any $\alpha \in E$ the set $\bigcap_{x \in A} \{y \in M : \varphi(x, y) < \alpha\} \in \mathcal{H}(Y)$, and*

$$\sup_{x \notin K} \inf_{y \in Y_0} \phi(x, y) \leq \sup_{x \in X} \inf_{y \in Y} \phi(x, y), \quad (3.18)$$

then $\inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq \sup_{x \in X} \inf_{y \in Y} \phi(x, y)$.

PROOF. Letting $z^* = \inf_{y \in Y} \sup_{x \in X} \varphi(x, y)$, $z_* = \sup_{x \in X} \inf_{y \in Y} \phi(x, y)$. From the completeness of E , we know that z_* and z^* both exist. If $z^* > z_*$, then it follows from the density of E that there exists a $\bar{z} \in E$ such that $z^* > \bar{z} > z_*$. Let

$$F(y) = \{x \in X : \phi(x, y) \geq \bar{z}\}, \quad y \in Y. \quad (3.19)$$

By condition (3), $F(\gamma)$ is closed for all $\gamma \in Y$. From condition (4), for any $\gamma_1, \dots, \gamma_n \in Y$ there exists a compact set M containing $Y_0 \cup \{\gamma_1, \dots, \gamma_n\}$. It is easy to see that $\phi, \varphi : X \times M \rightarrow E$ satisfy all the conditions in [Theorem 3.3](#). By [Theorem 3.3](#), we know

$$\inf_{\gamma \in M} \sup_{x \in X} \varphi(x, \gamma) \leq \sup_{x \in X} \inf_{\gamma \in M} \phi(x, \gamma). \quad (3.20)$$

On the other hand, since $\inf_{\gamma \in M} \sup_{x \in X} \varphi(x, \gamma) \geq \inf_{\gamma \in Y} \sup_{x \in X} \varphi(x, \gamma) = z^* > \bar{z}$, $\sup_{x \in X} \inf_{\gamma \in M} \phi(x, \gamma) > \bar{z}$, and so $\bigcap_{\gamma \in M} F(\gamma) \neq \emptyset$. Hence we have $\bigcap_{\gamma \in Y_0} F(\gamma) \supset \bigcap_{\gamma \in M} F(\gamma) \neq \emptyset$. By condition (4), we have $\sup_{x \notin K} \inf_{\gamma \in Y_0} \phi(x, \gamma) \leq z_* < \bar{z}$. Hence, for all $x \in X \setminus K$ we have $\inf_{\gamma \in Y_0} \phi(x, \gamma) \leq z_* < \bar{z}$, and so there exists a $\gamma_0 \in Y_0$ such that $\phi(x, \gamma_0) < \bar{z}$, that is, $x \notin F(\gamma_0)$. Hence $x \notin \bigcap_{\gamma \in Y_0} F(\gamma) = D$. This implies that $D \subset K$. Since K is compact and D is closed, then D is a compact set.

Summing up the above arguments, we have proved that $\{F(\gamma) \cap D : \gamma \in Y\}$ is a family of compact sets in D having the finite intersection property. Hence, we have

$$\bigcap_{\gamma \in Y} F(\gamma) = \bigcap_{\gamma \in Y} (F(\gamma) \cap D) \neq \emptyset. \quad (3.21)$$

Therefore $z_* = \sup_{x \in X} \inf_{\gamma \in Y} \phi(x, \gamma) \geq \bar{z}$. This contradicts the choice of \bar{z} . Therefore $z^* \leq z_*$. This completes the proof. \square

REMARK 3.12. [Theorem 3.11](#) improves the corresponding results in Ha [5] and Lin and Quan [10]. From [Theorem 3.11](#) we can obtain the following results.

COROLLARY 3.13. *Let X be a generalized interval space, Y a topological space, E a completely dense linear ordered space, and $\varphi : X \times Y \rightarrow E$ a mapping satisfying the following conditions:*

- (1) $\varphi(x, \gamma)$ is T -quasi-concave and upper semi-continuous in x and it is lower semi-continuous in γ ;
- (2) there exist a compact subset K in X and a nonempty subset Y_0 in Y satisfying the following condition: for any $B \in \mathcal{D}(Y)$ there exists a compact set M containing $Y_0 \cup B$ such that for any $A \in \mathcal{D}(X)$ and for any $\alpha \in E$, $\bigcap_{x \in A} \{\gamma \in M : \varphi(x, \gamma) < \alpha\} \in \mathcal{H}(Y)$ and

$$\sup_{x \notin K} \inf_{\gamma \in Y_0} \varphi(x, \gamma) \leq \sup_{x \in X} \inf_{\gamma \in Y} \varphi(x, \gamma), \quad (3.22)$$

then $\inf_{\gamma \in Y} \sup_{x \in X} \varphi(x, \gamma) = \sup_{x \in X} \inf_{\gamma \in Y} \varphi(x, \gamma)$.

THEOREM 3.14. *Let X be a generalized interval space, Y a topological space, E a completely dense linear ordered space, $\varphi : X \times Y \rightarrow E$ lower semi-continuous, and $\phi : X \times Y \rightarrow E$ upper semi-continuous in x . Suppose that the following conditions are satisfied:*

- (1) $\varphi(x, \gamma)$ is T -quasi-concave in x and $\varphi(x, \gamma) \leq \phi(x, \gamma)$ for all $(x, \gamma) \in X \times Y$;
- (2) there exist a compact subset K in X and a nonempty subset Y_0 in Y satisfying the following condition: for any $B \in \mathcal{D}(Y)$ there exists a compact set M containing $Y_0 \cup B$ such that for any $A \in \mathcal{D}(X)$ and for all $\alpha \in E$, $\bigcap_{x \in A} \{\gamma \in M : \varphi(x, \gamma) \leq \alpha\} \in \mathcal{H}(Y)$ and

$$\sup_{x \notin K} \inf_{\gamma \in Y_0} \phi(x, \gamma) \leq \sup_{x \in X} \inf_{\gamma \in Y} \phi(x, \gamma), \quad (3.23)$$

then $\inf_{y \in Y} \sup_{x \in X} \varphi(x, y) \leq \sup_{x \in X} \inf_{y \in Y} \phi(x, y)$.

PROOF. Letting $z^* = \inf_{y \in Y} \sup_{x \in X} \varphi(x, y)$, $z_* = \sup_{x \in X} \inf_{y \in Y} \phi(x, y)$, if $z^* > z_*$, then by the density of E , there exists a $\bar{z} \in E$ such that $z^* > \bar{z} > z_*$. Defining $F(y) = \{x \in X : \phi(x, y) \geq \bar{z}\}$, $y \in Y$, since $x \mapsto \phi(x, y)$ is upper semi-continuous, we know that $F(y)$ is closed for all $y \in Y$. Defining $D = \bigcap_{y \in Y_0} F(y)$, then D is closed.

On the other hand, for any finite set $\{y_1, \dots, y_n\} \subset Y$, it follows from condition (2) that there exists a compact subset M which contains $Y_0 \cup \{y_1, \dots, y_n\}$. Obviously, we can prove that $\varphi : X \times M \rightarrow E$ satisfies all the conditions in [Theorem 3.1](#), we have

$$\inf_{y \in M} \sup_{x \in X} \varphi(x, y) = \sup_{x \in X} \inf_{y \in M} \varphi(x, y), \quad (3.24)$$

and so we have

$$\begin{aligned} \sup_{x \in X} \inf_{y \in M} \phi(x, y) &\geq \sup_{x \in X} \inf_{y \in M} \varphi(x, y) = \inf_{y \in M} \sup_{x \in X} \varphi(x, y) \\ &\geq \inf_{y \in Y} \sup_{x \in X} \varphi(x, y) = z^* > \bar{z}. \end{aligned} \quad (3.25)$$

By the same method as in [Theorem 3.11](#), we can prove $z_* \geq \bar{z}$. This contradicts the choice of \bar{z} . Therefore we have $z^* \leq z_*$. This completes the proof. \square

From [Theorem 3.1](#) we can obtain the following theorem.

THEOREM 3.15. *Let X be a compact generalized interval space, E a completely dense linear ordered space, and $\varphi : X \times X \rightarrow E$ a lower semi-continuous function satisfying the following conditions:*

- (1) $x \mapsto \varphi(x, y)$ is T -quasi-concave;
- (2) for any $A \in \mathcal{I}(X)$ and for any $\alpha \in E$, $\bigcap_{x \in A} \{y \in X : \varphi(x, y) \leq \alpha\} \in \mathcal{K}(X)$,

then

- (1) there exists an $\bar{x} \in X$ such that $\varphi(\bar{x}, \bar{x}) > \theta$; or
- (2) there exists an $\bar{y} \in X$ such that $\varphi(x, \bar{y}) \leq \theta$ for all $x \in X$.

PROOF. Suppose the conclusion (1) is not true, then for any $x \in X$ we have $\varphi(x, x) \leq \theta$. By [Theorem 3.1](#) the following equation holds:

$$\sup_{x \in X} \inf_{y \in X} \varphi(x, y) = \inf_{y \in X} \sup_{x \in X} \varphi(x, y). \quad (3.26)$$

Since for any $x \in X$, $\inf_{y \in X} \varphi(x, y) \leq \varphi(x, x) \leq \theta$, we have $\sup_{x \in X} \inf_{y \in X} \varphi(x, y) \leq \theta$, and so we have $\inf_{y \in X} \sup_{x \in X} \varphi(x, y) \leq \theta$. By the compactness of X and the lower semi-continuity of $\sup_{x \in X} \varphi(x, y)$ in y , there exists a $\bar{y} \in X$ such that

$$\sup_{x \in X} \varphi(x, \bar{y}) = \inf_{y \in X} \sup_{x \in X} \varphi(x, y) \leq \theta, \quad (3.27)$$

that is, $\varphi(x, \bar{y}) \leq \theta$ for all $x \in X$. Therefore the conclusion (2) is true. \square

4. Application to coincidence problem and fixed point problem

THEOREM 4.1. *Let X be a compact generalized interval space, Y a topological space, $F : X \rightarrow 2^Y$ a mapping with closed values, and $s \in C(X, Y)$ a given mapping. If the*

following conditions are satisfied:

- (1) for any $x \in X$, $s^{-1}F(x) \neq \emptyset$ and $F^{-1}(s(x))$ is open;
- (2) for any $A \in \mathcal{A}(X)$, $X \setminus (\bigcup_{x \in A} F^{-1}(s(x))) \in \mathcal{K}(X)$;
- (3) for any $x_1, x_2 \in X$ and for any $y \in \bigcap_{i=1}^2 F^{-1}(s(x_i))$, $s(\Gamma(x_1, x_2)) \subset F(y)$ there exist $x'_1, x'_2 \in \Gamma(x_1, x_2)$ such that $F^{-1}(s(x_i)) \subset F^{-1}(s(x'_i))$, $i = 1, 2$,

then there exists an $\bar{x} \in X$ such that $s(\bar{x}) \in F(\bar{x})$.

PROOF. For any $(x, z) \in X \times X$, let

$$f(x, z) = \begin{cases} 0, & s(x) \notin F(z), \\ 1, & s(x) \in F(z). \end{cases} \tag{4.1}$$

If the conclusion is not true, then for any $x \in X$ we have $f(x, x) = 0$.

Since for any $x \in X$ and for any $r \in \mathbb{R}$,

$$\{z \in X : f(x, z) \leq r\} = \begin{cases} \emptyset, & r < 0, \\ F^*(s(x)), & 0 \leq r < 1, \\ X, & r \geq 1, \end{cases} \tag{4.2}$$

where $F^*(x) = X \setminus F^{-1}(x)$, it follows from conditions (1) and (2) that $f(x, z)$ is lower semi-continuous in z and for any $A \in \mathcal{A}(X)$ and for any $r \in \mathbb{R}$, $\bigcap_{x \in A} \{z \in X : f(x, z) < r\} \in \mathcal{K}(X)$. On the other hand, since for any $z \in X$ and for any $r \in \mathbb{R}$,

$$\{x \in X : f(x, z) \geq r\} = \begin{cases} X, & r \leq 0, \\ s^{-1}(F(z)), & 0 < r \leq 1, \\ \emptyset, & r > 1, \end{cases} \tag{4.3}$$

and F is a mapping with closed values, $f(x, z)$ is upper semi-continuous in x .

Letting $G(x) = F^*(s(x))$, then G is a mapping from X into 2^X and for any $(x, z) \in X \times X$,

$$f(x, z) = \begin{cases} 0, & z \in G(x), \\ 1, & z \notin G(x). \end{cases} \tag{4.4}$$

Next we prove that G is a T -KKM mapping. In fact, from condition (3) for any $x_1, x_2 \in X$, we have $s(\Gamma(x_1, x_2)) \subset F(z)$ for all $z \in \bigcap_{i=1}^2 F^{-1}(s(x_i))$. Hence for any $x \in \Gamma(x_1, x_2)$, and for any $z \in \bigcap_{i=1}^2 F^{-1}(s(x_i))$, $s(x) \in F(z)$.

Therefore, for all $x \in \Gamma(x_1, x_2)$ and for any $z \in \bigcap_{i=1}^2 F^{-1}(s(x_i))$, we have $z \in F^{-1}(s(x))$. This implies that $\bigcap_{i=1}^2 F^{-1}(s(x_i)) \subset F^{-1}(s(x))$ for all $x \in \Gamma(x_1, x_2)$. Therefore, $G(x) \subset \bigcup_{i=1}^2 G(x_i)$ for all $x \in \Gamma(x_1, x_2)$. Again by condition (3) there exist $x'_1, x'_2 \in \Gamma(x_1, x_2)$ such that $F^{-1}(s(x_i)) \subset F^{-1}(s(x'_i))$, $i = 1, 2$. Thus $G(x_i) \subset G(x'_i)$, $i = 1, 2$. This shows that G is a T -KKM mapping. By [Lemma 1.8](#) it is easy to prove that $f(x, y)$ is a T -quasi-concave mapping in x . Since X is compact, f satisfies all the conditions in [Corollary 3.7](#). By [Corollary 3.7](#) there exists a $\bar{z} \in X$ such that $f(x, \bar{z}) \leq 0$ for all $x \in X$, and so $f(x, \bar{z}) = 0$ for all $x \in X$, that is, $s(x) \notin F(\bar{z})$ for all $x \in X$. Hence $x \notin s^{-1}F(\bar{z})$ for all $x \in X$, and so $s^{-1}F(\bar{z}) = \emptyset$.

This contradicts condition (1). This completes the proof. □

From [Theorem 4.1](#) we obtain the following fixed point theorem.

COROLLARY 4.2. *Let X be a generalized interval space, C be a compact T -convex subset of X , and $F : C \rightarrow 2^C$ be a set-valued mapping with nonempty closed values. If the following conditions are satisfied:*

- (1) *for any $x \in C$, $F^{-1}(x)$ is open in C ;*
- (2) *for any $A \in \mathcal{A}(C)$, $C \setminus (\bigcup_{x \in A} F^{-1}(x)) \in \mathcal{K}(C)$;*
- (3) *for any $x_1, x_2 \in X$, $\Gamma(x_1, x_2) \subset F(y)$ for all $y \in \bigcap_{i=1}^2 F^{-1}(x_i)$, and there exist $x'_1, x'_2 \in \Gamma(x_1, x_2)$ such that $F^{-1}(x_i) \subset F^{-1}(x'_i)$, $i = 1, 2$,*

then there exists an $\bar{x} \in C$ such that $\bar{x} \in F(\bar{x})$.

THEOREM 4.3. *Let X be a generalized interval space, Y a topological space, $F : X \rightarrow 2^Y$ and $s \in C(X, Y)$ two mappings satisfying the following conditions:*

- (1) *for any $x \in X$, $F^{-1}(s(x))$ is nonempty closed and $X \setminus s^{-1}(F(x))$ is T -convex;*
- (2) *for any $A \in \mathcal{A}(X)$, $\bigcap_{x \in A} s^{-1}(F(x)) \in \mathcal{K}(X)$;*
- (3) *$F^{-1} : Y \rightarrow 2^X$ is upper semi-continuous and for any $x_1, x_2 \in X$, there exist $x'_1, x'_2 \in \Gamma(x_1, x_2)$ such that $F^{-1}(s(x'_i)) \subset F^{-1}(s(x_i))$, $i = 1, 2$;*
- (4) *there exists an $A_0 \in \mathcal{A}(X)$ such that $\bigcap_{x \in A_0} F^{-1}(s(x))$ is compact,*

then there exists an $\bar{x} \in X$ such that $s(\bar{x}) \in F(\bar{x})$.

PROOF. Letting $G(x) = F^{-1}(s(x))$, then $G : X \rightarrow 2^X$ is a nonempty mapping and closed valued and G satisfies condition (2) in [Theorem 2.1](#).

On the other hand, for any $x \in X$, let W be an open set containing $G(x)$. Since F^{-1} is upper semi-continuous, there exists an open neighborhood V of $s(x)$ such that for any $y \in V$, we have $F^{-1}(y) \subset W$. Since $s \in C(x, y)$, there exists an open neighborhood U of x such that for any $z \in U$ we have $s(z) \in V$, and so for $z \in U$, we have $G(z) = F^{-1}(s(z)) \subset W$. This shows G is an upper semi-continuous mapping.

Next we prove that $G : X \rightarrow 2^X$ is a T -KKM mapping. In fact, for any $x_1, x_2 \in X$ and for any $x \in \Gamma(x_1, x_2)$, if $z \notin \bigcap_{i=1}^2 G(x_i)$, then $z \notin F^{-1}(s(x_i))$, $i = 1, 2$, that is, $s(x_i) \notin F(z)$, $i = 1, 2$, and so $\{x_1, x_2\} \subset X \setminus s^{-1}F(z)$. By the assumption that $X \setminus s^{-1}F(z)$ is T -convex, hence $\Gamma(x_1, x_2) \subset X \setminus s^{-1}F(z)$, and so $z \notin F^{-1}(s(x))$ for all $x \in \Gamma(x_1, x_2)$. This implies that

$$G(x) = F^{-1}(s(x)) \subset \bigcup_{i=1}^2 G(x_i). \quad (4.5)$$

Again by condition (3), there exist $x'_1, x'_2 \in \Gamma(x_1, x_2)$ such that $G(x'_i) \subset G(x_i)$, $i = 1, 2$. This means that G is a T -KKM mapping.

In view of condition (4), we know that G satisfies all the conditions in [Theorem 2.1](#). By [Theorem 2.1](#) $\bigcap_{x \in X} G(x) \neq \emptyset$, and so there exists an $\bar{x} \in X$ such that $\bar{x} \in G(x)$ for all $x \in X$. Hence $\bar{x} \in G(\bar{x})$, that is, $\bar{x} \in F^{-1}(s(\bar{x}))$, and so $s(\bar{x}) \in F(\bar{x})$. \square

THEOREM 4.4. *Let K be a generalized interval space, $F : X \rightarrow 2^X$ a mapping with nonempty values. If X is normal and the following conditions are satisfied:*

- (1) *F is transfer closed valued;*
- (2) *F is almost upper semi-continuous, that is, for any $x \in X$ and for any open set V containing $F(x)$, there exists an open neighborhood U of x such that $F(y) \subset \text{cl}V$ for all $y \in U$;*
- (3) *for any $A \in \mathcal{A}(X)$, $\bigcap_{x \in A} \text{cl}F(x) \in \mathcal{K}(X)$;*

- (4) *there exists an $A_0 \in \mathcal{D}(X)$ such that $\bigcap_{x \in A_0} \text{cl}F(x)$ is compact;*
- (5) *for any $x \in X$, $X \setminus F^{-1}(x) = F^\bullet(x)$ is T -convex,*

then F has a fixed point in X .

PROOF. For any $x \in X$, if W is any open set containing $\text{cl}F(x)$, by the normality of X , there exists an open set V containing $\text{cl}F(x)$ such that $\text{cl}V \subset W$. It follows from the almost upper semi-continuity of F that there exists an open neighborhood U of x such that $F(y) \subset \text{cl}V$ for all $y \in U$. Hence $\text{cl}F(y) \subset \text{cl}V \subset W$. This means that F is upper semi-continuous.

Next we prove that F is a T -KKM mapping. In fact, for any $x_1, x_2 \in X$ if $x \notin \bigcup_{i=1}^2 F(x_i)$, then $x_1, x_2 \notin F^{-1}(y)$. Since $F^\bullet(y)$ is T -convex, $\Gamma(x_1, x_2) \subset F^\bullet(y)$, and so for any $x \notin \Gamma(x_1, x_2)$ we have $x \notin F^{-1}(y)$, that is, $y \notin F(x)$. Therefore, $F(x) \subset \bigcup_{i=1}^2 F(x_i)$ for all $x \in \Gamma(x_1, x_2)$, and so $\text{cl}F(x) \subset \bigcup_{i=1}^2 \text{cl}F(x_i)$ for all $x \in \Gamma(x_1, x_2)$. In view of condition (5) we know that F is a T -KKM mapping.

Moreover, it is easy to prove that all conditions in [Theorem 2.2](#) are satisfied. By [Theorem 2.2](#), $\bigcap_{x \in X} F(x) \neq \emptyset$. Hence there exists an $\tilde{x} \in X$ such that $\tilde{x} \in F(x)$ for all $x \in X$, and so we have $\tilde{x} \in F(\tilde{x})$. This completes the proof. □

5. Application to economy equilibrium problem

THEOREM 5.1. *Let X be a generalized interval space, $A, B, P : X \rightarrow 2^X$. If the following conditions are satisfied:*

- (1) *for any $x \in X$, $A(x) \subset B(x)$, $x \notin P(x)$, $A(x)$, and $P(x)$ are nonempty closed sets;*
- (2) *A and P are upper semi-continuous;*
- (3) *for any $y \in X$, $A^\bullet(y) \cup P^\bullet(y)$ and $A^\bullet(y) \cup \{x \in X : x \in \text{cl}B(x)\}$ are T -convex sets;*
- (4) *for any $M, N \in \mathcal{I}(X)$, $(\bigcap_{x \in M} A(x)) \cap (\bigcap_{x \in N} P(x)) \in \mathcal{H}(X)$;*
- (5) *there exists $M_0 \in \mathcal{I}(X)$ such that $\bigcap_{x \in M_0} A(x)$ is compact,*

then (X, A, B, P) has a Shafer-Sonnenschein equilibrium, that is, there exists an $\tilde{x} \in X$ such that $\tilde{x} \in \text{cl}B(\tilde{x})$ and $A(\tilde{x}) \cap P(\tilde{x}) \neq \emptyset$.

PROOF. Suppose the contrary, for any $x \in X$ we have $x \notin \text{cl}B(x)$ or $A(x) \cap P(x) = \emptyset$. Letting

$$D = \{x \in X : x \in \text{cl}B(x)\},$$

$$F(x) = \begin{cases} A(x) \cap P(x), & x \in D, \\ A(x), & x \notin D. \end{cases} \tag{5.1}$$

By condition (1) and (2), F is an upper semi-continuous mapping with nonempty closed values. For any $y \in X$,

$$F^{-1}(y) = \{x \in X : y \in F(x)\}$$

$$= \{x \in X : x \in A^{-1}(y) \cap P^{-1}(y)\} \cup \{x \notin D : x \in A^{-1}(y)\},$$

$$F^\bullet(y) = \{x \in X : x \in A^\bullet(y) \cup P^\bullet(y)\} \cap \{x \in X : x \in A^\bullet(y) \cup D\}$$

$$= (A^\bullet(y) \cup P^\bullet(y)) \cap (A^\bullet(y) \cup D). \tag{5.2}$$

By condition (3), $F^*(\gamma)$ is a T -convex set; and by condition (4), $\bigcap_{x \in M} F(x) \in \mathcal{H}(X)$. It is obvious that F satisfies all conditions in [Theorem 2.1](#). By [Theorem 2.1](#), there exists a $\bar{y} \in X$ such that $\bar{y} \in F(\bar{y})$. If $\bar{y} \notin D$, then $\bar{y} \in A(\bar{y}) \subset B(\bar{y})$ which is impossible; if $\bar{y} \in D$, then $\bar{y} \in P(\bar{y})$ which contradicts condition (1). Therefore, there exists an $\bar{x} \in X$ such that $\bar{x} \in \text{cl}B(\bar{x})$ and $A(\bar{x}) \cap P(\bar{x}) \neq \emptyset$. \square

REFERENCES

- [1] H. Brézis, L. Nirenberg, and G. Stampacchia, *A remark on Ky Fan's minimax principle*, Boll. Un. Mat. Ital. (4) **6** (1972), 293–300. [MR 48#2850](#). [Zbl 264.49013](#).
- [2] C. Z. Cheng and Y. H. Lin, *A generalization of minimax theorems of fixed-point type*, Acta Math. Sinica **34** (1991), no. 4, 502–507. [MR 93c:49008](#). [Zbl 0833.49008](#).
- [3] K. Fan, *Fixed-point and minimax theorems in locally convex topological linear spaces*, Proc. Nat. Acad. Sci. U.S.A. **38** (1952), 121–126. [MR 13,858d](#). [Zbl 0047.35103](#).
- [4] M. A. Geraghty and B.-L. Lin, *Topological minimax theorems*, Proc. Amer. Math. Soc. **91** (1984), no. 3, 377–380. [MR 86c:49009](#). [Zbl 533.90095](#).
- [5] C. W. Ha, *A noncompact minimax theorem*, Pacific J. Math. **97** (1981), no. 1, 115–117. [MR 83g:49024](#). [Zbl 474.49015](#).
- [6] J. Hartung, *An extension of Sion's minimax theorem with an application to a method for constrained games*, Pacific J. Math. **103** (1982), no. 2, 401–408. [MR 84k:90101](#). [Zbl 534.49009](#).
- [7] T. Hoang, *On a general minimax theorem*, Soviet Math. Dokl. **15** (1974), 1689–1693 (Russian). [Zbl 314.90088](#).
- [8] J. Kindler and R. Trost, *Minimax theorems for interval spaces*, Acta Math. Hungar. **54** (1989), no. 1-2, 39–49. [MR 91f:49009](#). [Zbl 688.90061](#).
- [9] V. Komornik, *Minimax theorems for upper semicontinuous functions*, Acta Math. Acad. Sci. Hungar. **40** (1982), no. 1-2, 159–163. [MR 84h:49023](#). [Zbl 505.49007](#).
- [10] B.-L. Lin and X. C. Quan, *A noncompact topological minimax theorem*, J. Math. Anal. Appl. **161** (1991), no. 2, 587–590. [MR 92i:49011](#). [Zbl 779.49009](#).
- [11] ———, *A two functions nonlinear minimax theorem*, Fixed Point Theory and Applications (Marseille, 1989), Pitman Res. Notes Math. Ser., vol. 252, Longman Scientific & Technical, Harlow, 1991, pp. 321–325. [MR 92j:54023](#). [Zbl 771.49003](#).
- [12] M. Sion, *On general minimax theorems*, Pacific J. Math. **8** (1958), 171–176. [MR 20#3506](#). [Zbl 081.11502](#).
- [13] L. L. Stachó, *Minimax theorems beyond topological vector spaces*, Acta Sci. Math. (Szeged) **42** (1980), no. 1-2, 157–164. [MR 82a:49017](#). [Zbl 436.49017](#).
- [14] G. Q. Tian, *Generalizations of the FKKM theorem and the Ky Fan minimax inequality, with applications to maximal elements, price equilibrium, and complementarity*, J. Math. Anal. Appl. **170** (1992), no. 2, 457–471. [MR 94b:90014](#). [Zbl 767.49007](#).
- [15] W.-C. Wu, *A remark on the fundamental theorem in the theory of games*, Sci. Record (N.S.) **3** (1959), 229–233. [MR 22#13311](#).

DA-CHENG WANG: DEPARTMENT OF COMPUTERS, CHONGQING JIAOTONG INSTITUTE, CHONGQING 400074, CHINA

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk