

INTEGRAL MEAN ESTIMATES FOR POLYNOMIALS WHOSE ZEROS ARE WITHIN A CIRCLE

K. K. DEWAN, ABDULLAH MIR, and R. S. YADAV

(Received 3 November 2000)

ABSTRACT. Let $p(z)$ be a polynomial of degree n having all its zeros in $|z| \leq k$; $k \leq 1$, then for each $r > 0$, $p > 1$, $q > 1$ with $p^{-1} + q^{-1} = 1$, Aziz and Ahemad (1996) recently proved that $n \left\{ \int_0^{2\pi} |p(e^{i\theta})|^r d\theta \right\}^{1/r} \leq \left\{ \int_0^{2\pi} |1 + ke^{i\theta}|^{pr} d\theta \right\}^{1/pr} \left\{ \int_0^{2\pi} |p'(e^{i\theta})|^{qr} d\theta \right\}^{1/qr}$. In this paper, we extend the above inequality to the class of polynomials $p(z) = a_n z^n + \sum_{v=\mu}^n a_{n-v} z^{n-v}$; $1 \leq \mu \leq n$ having all its zeros in $|z| \leq k$; $k \leq 1$ and obtain a generalization as well as a refinement of the above result.

2000 Mathematics Subject Classification. 30A10, 30C10, 30C15.

1. Introduction and statement of results. Let $p(z)$ be a polynomial of degree n and $p'(z)$ its derivative. If $p(z)$ has all its zeros in $|z| \leq 1$, then it was shown by Turan [7] that

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{2} \max_{|z|=1} |p(z)|. \quad (1.1)$$

Inequality (1.1) is best possible with equality for $p(z) = \alpha z^n + \beta$, where $|\alpha| = |\beta|$. As an extension of (1.1) Malik [4] proved that if $p(z)$ has all its zeros in $|z| \leq k$, where $k \leq 1$, then

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{1+k} \max_{|z|=1} |p(z)|. \quad (1.2)$$

Malik [5] obtained a generalization of (1.1) in the sense that the right-hand side of (1.1) is replaced by a factor involving the integral mean of $|p(z)|$ on $|z| = 1$. In fact he proved the following theorem.

THEOREM 1.1. *If $p(z)$ has all its zeros in $|z| \leq 1$, then for each $r > 0$*

$$n \left\{ \int_0^{2\pi} |p(e^{i\theta})|^r d\theta \right\}^{1/r} \leq \left\{ \int_0^{2\pi} |1 + e^{i\theta}|^r d\theta \right\}^{1/r} \max_{|z|=1} |p'(z)|. \quad (1.3)$$

The result is sharp and equality in (1.3) holds for $p(z) = (z+1)^n$.

If we let $r \rightarrow \infty$ in (1.3) we get (1.1). Aziz and Ahemad [1] generalized (1.3) in the sense that $\max_{|z|=1} |p'(z)|$ on $|z| = 1$ on the right-hand side of (1.3) is replaced by a factor involving the integral mean of $|p'(z)|$ on $|z| = 1$ and proved the following result.

THEOREM 1.2. *If $p(z)$ is a polynomial of degree n having all its zeros in $|z| \leq k \leq 1$, then for $r > 0, p > 1, q > 1$ with $1/p + 1/q = 1$,*

$$n \left\{ \int_0^{2\pi} |p(e^{i\theta})|^r d\theta \right\}^{1/r} \leq \left\{ \int_0^{2\pi} |1 + ke^{i\theta}|^{qr} d\theta \right\}^{1/qr} \left\{ \int_0^{2\pi} |p'(e^{i\theta})|^{pr} d\theta \right\}^{1/pr}. \tag{1.4}$$

If we let $r \rightarrow \infty$ and $p \rightarrow \infty$ (so that $q \rightarrow 1$) in (1.4) we get (1.2).

In this paper, we will first extend Theorem 1.2 to the class of polynomials $p(z) = a_n z^n + \sum_{v=\mu}^n a_{n-v} z^{n-v}, 1 \leq \mu \leq n$, having all the zeros in $|z| \leq k; k \leq 1$, and thereby obtain a generalization of it. More precisely, we prove the following result.

THEOREM 1.3. *If $p(z) = a_n z^n + \sum_{v=\mu}^n a_{n-v} z^{n-v}, 1 \leq \mu \leq n$ is a polynomial of degree n having all its zeros in $|z| \leq k; k \leq 1$, then for each $r > 0, p > 1, q > 1$ with $1/p + 1/q = 1$,*

$$n \left\{ \int_0^{2\pi} |p(e^{i\theta})|^r d\theta \right\}^{1/r} \leq \left\{ \int_0^{2\pi} |1 + k^\mu e^{i\theta}|^{pr} d\theta \right\}^{1/pr} \left\{ \int_0^{2\pi} |p'(e^{i\theta})|^{qr} d\theta \right\}^{1/qr}. \tag{1.5}$$

REMARK 1.4. If we let $r \rightarrow \infty$ and $q \rightarrow \infty$ (so that $p \rightarrow 1$) in (1.5) we get (1.2) for $\mu = 1$.

Our next result is an improvement of Theorem 1.3 which in turn gives a generalization as well as a refinement of Theorem 1.2.

THEOREM 1.5. *If $p(z) = a_n z^n + \sum_{v=\mu}^n a_{n-v} z^{n-v}, 1 \leq \mu \leq n$, is a polynomial of degree n having all its zeros in $|z| \leq k; k \leq 1$ and $m = \min_{|z|=k} |p(z)|$, then for every real or complex number β with $|\beta| \leq 1, r > 0, p > 1, q > 1$ with $1/p + 1/q = 1$,*

$$\begin{aligned} n \left\{ \int_0^{2\pi} \left| p(e^{i\theta}) + \frac{\beta m e^{i(n-1)\theta}}{k^{n-\mu}} \right|^r d\theta \right\}^{1/r} \\ \leq \left\{ \int_0^{2\pi} |1 + k^\mu e^{i\theta}|^{pr} d\theta \right\}^{1/pr} \left\{ \int_0^{2\pi} |p'(e^{i\theta})|^{qr} d\theta \right\}^{1/qr}. \end{aligned} \tag{1.6}$$

REMARK 1.6. Letting $r \rightarrow \infty$ and $q \rightarrow \infty$ (so that $p \rightarrow 1$) in (1.6) and choosing the argument of β suitably with $|\beta| = 1$, it follows that, if $p(z) = a_n z^n + \sum_{v=\mu}^n a_{n-v} z^{n-v}, 1 \leq \mu \leq n$, is a polynomial of degree n having all its zeros in $|z| \leq k; k \leq 1$, then

$$\max_{|z|=1} |p'(z)| \geq \frac{n}{1 + k^\mu} \left\{ \max_{|z|=1} |p(z)| + \frac{1}{k^{n-\mu}} \min_{|z|=1} |p(z)| \right\}. \tag{1.7}$$

Inequality (1.7) was recently proved by Aziz and Shah [2].

2. Lemmas. For the proof of Theorem 1.5 we will make use of the following lemmas.

LEMMA 2.1 (see Aziz and Shah [2, Lemma 2]). *If $p(z) = a_n z^n + \sum_{v=\mu}^n a_{n-v} z^{n-v}$ is a polynomial of degree n having all its zeros in $|z| \leq k \leq 1$, then*

$$|q'(z)| \leq k^\mu |p'(z)| \quad \text{for } |z| = 1, 1 \leq \mu \leq n, \tag{2.1}$$

where here and throughout $q(z) = z^n \overline{p(1/\bar{z})}$.

LEMMA 2.2 (see Rather [6]). *If $p(z) = a_n z^n + \sum_{v=\mu}^n a_{n-v} z^{n-v}$, $1 \leq \mu \leq n$, is a polynomial of degree n having all its zeros in $|z| \leq k \leq 1$, and $m = \min_{|z|=k} |p(z)|$, then*

$$k^\mu |p'(z)| \geq |q'(z)| + \frac{mn}{k^{n-\mu}} \quad \text{for } |z| = 1. \tag{2.2}$$

3. Proof of theorems

PROOF OF THEOREM 1.3. Suppose that $p(z)$ has all its zeros in $|z| \leq k \leq 1$, therefore, by Lemma 2.1 we have

$$k^\mu |p'(z)| \geq |q'(z)| \quad \text{for } |z| = 1. \tag{3.1}$$

Also $q(z) = z^n \overline{p(1/\bar{z})}$ so that $p(z) = z^n \overline{q(1/\bar{z})}$, we have

$$p'(z) = n z^{n-1} \overline{q\left(\frac{1}{\bar{z}}\right)} - z^{n-2} \overline{q'\left(\frac{1}{\bar{z}}\right)}. \tag{3.2}$$

Equivalently,

$$z p'(z) = n z^n \overline{q\left(\frac{1}{\bar{z}}\right)} - z^{n-1} \overline{q'\left(\frac{1}{\bar{z}}\right)} \tag{3.3}$$

which implies

$$|p'(z)| = |nq(z) - zq'(z)| \quad \text{for } |z| = 1. \tag{3.4}$$

Using (3.1) in (3.4) we get

$$|q'(z)| \leq k^\mu |nq(z) - zq'(z)| \quad \text{for } |z| = 1; 1 \leq \mu \leq n. \tag{3.5}$$

Since $p(z)$ has all its zeros in $|z| \leq k \leq 1$, by the Gauss-Lucas theorem all the zeros of $p'(z)$ also lie in $|z| \leq 1$. This implies that the polynomial

$$z^{n-1} \overline{p'\left(\frac{1}{\bar{z}}\right)} = nq(z) - zq'(z) \tag{3.6}$$

has all its zeros in $|z| \geq 1/k \geq 1$.

Therefore, it follows from (3.5) that the function

$$w(z) = \frac{zq'(z)}{k^\mu (nq(z) - zq'(z))} \tag{3.7}$$

is analytic for $|z| \leq 1$ and $|w(z)| \leq 1$ for $|z| \leq 1$. Furthermore $w(0) = 0$. Thus the function $1 + k^\mu w(z)$ is subordinate to the function $1 + k^\mu z$ in $|z| \leq 1$. Hence by a well-known property of subordination [3] we have for $r > 0$ and for $0 \leq \theta < 2\pi$,

$$\int_0^{2\pi} |1 + k^\mu w(e^{i\theta})|^r d\theta \leq \int_0^{2\pi} |1 + k^\mu e^{i\theta}|^r d\theta. \tag{3.8}$$

Also from (3.7), we have

$$1 + k^\mu w(z) = \frac{nq(z)}{nq(z) - zq'(z)} \tag{3.9}$$

or

$$|nq(z)| = |1 + k^\mu w(z)| |nq(z) - zq'(z)|. \tag{3.10}$$

Using (3.4) and also $|p(z)| = |q(z)|$ in (3.10), we have

$$n|p(z)| = |1 + k^\mu w(z)| |p'(z)| \quad \text{for } |z| = 1. \tag{3.11}$$

Combining (3.8) and (3.11) we get

$$n^r \int_0^{2\pi} |p(e^{i\theta})|^r d\theta \leq \int_0^{2\pi} |1 + k^\mu e^{i\theta}|^r |p'(e^{i\theta})|^r d\theta \quad \text{for } r > 0. \tag{3.12}$$

Now applying Hölder's inequality for $p > 1, q > 1$ with $1/p + 1/q = 1$ to (3.12), we get

$$\begin{aligned} n^r \int_0^{2\pi} |p(e^{i\theta})|^r d\theta &\leq \left\{ \int_0^{2\pi} |1 + k^\mu e^{i\theta}|^{pr} d\theta \right\}^{1/p} \left\{ \int_0^{2\pi} |p'(e^{i\theta})|^{qr} d\theta \right\}^{1/q} \quad \text{for } r > 0 \end{aligned} \tag{3.13}$$

which is equivalent to

$$\begin{aligned} n \left\{ \int_0^{2\pi} |p(e^{i\theta})|^r d\theta \right\}^{1/r} &\leq \left\{ \int_0^{2\pi} |1 + k^\mu e^{i\theta}|^{pr} d\theta \right\}^{1/pr} \left\{ \int_0^{2\pi} |p'(e^{i\theta})|^{qr} d\theta \right\}^{1/qr} \quad \text{for } r > 0 \end{aligned} \tag{3.14}$$

which proves the desired result. □

PROOF OF THEOREM 1.5. Since $p(z)$ has all its zeros in $|z| \leq k \leq 1$, therefore, by Lemma 2.2 we get

$$k^\mu |p'(z)| \geq |q'(z)| + \frac{mn}{k^{n-\mu}} \quad \text{for } |z| = 1, 1 \leq \mu \leq n. \tag{3.15}$$

Also by (3.4) for $|z| = 1$, we have

$$|p'(z)| = |nq(z) - zq'(z)|. \tag{3.16}$$

Now using (3.15) for every complex β with $|\beta| \leq 1$, we get

$$\begin{aligned} \left| q'(z) + \beta \frac{mn}{k^{n-\mu}} \right| &\leq |q'(z)| + \frac{mn}{k^{n-\mu}} \\ &\leq k^\mu |p'(z)| \\ &= k^\mu |nq(z) - zq'(z)| \quad \text{for } |z| = 1. \end{aligned} \tag{3.17}$$

Since $p(z)$ has all its zeros in $|z| \leq k \leq 1$, the result follows on the same lines as that of [Theorem 1.3](#). Hence we omit the proof. \square

REFERENCES

- [1] A. Aziz and N. Ahemad, *Integral mean estimates for polynomials whose zeros are within a circle*, Glas. Mat. Ser. III **31(51)** (1996), no. 2, 229–237. [MR 98d:30008](#). [Zbl 874.30001](#).
- [2] A. Aziz and W. M. Shah, *An integral mean estimate for polynomials*, Indian J. Pure Appl. Math. **28** (1997), no. 10, 1413–1419. [MR 99a:30005](#). [Zbl 895.30004](#).
- [3] E. Hille, *Analytic Function Theory. Vol. II*, Introductions to Higher Mathematics, Ginn and Company, New York, 1962. [MR 34#1490](#). [Zbl 102.29401](#).
- [4] M. A. Malik, *On the derivative of a polynomial*, J. London Math. Soc. (2) **1** (1969), 57–60. [MR 40#2827](#). [Zbl 179.37901](#).
- [5] ———, *An integral mean estimate for polynomials*, Proc. Amer. Math. Soc. **91** (1984), no. 2, 281–284. [MR 85c:30007](#). [Zbl 543.30002](#).
- [6] N. A. Rather, *Extremal properties and location of the zeros of polynomials*, Ph.D. thesis, University of Kashmir, 1988.
- [7] P. Turan, *Über die Ableitung von Polynomen*, Compositio Math. **7** (1939), 89–95 (German). [MR 1,37b](#).

K. K. DEWAN: DEPARTMENT OF MATHEMATICS, FACULTY OF NATURAL SCIENCES, JAMIA MILLIA ISLAMIA, NEW DELHI-110025, INDIA
E-mail address: kkdewan123@rediffmail.com

ABDULLAH MIR AND R. S. YADAV: DEPARTMENT OF MATHEMATICS, FACULTY OF NATURAL SCIENCES, JAMIA MILLIA ISLAMIA, NEW DELHI-110025, INDIA

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru