

ON SOME PROPERTIES OF THE LÜROTH-TYPE ALTERNATING SERIES REPRESENTATIONS FOR REAL NUMBERS

C. GANATSIU

(Received 3 November 2000)

ABSTRACT. We investigate some properties connected with the alternating Lüroth-type series representations for real numbers, in terms of the integer digits involved. In particular, we establish the analogous concept of the asymptotic density and the distribution of the maximum of the first n denominators, by applying appropriate limit theorems.

2000 Mathematics Subject Classification. 60A10, 11K55, 37A30, 37A50, 60K99, 60G10, 60J10.

1. Introduction. Let x be any number in the interval $I := (0, 1)$. Then by using a general alternating series algorithm introduced by Knopfmacher and Knopfmacher [6, 7], analogous to a positive one of Oppenheim [8], we may prove that x has a unique finite or infinite representation in the form

$$\begin{aligned} x &= \frac{1}{\alpha_1} - \frac{1}{(\alpha_1 + 1)\alpha_1} \frac{1}{\alpha_2} + \frac{1}{(\alpha_1 + 1)\alpha_1(\alpha_2 + 1)\alpha_2} \frac{1}{\alpha_3} - \dots \\ &= ((\alpha_1, \alpha_2, \dots, \alpha_n, \dots)), \end{aligned} \tag{1.1}$$

where $\alpha_n \geq 1$, $n \geq 1$.

Representation (1.1) is called *Lüroth-type* alternating expansion, while the positive integers $\alpha_1, \alpha_2, \dots, \alpha_n, \dots$ are called the *digits* (or the *denominators*) of the above mentioned expansion. It is obvious that the digits α_n are functions $\alpha_n(x)$ of x ; therefore, it can be easily seen that the α_n 's may be considered as random variables defined almost surely on I with respect to any probability measure on the σ -algebra B_I (in particular, with respect to the Lebesgue measure λ).

A lot of research has been carried out into the ergodic properties of the denominators in the Lüroth expansions of real numbers in $(0, 1)$. In particular, it was studied independently by Šalát [9] and Jager and de Vroedt [3] not only the stochastic behaviour of the digits d_n (which are independent random variables on a probability space S , where the basic set is the unit interval $(0, 1)$ and the probability is the Lebesgue measure) but some other important metric properties concerning the sequence $\{d_n\}_n$.

Similar further results were derived later by Kalpazidou, Knopfmacher, and Knopfmacher (see [4, 5]), for the alternating Lüroth-type series.

The aim of the present paper is to give some sharper properties for the alternating Lüroth-type expansions related with the order of magnitude of the digits α_n . For the ordinary Lüroth expansions, an analogous problem has been investigated by Šalát [9].

2. Preliminaries. Let

$$I_n \equiv I_n(k_1, \dots, k_n) = \{x \in I \mid \alpha_1(x) = k_1, \dots, \alpha_n(x) = k_n\}, \quad \text{for any } k_1, k_2, \dots, k_n \in \mathbb{N}^*, \tag{2.1}$$

be the set of all $x \in I$ which have a unique expansion of the form (1.1) such that the digits $\alpha_1(x), \dots, \alpha_n(x)$ have the concrete values k_1, \dots, k_n .

Then according to a result of [5] concerning the stochastic behaviour of the α_n 's, we have the following proposition.

PROPOSITION 2.1. *The digits $\alpha_n(\cdot), n \in \mathbb{N}^*$, are stochastically independent and identically distributed random variables with respect to Lebesgue measure λ , with*

$$\lambda(\alpha_n = k) = \frac{1}{k(k+1)}, \quad k \in \mathbb{N}^*. \tag{2.2}$$

Evidently the mean value of the digits α_n is given by

$$E(\alpha_n) = \sum_{k=1}^{+\infty} k\lambda(\alpha_n = k) = \sum_{k=1}^{+\infty} \frac{1}{k+1} = +\infty, \tag{2.3}$$

which means that the usual limit theorems do not apply.

Then, according to an interesting result which is a kind of converse to the strong law of large numbers due to Chow and Robbins [1], it may be obtained that, for a.a. x ,

$$\limsup \frac{(\alpha_1 + \alpha_2 + \dots + \alpha_n)}{n} = +\infty. \tag{2.4}$$

It is obvious that if we take the functions $u(\alpha_n)$ of α_n for which $\sum_{k=1}^{+\infty} u(\alpha_n)/k(k+1) < +\infty$, then we can apply the usual theorems, obtaining strong laws and asymptotic normality.

In Section 3, we investigate some sharper results for the digits α_n of the alternating Lüroth-type series following the spirit of Šalát [9], while in Section 4, we investigate the asymptotic behaviour of $M_n = M_n(x) = \max(\alpha_1, \dots, \alpha_n)$ by using Proposition 2.1.

3. Some remarks on the digits of the alternating Lüroth series. Let $\{c_n\}_n$ be an arbitrarily chosen sequence of real numbers and X_n the indicator variable of the fact $\{x \mid \alpha_n(x) > c_n(x)\}$, that is,

$$X_n = \begin{cases} 1, & \text{if } \alpha_n > c_n, \\ 0, & \text{otherwise.} \end{cases} \tag{3.1}$$

By applying Proposition 2.1 we obtain that X_n are independent random variables and; moreover,

$$P(X_n = 1) = \sum_{k=[c_n]+1}^{+\infty} \frac{1}{k(k+1)} = \sum_{k=[c_n]+1}^{+\infty} \left(\frac{1}{k} - \frac{1}{(k+1)} \right) = \frac{1}{[c_n]+1}, \tag{3.2}$$

$$P(X_n = 0) = 1 - P(X_n = 1) = 1 - \frac{1}{[c_n]+1},$$

if we assume that $c_n \geq 0$. (This assumption does not restrict the generality of our investigation, since from the general alternating series algorithm we have $\alpha_n \geq 1$, for all n .) At first we prove the following theorems.

THEOREM 3.1. *The series $\sum X_n$ converges a.e. if and only if*

$$\sum \frac{1}{c_n + 1} < +\infty. \tag{3.3}$$

Moreover, if the mean value

$$E_N = \sum_{n=1}^N \frac{1}{[c_n] + 1} \rightarrow +\infty, \tag{3.4}$$

then by setting

$$Z_N = \sum_{n=1}^N X_n, \quad V_N = \sum_{n=1}^N \frac{1}{[c_n] + 1} \cdot \left(1 - \frac{1}{[c_n] + 1}\right), \tag{3.5}$$

we take

$$\lambda(Z_N - E_N < z\sqrt{V_N}) \rightarrow \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} \cdot e^{-t^2/2} dt \tag{3.6}$$

as $N \rightarrow +\infty$, in case $V_N \rightarrow +\infty$.

PROOF. The first part of the above theorem is a consequence of the Borel-Cantelli lemma (see [2]). Furthermore, since

$$E(X_n) = \frac{1}{[c_n] + 1}, \quad \text{Var}(X_n) = \frac{1}{[c_n] + 1} \cdot \left(1 - \frac{1}{[c_n] + 1}\right), \tag{3.7}$$

we may apply the central limit theorem under Lindeberg's conditions, that is,

$$\lambda\left(-\infty < \frac{Z_N - E_N}{\sqrt{V_N}} < z\right) \xrightarrow{N \rightarrow +\infty} \int_{-\infty}^z f(t) dt = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} \cdot e^{-t^2/2} dt \tag{3.8}$$

and the proof is complete. □

Relation (3.6) implies that Z_N converges in probability to E_N if both E_N and V_N tend to $+\infty$. Although in the general case much more important information may be derived, we take in a special case the following theorem.

THEOREM 3.2. *Assume that*

$$\lim_{N \rightarrow +\infty} \frac{E_N}{N} = c \tag{3.9}$$

exists and is positive, then for a.a. x in $(0, 1)$

$$\lim_{N \rightarrow +\infty} \frac{(X_1 + X_2 + \dots + X_N)}{N} = c. \tag{3.10}$$

The statement holds also in case $c = 0$.

PROOF. We define the random variables K_n by the relation $K_n = X_n - 1/([c_n] + 1)$. Consequently, we have

$$\begin{aligned} E(K_n) &= E(X_n) - \frac{1}{[c_n] + 1} = 0, \\ V(K_n) &= \text{Var}(X_n) = \frac{1}{[c_n] + 1} \cdot \left(1 - \frac{1}{[c_n] + 1}\right). \end{aligned} \tag{3.11}$$

By using the Kolmogorov's theorem [2] we obtain that, for a.a. x in $(0, 1)$,

$$\lim_{N \rightarrow +\infty} \frac{(K_1 + \dots + K_N)}{N} = 0, \tag{3.12}$$

which gives the proof of our theorem. □

Theorem 3.2 is related to the concept of asymptotic density, which in the case of alternating Lüroth-type expansions is defined as follows. Let $\{b_n\}$ be an increasing sequence of positive integers and let $K(M)$ be the number of elements of the sequence $\{b_n\}$ for which $b_t \leq M$. If the limit of $K(M)/M$ as $M \rightarrow +\infty$ exists, then we may say that the sequence $\{b_n\}$ has asymptotic density. **Theorem 3.2** provides the criterion for the sequence $\{n/\alpha_n > c_n\}$ to have asymptotic density, for a.a. x . This means that applying **Theorem 3.2** with $\alpha_n = 1$, for all $n \geq 1$, then with $\alpha_n = 2$, for all $n \geq 1$ we get successively the densities of $\{n \mid \alpha_n = 1\}$, $\{n \mid \alpha_n = 2\}$, and so on. Note that if $c_n \rightarrow +\infty$, then by **Theorem 3.2**, c exists and equals zero.

Using **Proposition 2.1** and Borel-Cantelli lemma, we can have a sharper result about the behaviour of the sequence $\{\alpha_n\}$ according to the following theorem.

THEOREM 3.3. *Let $\{c_n\}_n, \{d_n\}_n$, with $0 < c_n \leq d_n$, be two sequences of real numbers which tend to $+\infty$. Moreover, assume that*

$$\limsup \frac{(c_n + 1)}{d_n} = u < 1 \tag{3.13}$$

and that

$$\sum_{n=1}^{+\infty} \frac{1}{d_n(c_n + 1)} = +\infty. \tag{3.14}$$

Then for a.a. x , the inequalities $d_n < \alpha_n \leq d_n(1 + 1/(c_n + 1))$ hold for many infinite values of n .

PROOF. It is known that

$$\lambda(u < \alpha_n \leq w) = \lambda(u < \alpha_n) - \lambda(w < \alpha_n). \tag{3.15}$$

Then from (3.2) we have

$$\begin{aligned} \lambda\left(d_n < \alpha_n \leq d_n + \frac{d_n}{c_n + 1}\right) &= \frac{1}{[d_n] + 1} - \frac{1}{[d_n + d_n/(c_n + 1)] + 1} \\ &\geq \frac{1}{d_n + 1} - \frac{1}{d_n + d_n/(c_n + 1)} \\ &\geq \frac{1 - (c_n + 1)/d_n}{3d_n(c_n + 1)} \geq \frac{a}{d_n(c_n + 1)}, \end{aligned} \tag{3.16}$$

where a is a suitable constant, which in view of (3.13), for n sufficiently large, can be chosen arbitrarily close to $1 - u$, hence $a > 0$. Now by using Proposition 2.1, the Borel-Cantelli lemma is applicable. So (3.14) and (3.16) imply the statement of Theorem 3.3, and the proof is complete. \square

Theorem 3.3 states that if for a sequence d_n tending to $+\infty$, there is a sequence c_n such that (3.13) and (3.14) hold, then for a.a. x , infinitely often $\alpha_n \sim d_n$.

This raises the problem whether $M_n = \max(\alpha_1, \alpha_2, \dots, \alpha_n)$ follows an asymptotic law. We will deal with this problem in the next section.

4. The distribution of the maximum of the first n digits. If M_n is the maximum of the first n digits, then we take the following theorem.

THEOREM 4.1. For any fixed $y > 0$,

$$\lim_{n \rightarrow \infty} \lambda\left(\frac{M_n}{n} \leq y\right) = \exp\left(-\frac{1}{y+1}\right). \tag{4.1}$$

PROOF. We define the events $A_i = \{x \mid \alpha_i/n \leq y\}$, $1 \leq i \leq n$. It is obvious that

$$\left\{x \mid \frac{M_n}{n} \leq y\right\} = \bigcap_{i=1}^n A_i, \tag{4.2}$$

and therefore by using Proposition 2.1 and (3.2), we have

$$\lambda\left(\frac{M_n}{n} \leq y\right) = \prod_{i=1}^n \lambda(A_i) = \prod_{i=1}^n \lambda\left(\frac{\alpha_i}{n} \leq y\right) = \prod_{i=1}^n \left[1 - \lambda\left(\frac{\alpha_i}{n} > y\right)\right]. \tag{4.3}$$

But

$$\lambda\left(\frac{\alpha_i}{n} > y\right) = \lambda(\alpha_i > ny) = \sum_{k=[ny]+1}^{+\infty} \frac{1}{k(k+1)} = \frac{1}{[ny]+1}. \tag{4.4}$$

So

$$\lambda\left(\frac{M_n}{n} \leq y\right) = \prod_{i=1}^n \left[1 - \frac{1}{[ny]+1}\right] = \left(1 - \frac{1}{[ny]+1}\right)^n. \tag{4.5}$$

Using a well-known characteristic limit relation, the proof is complete. \square

Hence we can obtain the following corollary.

COROLLARY 4.2. Let K_n be a random variable defined on the probability space (I, B_I, λ) , and assume that K_n converges in probability to 1. Then

$$\lim_{n \rightarrow +\infty} \lambda\left(\frac{K_n}{M_n} < y\right) = 1 - e^{-y/(y+1)}. \tag{4.6}$$

PROOF. We may write

$$\frac{K_n}{M_n} = \frac{K_n}{n} \cdot \frac{n}{M_n} = \frac{n}{M_n} + \frac{n}{M_n} \left(\frac{K_n}{n} - 1\right). \tag{4.7}$$

Using (4.7) we have only to show that the second term tends in probability to 0, since then using a well-known result of Cramer we may obtain the statement of [Corollary 4.2](#).

More precisely we have to show that, for any positive real number u ,

$$\lim_{n \rightarrow +\infty} \lambda\left(\left|\frac{n}{M_n} \cdot \left(\frac{K_n}{n} - 1\right)\right| \geq u\right) = 0. \quad (4.8)$$

If we apply this relation, for any fixed U , then we obtain

$$\lambda\left(\left|\frac{n}{M_n} \cdot \left(\frac{K_n}{n} - 1\right)\right| \geq u\right) \leq \lambda\left(\frac{n}{M_n} > U\right) + \lambda\left(\left|\frac{K_n}{n} - 1\right| \geq \frac{u}{U}\right). \quad (4.9)$$

From (4.9) we get that the first and the second terms are smaller than any prescribed real number by using [Theorem 4.1](#) and the assumption on K_n , respectively. So the proof is complete. \square

From the occurrence of the exponential distribution in [Theorem 4.1](#) and [Corollary 4.2](#) it can be derived that the number of terms of the sequence $\{\alpha_n\}$, which are of the same order as M_n , follows a Poisson distribution. This is given in the following theorem.

THEOREM 4.3. *Let Y_n denote the number of terms in the sequence $\{\alpha_n\}$ for which $\alpha_n > \gamma$. Then its asymptotic probability function is given by*

$$\lim_{n \rightarrow +\infty} \lambda(Y_n = K) = \frac{e^{-1(\gamma+1)}}{K!(\gamma+1)^K}. \quad (4.10)$$

PROOF. Using [Proposition 2.1](#) and relation (3.2) we have that the random variables Z_N follow a binomial distribution with parameters n and $1/([n\gamma] + 1)$. Since $n/([n\gamma] + 1) \rightarrow 1/(\gamma + 1)$, as $n \rightarrow \infty$, by the result known as ‘‘Poisson approximation to the binomial distribution’’ (see [2]), we obtain that the distribution of Y_n is Poisson with parameter $1/(\gamma + 1)$. So the proof is complete. \square

REFERENCES

- [1] Y. S. Chow and H. Robbins, *On sums of independent random variables with infinite moments and ‘‘fair’’ games*, Proc. Nat. Acad. Sci. U.S.A. **47** (1961), 330–335. [MR 23#A2908](#). [Zbl 099.35103](#).
- [2] W. Feller, *An Introduction to Probability Theory and Its Applications. Vol. I*, 3rd ed., John Wiley & Sons, New York, 1968. [MR 37#3604](#). [Zbl 155.23101](#).
- [3] H. Jager and C. de Vroedt, *Lüroth series and their ergodic properties*, Nederl. Akad. Wetensch. Proc. Ser. A **31** (1969), 31–42. [MR 39#157](#). [Zbl 167.32201](#).
- [4] S. Kalpazidou and C. Ganatsiou, *Knopfmacher expansions in number theory*, to appear in *Quaestiones Math.*
- [5] S. Kalpazidou, A. Knopfmacher, and J. Knopfmacher, *Lüroth-type alternating series representations for real numbers*, Acta Arith. **55** (1990), no. 4, 311–322. [MR 91i:11011](#). [Zbl 702.11048](#).
- [6] A. Knopfmacher and J. Knopfmacher, *Two constructions of the real numbers via alternating series*, Int. J. Math. Math. Sci. **12** (1989), no. 3, 603–613. [MR 90k:26003b](#). [Zbl 683.10008](#).
- [7] ———, *New series and product representations for real numbers*, Manuscript, 1990.

- [8] A. Oppenheim, *The representation of real numbers by infinite series of rationals*, Acta Arith. **21** (1972), 391–398. [MR 46#8982](#). [Zbl 258.10003](#).
- [9] T. Šalát, *Zur Metrischen theorie der Lürothschen Entwicklungen der reellen Zahlen*, Czechoslovak Math. J. **18** (93) (1968), 489–522 (German). [MR 37#5179](#). [Zbl 162.34703](#).

C. GANATSIOU: TECHNOLOGICAL EDUCATIONAL INSTITUTE OF LARISSA, SCHOOL OF TECHNOLOGICAL APPLICATIONS, DEPARTMENT OF EXACT SCIENCES, SECTION OF MATHEMATICS, LARISSA, GREECE

E-mail address: ganatsio@teilar.gr

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mssklai@cityu.edu.hk