

A NOTE ON HOMOMORPHISMS OF HILBERT ALGEBRAS

SERGIO CELANI

Received 21 November 2000 and in revised form 14 May 2001

We give a representation theorem for Hilbert algebras by means of ordered sets and characterize the homomorphisms of Hilbert algebras in terms of applications defined between the sets of all irreducible deductive systems of the associated algebras. For this purpose we introduce the notion of order-ideal in a Hilbert algebra and we prove a separation theorem. We also define the concept of semi-homomorphism as a generalization of the similar notion of Boolean algebras and we study its relation with the homomorphism and with the deductive systems.

2000 Mathematics Subject Classification: 08A30.

1. Introduction and preliminaries. Hilbert algebras correspond to the algebraic counterpart of the implicative fragment of Intuitionistic Propositional Logic. For the general development of Hilbert algebras, the notion of deductive system (also called implicative filters) plays an important role. For example, it is known that the set of all deductive systems of a Hilbert algebra is a distributive lattice isomorphic to the lattice of congruences (see [2, 4]). In [4], Diego proved that every Hilbert algebra can be represented as a subalgebra of the Hilbert algebra of the open subsets of a suitable T_0 topological space. Our central aim is to prove a new representation theorem for Hilbert algebras by means of ordered sets and give some results relating homomorphisms of Hilbert algebras to deductive systems and a weaker version of homomorphisms.

In Section 2, we start recalling the notion of deductive system and irreducible deductive system. We prove a theorem of separation, and a representation theorem for Hilbert algebras by means of ordered sets. In Section 3, we introduce the notion of semi-homomorphism between Hilbert algebras. We prove a characterization of the semi-homomorphism by means of deductive systems and we prove a characterization of the homomorphism by means of irreducible deductive systems. To conclude, we deal with the study of injective and surjective homomorphism of Hilbert algebras.

We note that Hilbert algebras are dual to positive implicative BCK-algebras. This duality was first noted in [3], and fully proved in [1, 5]. Thus, the results given in this note can be applied straightly to the positive implicative BCK-algebras.

DEFINITION 1.1. A *Hilbert algebra* is an algebra $\langle A, \rightarrow, 1 \rangle$ where A is a nonempty set, \rightarrow is a binary operation on A , 1 is an element of A such that the following conditions are satisfied for every $a, b, c \in A$:

(H1) $a \rightarrow (b \rightarrow a) = 1$.

(H2) $(a \rightarrow (b \rightarrow c)) \rightarrow ((a \rightarrow b) \rightarrow (a \rightarrow c)) = 1$.

(H3) If $a \rightarrow b = 1$ and $b \rightarrow a = 1$, then $a = b$.

Let A be a Hilbert algebra. It is known that the relation \leq defined in A by $a \leq b$ if and only if $a \rightarrow b = 1$ is a partial order on A where 1 is the greatest element of A with respect to this ordering.

PROPOSITION 1.2. *Let A be a Hilbert algebra. Then the following properties are valid for any $a, b, c \in A$:*

- (1) $a \leq b \rightarrow a$.
- (2) $a \rightarrow 1 = 1$.
- (3) $1 \rightarrow a = a$.
- (4) $a \rightarrow (b \rightarrow c) = b \rightarrow (a \rightarrow c)$.
- (5) $a \rightarrow (b \rightarrow c) = (a \rightarrow b) \rightarrow (a \rightarrow c)$.
- (6) If $a \leq b$, then $c \rightarrow a \leq c \rightarrow b$ and $b \rightarrow c \leq a \rightarrow c$.

PROOF. See [2] or [4]. □

2. Representation theorem

DEFINITION 2.1. Let A be a Hilbert algebra. A *deductive system* of A is a subset D of A such that

- (D1) $1 \in D$,
- (D2) if $a, a \rightarrow b \in D$, then $b \in D$.

Let A be a Hilbert algebra. The set of all deductive systems of A is denoted by $Ds(A)$. It is known that $Ds(A)$ is a distributive lattice (see [4]). If $X \subseteq A$, then the deductive system generated by X , in symbols $D(X)$, is the set (see [2] or [4])

$$D(X) = \{a \in A : \exists \{x_0, x_1, \dots, x_n\} \subseteq X : x_0 \rightarrow (x_1 \rightarrow \dots (x_n \rightarrow a) \dots) = 1\}. \quad (2.1)$$

DEFINITION 2.2. Let A be a Hilbert algebra. Let $D \in Ds(A)$. Then we say that

- (1) D is *irreducible* if and only if for any $D_1, D_2 \in Ds(A)$ such that $D = D_1 \cap D_2$, then $D = D_1$ or $D = D_2$.
- (2) D is *completely irreducible* if and only if for any family $H = \{D_i : i \in I\} \subseteq Ds(A)$ such that $D = \bigcap \{D_i : i \in I\}$, then $D = D_i$ for some $i \in I$.

We denote the set of all irreducible and the set of all completely irreducible deductive systems of A , by $X(A)$ and $X_c(A)$, respectively.

The following results are known (see [2, 4]).

THEOREM 2.3. *Let A be a Hilbert algebra. Then*

- (1) $D \in X(A)$ if and only if for every $a, b \notin D$ there exists $c \notin D$ such that $a \leq c$ and $b \leq c$.
- (2) $D \in X_c(A)$ if and only if there exists $c \notin D$ such that $a \rightarrow c \in D$ for every $a \notin D$.
- (3) For all $a, b \in A$ if $a \not\leq b$ there exists $D \in X_c(A)$ such that $a \in D$ and $b \notin D$.

The following result gives another characterization of irreducible deductive systems in a Hilbert algebra.

LEMMA 2.4. *Let A be a Hilbert algebra. Let $D \in Ds(A)$. Then D is irreducible if and only if for every $a, b \notin D$ there exists $c \notin D$ such that $a \rightarrow c, b \rightarrow c \in D$.*

PROOF. The direction (\Rightarrow) follows by assertion 1 of [Theorem 2.3](#).

(\Leftarrow) . Suppose that $D_1, D_2 \in \text{Ds}(A)$ such that $D = D_1 \cap D_2$, $D \neq D_1$ and $D \neq D_2$. Then there exist $a \in D_1 - D$ and $b \in D_2 - D$. Then, by assumption, there exists $c \notin D$ such that $a \rightarrow c \in D$ and $b \rightarrow c \in D$. So, $a, a \rightarrow c \in D_1$ and as $D_1 \in \text{Ds}(A)$, then $c \in D_1$. Similarly, $c \in D_2$. Thus, $c \in D_1 \cap D_2 = D$, which is a contradiction. \square

DEFINITION 2.5. Let A be a Hilbert algebra. A subset I of A is called an *order-ideal* of A if

(ID1) If $b \in I$ and $a \leq b$, then $a \in I$.

(ID2) If $a, b \in I$ then there exists $c \in I$ such that $a \leq c$ and $b \leq c$.

The set of all order-ideals of A will be denoted by $\text{Id}(A)$.

Now, we prove the mentioned separation theorem.

THEOREM 2.6. Let A be a Hilbert algebra. Let $D \in \text{Ds}(A)$ and $I \in \text{Id}(A)$ such that $D \cap I = \emptyset$. Then there exists $P \in X(A)$ such that $D \subseteq P$ and $P \cap I = \emptyset$.

PROOF. Consider the following subset of $\text{Ds}(A)$:

$$\mathcal{F} = \{H \in \text{Ds}(A) : D \subseteq H \text{ and } H \cap I = \emptyset\}. \quad (2.2)$$

Since $D \in \mathcal{F}$, then $\mathcal{F} \neq \emptyset$. It is clear that the union of a chain of elements of \mathcal{F} is also in \mathcal{F} . So, by Zorn's lemma, there exists a deductive systems P maximal in \mathcal{F} . We prove that $P \in X(A)$. Let $a, b \notin P$ and consider the deductive systems $P_a = D(P \cup \{a\})$ and $P_b = D(P \cup \{b\})$. Clearly, $P \subset P_a \cap P_b$. Then, $P_a, P_b \notin \mathcal{F}$. Thus, $P_a \cap I \neq \emptyset$ and $P_b \cap I \neq \emptyset$. It follows that there exist $x, y \in I$ such that $a \rightarrow x \in P$ and $b \rightarrow y \in P$. Since I is an order-ideal, there exists $c \in I$ such that $x \leq c$ and $y \leq c$. So, $a \rightarrow x \leq a \rightarrow c \in P$ and $b \rightarrow y \leq b \rightarrow c \in P$. Therefore, by [Lemma 2.4](#), we conclude that $P \in X(A)$. \square

Let $\langle X, \leq \rangle$ be a poset. For $Y \subseteq X$, we put $[Y] = \{x \in X : y \leq x \text{ for some } y \in Y\}$. A subset Y of X is *increasing* if $Y = [Y]$. The sets of all increasing subsets of X will be denoted by $\mathcal{P}_i(X)$.

LEMMA 2.7. Let $\langle X, \leq \rangle$ be a poset. Then $\langle \mathcal{P}_i(X), \Rightarrow, X \rangle$ is a Hilbert algebra where the implication \Rightarrow is defined by

$$U \Rightarrow V = \{x \in X : [x] \cap U \subseteq V\}, \quad (2.3)$$

for $U, V \in \mathcal{P}_i(X)$.

PROOF. The proof is immediate. \square

Let A be a Hilbert algebra. Consider the poset $\langle X(A), \subseteq \rangle$ and consider the mapping

$$\beta : A \rightarrow \mathcal{P}_i(X(A)) \quad (2.4)$$

defined by

$$\beta(a) = \{P \in X(A) : a \in P\}. \quad (2.5)$$

THEOREM 2.8 (representation theorem). Let A be a Hilbert algebra. Then, A is isomorphic to the subalgebra $\beta(A) = \{\beta(a) : a \in A\}$ of $\mathcal{P}_i(X(A))$.

PROOF. It is clear that the map $\beta : A \rightarrow \mathcal{P}_i(X(A))$ is injective. We prove that $\beta(a \rightarrow b) = \beta(a) \Rightarrow \beta(b)$. Let $P, Q \in X(A)$ and $a, b \in A$ such that $a \rightarrow b \in P$. If $Q \in [P] \cap \beta(a)$, then $Q \in \beta(b)$, that is, $b \in Q$.

Suppose that $a \rightarrow b \notin P$. Consider the deductive systems $P_a = D(P \cup \{a\})$. Since $b \notin P_a$, then there exists $Q \in X(A)$ such that $P \subseteq Q$, $a \in Q$ and $b \notin Q$. Thus, $P \notin \beta(a) \Rightarrow \beta(b)$. \square

3. Homomorphisms of Hilbert algebras. The purpose of this section is to study the homomorphism between Hilbert algebras.

DEFINITION 3.1. Let A and B be Hilbert algebras. A *semi-homomorphism* is a mapping $h : A \rightarrow B$ such that

$$(SH1) \quad h(1) = 1,$$

$$(SH2) \quad h(a \rightarrow b) \leq h(a) \rightarrow h(b), \text{ for every } a, b \in A.$$

THEOREM 3.2. Let A and B be Hilbert algebras. Let $h : A \rightarrow B$ be a mapping. Then the following conditions are equivalent:

(1) h is a semi-homomorphism

(2) $h^{-1}(D) \in \text{Ds}(A)$ for all $D \in \text{Ds}(B)$.

PROOF. (1) \Rightarrow (2). Let $D \in \text{Ds}(B)$. By condition (SH1), $1 \in h^{-1}(D)$. If $a, a \rightarrow b \in h^{-1}(D)$, then $h(a), h(a \rightarrow b) \leq h(a) \rightarrow h(b) \in D$. Since D is a deductive system, then $h(b) \in D$, that is, $b \in h^{-1}(D)$. Thus, $h^{-1}(D) \in \text{Ds}(A)$.

(2) \Rightarrow (1). Suppose that $h^{-1}(D) \in \text{Ds}(A)$ for all $D \in \text{Ds}(B)$. If $h(1) \neq 1$, then there exists $D \in \text{Ds}(B)$ such that $h(1) \notin D$, that is, $1 \notin h^{-1}(D)$, which is impossible. Thus, $h(1) = 1$. Suppose that $h(a \rightarrow b) \not\leq h(a) \rightarrow h(b)$. Then, there exists $P \in X(B)$ such that $h(a \rightarrow b) \in P$ and $h(a) \rightarrow h(b) \notin P$. Consider the deductive system $D(P \cup h(a))$. Since $h(b) \notin D(P \cup h(a))$, there exists $Q \in X(A)$ such that $P \subseteq Q$, $h(a) \in Q$ and $h(b) \notin Q$. As $a \rightarrow b \in h^{-1}(P) \subseteq h^{-1}(Q)$ and $a \in h^{-1}(Q)$, then $b \in h^{-1}(Q)$, which is a contradiction. Therefore, $h(a \rightarrow b) \leq h(a) \rightarrow h(b)$. \square

Let A be a Hilbert algebra. The set complement of a subset P of A is denoted by P^c .

Let A and B be Hilbert algebras. Let $h : A \rightarrow B$ be a homomorphism. It is easy to see that, for each $P \in X(A)$, the subset of B

$$(h(P^c)) = \{y \in B : y \leq h(p) \text{ for some } p \notin P\} \quad (3.1)$$

is an order-ideal of B . This fact will be used in the following results.

THEOREM 3.3. Let A and B be Hilbert algebras. Let $h : A \rightarrow B$ be a semi-homomorphism. Then the following conditions are equivalent:

(1) h is a homomorphism.

(2) For every $(P, Q) \in X(B) \times X(A)$ such that $h^{-1}(P) \subseteq Q$ there exists $D \in X(B)$ such that $P \subseteq D$ and $h^{-1}(D) = Q$.

PROOF. (1) \Rightarrow (2). Suppose that $h : A \rightarrow B$ is a homomorphism. Let $(P, Q) \in X(B) \times X(A)$ such that $h^{-1}(P) \subseteq Q$. Consider the deductive system $D(P \cup h(Q))$. Since $Q \in X(A)$ then $(h(Q^c)) \in \text{Id}(B)$. We prove that

$$D(P \cup h(Q)) \cap (h(Q^c)) = \emptyset. \quad (3.2)$$

Suppose the contrary. Then there exists $p \in P, q \in Q$, and $a \notin Q$ such that $p \rightarrow (h(q) \rightarrow h(a)) = 1$. Since h is a homomorphism, then $p \rightarrow h(q \rightarrow a) = 1 \in P$. So, $h(q \rightarrow a) \in P$ and this implies that $q \rightarrow a \in h^{-1}(P) \subseteq Q$. Thus, $a \in Q$, which is a contradiction. Therefore, $D(P \cup h(Q)) \cap (h(Q^c)) = \emptyset$. By [Theorem 2.6](#), there exists $D \in X(B)$ such that $P \subseteq D$ and $h^{-1}(D) = Q$.

(2) \Rightarrow (1). Let $a, b \in A$ and assume that $h(a) \rightarrow h(b) \neq h(a \rightarrow b)$. Then there exists $P \in X(B)$ such that $h(a) \rightarrow h(b) \in P$ and $a \rightarrow b \notin h^{-1}(P)$. Since h is a semi-homomorphism, by [Theorem 3.2](#), $h^{-1}(P) \in \text{Ds}(A)$. So, there exists $Q \in X(A)$ such that $h^{-1}(P) \subseteq Q$, $a \in Q$ and $b \notin Q$. By assumption, there exists $D \in X(B)$ such that $P \subseteq D$ and $h^{-1}(D) = Q$. But since $h(a) \rightarrow h(b) \in P \subseteq D$ and $h(a) \in D$, we get that $h(b) \in D$, which is a contradiction. We conclude that h is a homomorphism. \square

Let A and B be Hilbert algebras and let $h : A \rightarrow B$ be a homomorphism. Consider the subset of $X(B)$

$$X_h(B) = \{P \in X(B) : h^{-1}(P) \in X(A)\}. \tag{3.3}$$

Consider the partial mapping

$$X(h) : X_h(B) \longrightarrow X(A) \tag{3.4}$$

defined by

$$X(h)(P) = h^{-1}(P), \tag{3.5}$$

with $P \in X_h(B)$.

THEOREM 3.4. *Let A and B be Hilbert algebras and let $h : A \rightarrow B$ be a homomorphism. Then*

- (1) h is injective if and only if $X(h)$ is surjective.
- (2) h is surjective if and only if $X(h)$ is injective with $X_h(B) = X(B)$.

PROOF. (1) \Rightarrow Let $P \in X(A)$. Consider the deductive system $D(h(P))$. We prove that

$$D(h(P)) \cap (h(P^c)) = \emptyset. \tag{3.6}$$

Suppose the contrary case. Then there exists $p \in P$ and $a \notin P$ such that $h(p) \rightarrow h(a) = 1$. Since h is homomorphism, $h(p \rightarrow a) = 1 = h(1)$, and as h is injective, we get $p \rightarrow a = 1$. So, $a \in P$, which is impossible. Thus, by [Theorem 2.6](#), there exists $Q \in X(B)$ such that $h(P) \subseteq Q$ and $h(P^c) \cap Q = \emptyset$, that is, $h^{-1}(Q) = P$. Therefore, $X(h)$ is surjective.

(\Leftarrow) Let $a, b \in A$. Suppose that $a \neq b$. Then there exists $P \in X(A)$ such that $a \in P$ and $b \notin P$. By assumption, there exists $Q \in X_h(B) \subseteq X(B)$ such that $P = h^{-1}(Q)$. Then, $h(a) \in Q$ and $h(b) \notin Q$. It follows that $h(a) \neq h(b)$, which implies that h is injective.

(2) \Rightarrow First, we prove that $X_h(B) = X(B)$. Let $Q \in X(B)$. Since h is a homomorphism, $h^{-1}(Q) \in \text{Ds}(A)$. Let $a, b \in A$ such that $h(a), h(b) \notin Q$. Then there exists $c \notin Q$ such that $h(a) \leq c$ and $h(b) \leq c$. Since h is onto, there exists $d \in B$ such that $c = h(d)$. So, $h(a) \leq h(d)$ and $h(b) \leq h(d)$ with $d \notin h^{-1}(Q)$. Thus, $h^{-1}(Q) \in X(A)$ and consequently $X_h(B) = X(B)$.

Let $P, Q \in X(B)$ such that $h^{-1}(P) \subseteq h^{-1}(Q)$. Then it is easy to check that $P \subseteq Q$. So, $X(h)$ is an order-isomorphism. Thus, $X(h)$ is injective.

(\Leftarrow) Let $b \in B$ and assume that $b \notin h(A) = \{h(a) : a \in A\}$. Consider the deductive system $D(b)$ and the deductive system $D(h(h^{-1}(D(b))))$. We prove that

$$D(h(h^{-1}(D(b)))) \cap (b) = \emptyset. \quad (3.7)$$

Suppose the contrary. Then there exists $h(x) \in D(b)$ such that $h(x) \leq b$. It follows that $b = h(x)$, which is a contradiction. So, by [Theorem 2.6](#), there exists $P \in X(B)$ such that $h(h^{-1}(D(b))) \subseteq P$ and $b \notin P$. We prove now that

$$(P^c \cap h(A)) = \{y \in B : y \leq x \text{ for some } x \in P^c \cap h(A)\} \quad (3.8)$$

is an order-ideal of B . It is enough to prove that if $x, y \in P^c \cap h(A)$, then there exists $c \in P^c \cap h(A)$ such that $x \leq c$ and $y \leq c$. Let $x, y \in P^c \cap h(A)$. Then $x = h(a)$ and $y = h(b)$ for some $a, b \in A$. Since $a, b \notin h^{-1}(P)$ and by assumption, $h^{-1}(P) \in X(B)$, then there exists $c \notin h^{-1}(P)$ such that $a \leq c$ and $b \leq c$. So, $x \leq h(c)$ and $y \leq h(c)$. Thus, $(P^c \cap h(A))$ is an order-ideal of B .

We prove that

$$D(P \cap h(A) \cup \{b\}) \cap (P^c \cap h(A)) = \emptyset. \quad (3.9)$$

Assume the contrary, then there exists $a \in D(P \cap h(A) \cup \{b\}) \cap (P^c \cap h(A))$, $p = h(q) \in P \cap h(A)$, and $d = h(z) \in P^c \cap h(A)$ such that $b \rightarrow (h(q) \rightarrow a) = 1$ and $a \leq h(z)$. Thus, $h(q \rightarrow z) \in D(b)$. It follows that $q \rightarrow z \in h^{-1}(D(b)) \subseteq h^{-1}(P)$. Since $q \in P$, then $z \in h^{-1}(P)$, that is, $h(z) = d \in P$, which is absurd. Thus, (3.9) is valid. By [Theorem 2.6](#), there exists $Q \in X(B)$ such that $Q \cap h(A) = P \cap h(A)$ and $b \in Q$. So, $h^{-1}(Q) = h^{-1}(P)$ and as $X(h)$ is injective we conclude that $Q = P$. Thus, $b \in P$, which is a contradiction. So $b \in h(A)$ and therefore h is surjective. \square

4. Conclusions. In this note, we give a representation theorem for Hilbert algebras and we give several results relating homomorphism of Hilbert algebras to deductive systems and semi-homomorphisms. One of the referees pointed out that these results can help to find a full duality for Hilbert algebras. This problem was one of the motivations to study the representation for Hilbert algebras. Unfortunately, we cannot give a full duality. However, using the results of this note, we have given a duality for implicative semilattice (or positive implicative BCK-algebras with condition (S) [6]). These results will be presented in a future paper.

ACKNOWLEDGEMENT. I would like to thank the referees for the comments and suggestions on the presentation of this note.

REFERENCES

- [1] C. Bărbăcioru, *Positive implicative BCK-algebras*, Math. Japon. **38** (1993), no. 3, 513-520.
- [2] D. Buşneag, *On the maximal deductive systems of a bounded Hilbert algebra*, Bull. Math. Soc. Sci. Math. R. S. Roumanie (N.S.) **31(79)** (1987), no. 1, 9-21.
- [3] W. H. Cornish, *On positive implicative BCK-algebras*, Math. Sem. Notes Kobe Univ. **8** (1980), no. 3, 455-468.

- [4] A. Diego, *Sur les algèbres de Hilbert*, Collection de Logique Mathématique, Sér. A, Fasc. vol. 21, Gauthier-Villars, Paris, 1966 (French), translated from Spanish by Luisa Iturrioz.
- [5] M. Kondo, *Hilbert algebras are dual isomorphic to positive implicative BCK-algebras*, Math. Japon. **49** (1999), no. 2, 265-268.
- [6] J. Meng, Y. B. Jun, and S. M. Hong, *Implicative semilattices are equivalent to positive implicative BCK-algebras with condition (S)*, Math. Japon. **48** (1998), no. 2, 251-255.

SERGIO CELANI: DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS, UNIVERSIDAD NACIONAL DEL CENTRO, PINTO 399, 7000 TANDIL, ARGENTINA

E-mail address: scelani@exa.unicen.edu.ar

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

| | |
|------------------------|------------------|
| Manuscript Due | December 1, 2008 |
| First Round of Reviews | March 1, 2009 |
| Publication Date | June 1, 2009 |

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru