

## SCHRÖDINGER OPERATORS WITH A SINGULAR POTENTIAL

TOKA DIAGANA

Received 13 April 2001

This note is devoted to the study of some Schrödinger operators with a singular real potential  $Q$ . The potential  $Q$  is chosen so that the algebraic sum  $L = -\Delta + Q$  is not defined. Next, we define the sum form operator which will be well defined and we show that this operator verifies the well-known Kato's square root problem.

2000 Mathematics Subject Classification: 35J05, 47B25, 47B44.

**1. Introduction.** Let  $H$  be a Hilbert space and let  $A$  and  $B$  be two unbounded linear operators on  $H$ . The algebraic sum of  $A$  and  $B$  is defined by

$$D(A+B) = D(A) \cap D(B), \quad (A+B)u = Au + Bu, \quad \forall u \in D(A) \cap D(B). \quad (1.1)$$

Thus, if  $A$  and  $B$  are chosen so that  $D(A) \cap D(B)$  is reduced to  $\{0\}$ , therefore the algebraic sum of  $A$  and  $B$  is not defined except in zero. So, the algebraic sum is not always well adapted to problems arising in mathematical analysis. This has motivated some mathematical developments where the notion of sum is weakened or relaxed, see for example [2, 3, 4]. Consider  $A$  and  $B$  be two linear operators given on  $L^2(\mathbb{R})$  by

$$\begin{aligned} D(A) &= H^2(\mathbb{R}^n), \quad Au = -\Delta u, \\ D(B) &= \{u \in L^2(\mathbb{R}^n) : Qu \in L^2(\mathbb{R}^n)\}, \quad Bu = Qu, \end{aligned} \quad (1.2)$$

where  $Q$  is a measurable real function. Let  $L$  be the operator of Schrödinger given by  $L = -\Delta + Q$ . Let  $\Phi$  and  $\Psi$  be the closed and densely defined sesquilinear forms associated, respectively, to  $-\Delta$  and  $Q$ , given by

$$\begin{aligned} \Phi(u, v) &= \int_{\mathbb{R}^n} \nabla u \bar{\nabla} v dx, \quad \forall u, v \in H^1(\mathbb{R}^n), \\ \Psi(u, v) &= \int_{\mathbb{R}^n} Qu \bar{v} dx, \quad \forall u, v \in D(B^{1/2}). \end{aligned} \quad (1.3)$$

We will put,  $\Upsilon(u, v) = (\Phi + \Psi)(u, v)$  for all  $u, v \in D(\Upsilon) = D(\Phi) \cap D(\Psi)$ , the sum of sesquilinear forms associated to  $-\Delta$  and  $Q$ .

### 2. The main result

**HYPOTHESIS ON  $Q$ .** We assume that the potential  $Q$  verifies  $H_Q$ , given by

$$Q > 0, \quad Q \in L^1(\mathbb{R}^n), \quad Q \notin L^2_{\text{loc}}(\mathbb{R}^n). \quad (2.1)$$

**PROPOSITION 2.1.** *Under hypothesis  $H_Q$  and if  $n < 4$ , then,*

$$D(A) \cap D(B) = \{0\}. \quad (2.2)$$

**PROOF.** Let  $u \in D(-\Delta) \cap D(Q)$  and suppose that  $u \neq 0$ . Since  $u \in H^2(\mathbb{R}^n)$  where  $n < 4$ , then  $u$  is a continuous function by Sobolev's theorem (see [1]). There exists an open subset  $\Omega$  of  $\mathbb{R}^n$  and there exists  $\delta > 0$  such that  $|u(x)| > \delta$  for all  $x \in \Omega$ . Let  $\Omega'$  be a compact subset of  $\Omega$ , equipped with the induced topology by  $\Omega$ . Thus  $\Omega'$  is also a compact subset of  $\mathbb{R}^n$ . It follows that

$$|Q|_{\Omega'} = \frac{(|Qu|)_{\Omega'}}{|u|_{\Omega'}} \in L^2(\Omega'), \quad (2.3)$$

because  $(|Qu|)_{\Omega'} \in L^2(\Omega')$  and  $1/(|u|)'_{\Omega'} \in L^\infty(\Omega')$ . Therefore  $Q \in L^2(\Omega')$ , that is impossible according to hypothesis  $H_Q$ , then  $u \equiv 0$ .  $\square$

**QUESTION 2.2.** Find a characterization of  $D(-\Delta) \cap D(Q)$ , when  $n \geq 4$ ?

**PROPOSITION 2.3.** *Under hypothesis  $H_Q$ , then*

- $D(A^{1/2}) \hookrightarrow D(B^{1/2})$  if  $n = 1$ ,
- $D(A^{1/2}) \cap D(B^{1/2}) \supseteq C_0^\infty(\mathbb{R}^n)$  if  $n > 1$ .

**EXAMPLE OF POTENTIAL  $Q$  VERIFYING  $H_Q$ .** Let  $\Omega$  be a compact subset of  $\mathbb{R}^n$  and let  $G$  be a complex function satisfying,  $\operatorname{Re} G > 0$ ,  $G \in L^1(\Omega)$ ,  $G \notin L^2(\Omega)$ , and  $G \equiv 0$  on  $\mathbb{R}^n - \Omega$ . Consider the following rational sequence  $\alpha_k = (\alpha_k^1, \alpha_k^2, \dots, \alpha_k^n) \in \mathbb{Q}^n$ . Then the function  $Q$ , given by

$$Q(x) = \sum_{k=1}^{+\infty} \frac{G(x - \alpha_k)}{k^2}, \quad (2.4)$$

verifies hypothesis  $H_Q$ .

**3. Generalized sum of  $A$  and  $B$ .** Consider the closed sesquilinear forms given by

$$\begin{aligned} \Phi(u, v) &= \int_{\mathbb{R}^n} \nabla u \overline{\nabla v} dx \quad \forall u, v \in H^1(\mathbb{R}^n), \\ \Psi(u, v) &= \int_{\mathbb{R}^n} Qu \bar{v} dx \quad \forall u, v \in D(B^{1/2}), \end{aligned} \quad (3.1)$$

and the sum of the forms  $\Phi$  and  $\Psi$  given by,  $\Upsilon = \Phi + \Psi$ , in other words,  $\Upsilon(u, v) = \int_{\mathbb{R}^n} (\nabla u \overline{\nabla v} + Qu \bar{v}) dx$  for all  $u, v \in D(A^{1/2}) \cap D(B^{1/2})$ . The sesquilinear form  $\Upsilon$  is a closed sectorial and densely defined form as sum of closed sectorial and densely defined forms, then there exists a unique  $m$ -sectorial operator  $A \oplus B$ , called sum form or generalized form of  $A$  and  $B$  associated to  $\Upsilon$  (see [3, 4]) and  $\Upsilon$  has the following representation:

$$\Upsilon(u, v) = \langle (A \oplus B)u, v \rangle \quad \forall u \in D(A \oplus B), v \in D(A^{1/2}) \cap D(B^{1/2}). \quad (3.2)$$

According to [3], the operator  $A \oplus B$  verifies the well-known condition of Kato (see [4]), in other words,

$$D((A \oplus B)^{1/2}) = D(\Upsilon) = D\left((A \oplus B)^*{}^{1/2}\right), \quad (3.3)$$

and the operator  $A \oplus B$  is given by

$$D(A \oplus B) = \{u \in H^1(\mathbb{R}^n) : Q|u|^2 \in L^1(\mathbb{R}^n), -\Delta u + Qu \in L^2(\mathbb{R}^n)\}, \quad (A \oplus B)u = -\Delta u + Qu. \quad (3.4)$$

**ACKNOWLEDGEMENT.** I would like to thank Prof Chandler Davis for his help.

#### REFERENCES

- [1] R. A. Adams, *Sobolev Spaces*, Pure and Applied Mathematics, vol. 65, Academic Press, New York, 1975.
- [2] P. R. Chernoff, *Product Formulas, Nonlinear Semigroups, and Addition of Unbounded Operators*, Memoirs of the American Mathematical Society, no. 140, American Mathematical Society, Rhode Island, 1974.
- [3] T. Diagana, *Sommes d'opérateurs et conjecture de Kato-McIntosh [Sums of linear operators and the Kato-McIntosh conjecture]*, C. R. Acad. Sci. Paris Sér. I Math. **330** (2000), no. 6, 461–464 (French).
- [4] T. Kato, *Perturbation Theory for Linear Operators*, Die Grundlehren der mathematischen Wissenschaften, vol. 132, Springer-Verlag, New York, 1966.

TOKA DIAGANA: DEPARTMENT OF MATHEMATICS, HOWARD UNIVERSITY, 2441 SIXTH STREET, NW, WASHINGTON, DC 20059, USA

*E-mail address:* [tdiagana@howard.edu](mailto:tdiagana@howard.edu)

## Special Issue on Intelligent Computational Methods for Financial Engineering

### Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Lean Yu**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [yulean@amss.ac.cn](mailto:yulean@amss.ac.cn)

**Shouyang Wang**, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; [sywang@amss.ac.cn](mailto:sywang@amss.ac.cn)

**K. K. Lai**, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; [mskklai@cityu.edu.hk](mailto:mskklai@cityu.edu.hk)