

THE CAYLEY TRANSFORM OF BANACH ALGEBRAS

ZHIDONG PAN

Received 30 July 2001

The main result of Haynes (1991) is that a square matrix is convergent ($\lim_{n \rightarrow \infty} D^n = 0$) if and only if it is the Cayley transform $C_A = (I - A)^{-1}(I + A)$ of a stable matrix A . In this note, we show, with a simple proof, that the above is true in a much more general setting of complex Banach algebras.

2000 Mathematics Subject Classification: 15A04, 15A24, 47A60, 47C05.

A square matrix D is called convergent if $\lim_{n \rightarrow \infty} D^n = 0$, or equivalently, $\lim_{n \rightarrow \infty} \|D^n\| = 0$, where $\|\cdot\|$ denotes the matrix norm. A square matrix is called stable if its characteristic values all have negative real parts. The above definitions can be generalized to elements in general complex Banach algebras as follows: let \mathcal{A} be a complex Banach algebra with norm $\|\cdot\|$, and let $\sigma(T)$ be the spectrum of T and $r(T)$ the spectral radius of T for any $T \in \mathcal{A}$. An element $D \in \mathcal{A}$ is called convergent if $\lim_{n \rightarrow \infty} \|D^n\| = 0$ and an element $A \in \mathcal{A}$ is called stable if $\sigma(A)$ is contained in the open left half of the complex plane. Inspired by the work of Stein and Taussky [3, 4] and Haynes [2] shows that a square matrix is convergent if and only if it is the Cayley transform of some stable matrix. In this note, we show that this holds for any complex Banach algebra and our proof is much simpler than that of [2].

To this end, we need the following elementary lemma.

LEMMA 1. *Let $D \in \mathcal{A}$, then D is convergent if and only if $r(D) < 1$.*

PROOF. (\Rightarrow). First, we show that if $T \in \mathcal{A}$ is convergent then $I - T$ is invertible. Suppose T is convergent, choose a fixed n large enough such that $\|T^n\| < 1$. Then $I - T^n$ is invertible. Since $I - T^n = (I - T) \sum_{k=0}^{n-1} T^k$, it follows that $I - T$ is invertible.

Now, note that if D is convergent then so is $(1/t)D$ for any complex number t such that $|t| \geq 1$. Thus $I - (1/t)D$ is invertible. Therefore, $tI - D$ is invertible. This implies that $r(D) < 1$.

(\Leftarrow). Suppose $r(D) < 1$. Then choose a fixed $\epsilon > 0$ such that $r(D) + \epsilon < 1$. For n large enough, we have $\|D^n\|^{1/n} < r(D) + \epsilon$. Thus, $\|D^n\| < (r(D) + \epsilon)^n$. Therefore, D is convergent. \square

For any stable element $A \in \mathcal{A}$, $I - A$ is invertible. In this case, we define the Cayley transform of A as $C_A = (I - A)^{-1}(I + A)$.

THEOREM 2. *For any complex Banach algebra \mathcal{A} , an element $D \in \mathcal{A}$ is convergent if and only if $D = C_A$ for some stable $A \in \mathcal{A}$.*

PROOF. (\Leftarrow). If $D = C_A$ for some stable $A \in \mathcal{A}$, let $f(z) = (1 + z)/(1 - z)$. Then $f(z)$ is analytic in the open left half of the complex plane. In fact, $f(z)$ is a Möbius

transform that maps the open left half of the complex plane onto the open unit disk and $f(A) = C_A$. Since $f(z)$ is analytic in a neighborhood of $\sigma(A)$. By the Spectral Mapping theorem (see [1, page 208]), $\sigma(f(A)) = f(\sigma(A))$. Since $f(\sigma(A))$ is contained in the open unit disk, $\sigma(D) = \sigma(f(A))$ is contained in the open unit disk. By [Lemma 1](#), D is convergent.

(\Rightarrow). Suppose D is convergent. By [Lemma 1](#), $\sigma(D)$ is contained in the open unit disk. Thus $D + I$ is invertible. Let $A = (D + I)^{-1}(D - I)$. Define $g(z) = (z - 1)/(z + 1)$. The map $g(z)$ is analytic in the open unit disk. In fact, $g(z)$ is a *Möbius* transform that maps the open unit disk onto the open left half of the complex plane and $g(D) = A$. Similarly, by the Spectral Mapping theorem, we can obtain $\sigma(A) = \sigma(g(D)) = g(\sigma(D))$. This implies that $\sigma(A)$ is contained in the open left half of the complex plane, that is, A is stable. Solving for D from $A = (D + I)^{-1}(D - I)$, we have $D = (I - A)^{-1}(I + A)$, that is, $D = C_A$. \square

REFERENCES

- [1] J. B. Conway, *A Course in Functional Analysis*, Graduate Texts in Mathematics, vol. 96, Springer-Verlag, New York, 1985.
- [2] T. Haynes, *Stable matrices, the Cayley transform, and convergent matrices*, Int. J. Math. Math. Sci. **14** (1991), no. 1, 77–81.
- [3] P. Stein, *On the ranges of two functions of positive definite matrices*, J. Algebra **2** (1965), 350–353.
- [4] O. Taussky, *Matrices C with $C^n \rightarrow 0$* , J. Algebra **1** (1964), 5–10.

ZHIDONG PAN: DEPARTMENT OF MATHEMATICS, SAGINAW VALLEY STATE UNIVERSITY, UNIVERSITY CENTER, MI 48710, USA

E-mail address: pan@svsu.edu

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru