

## ON CONSTRAINED UNIFORM APPROXIMATION

M. A. BOKHARI

Received 22 April 2001 and in revised form 8 October 2001

The problem of uniform approximants subject to Hermite interpolatory constraints is considered with an alternate approach. The uniqueness and the convergence aspects of this problem are also discussed. Our approach is based on the work of P. Kirchberger (1903) and a generalization of Weierstrass approximation theorem.

2000 Mathematics Subject Classification: 41A05, 41A10, 41A29, 41A52.

**1. Introduction.** Let  $\pi_m$  denote the set of all polynomials of degree less than or equal to  $m$ . Let  $[a, b]$  be a closed finite interval and  $C[a, b]$  the space of real continuous functions on  $[a, b]$  with uniform norm

$$\|h\|_{\infty} = \max_{x \in [a, b]} |h(x)|. \quad (1.1)$$

Here we discuss uniform approximation of a prescribed  $f \in C[a, b]$  by the polynomials that are also Hermite interpolants to a set of given data at a finite number of preassigned points in the interval  $[a, b]$ . More precisely, we consider the following problem due to Loeb et al. [6].

**PROBLEM 1.1.** Suppose that  $k$ ,  $m$ , and  $n_i$ , where  $i = 1, 2, \dots, k$ , are positive integers with  $m \geq (\sum_{i=1}^k n_i) - 1$  and that  $\{u_i\}_{i=1}^k$  is a subset of  $[a, b]$  satisfying  $a \leq u_1 < u_2 < \dots < u_k \leq b$ . Then the problem is to find a best uniform approximant to a given  $f \in C[a, b]$  from the class

$$\Phi_{m, \beta} = \{\phi \in \pi_m : \phi^{(j)}(u_i) = \beta_{ij}, 1 \leq i \leq k, 0 \leq j \leq n_i - 1\}, \quad (1.2)$$

where

$$S_{\beta} := \{\beta_{ij} : 1 \leq i \leq k, 0 \leq j \leq n_i - 1\} \quad (1.3)$$

is a subset of the real numbers with  $\beta_{i0} = f(u_i)$ .

This problem originates from the work of Paszkowski [8, 9] who studied it by imposing the interpolatory conditions only on the values of interpolating polynomials at  $k$  distinct points of  $[a, b]$  in the sense of Lagrange interpolation. To do this, he followed the classical Tchebycheff approach of approximating a continuous function by elements of an  $n$ -dimensional Haar subspace. In [2] Deutsch discussed Paszkowski's results [8, Theorems 2 and 5] with a different method. Deutsch's work is based on a characterization theorem of best approximation that involves extreme points of the closed unit ball in the dual of the underlying space [3, Corollary 2.6]. Later, Loeb et al. [6] extended the work of Deutsch by constraining the uniform approximants with

Hermite interpolatory conditions. In fact, they discussed **Problem 1.1** through the notions of the  $n$ -dimensional extended Haar subspace of order  $v$  of  $C[a, b]$  (see [4]) and the generalized weight functions (see [7]). They also established a convergence result by giving a generalization of a theorem of de La Vallée Poussin [1, page 77].

In the present paper we continue the study of **Problem 1.1**. Our approach for its solution is based on the work of Kirchberger [5] that deals with extreme values of the error function. Uniqueness and convergence problems are also addressed in our work taking into account an extension of Weierstrass approximation theorem [10].

**2. Notations and reformulation of Problem 1.1.** For the sake of convenience, we set  $I_k := \{1, 2, \dots, k\}$ ,  $N_i := \{0, 1, \dots, n_i - 1\}$ ,  $s := \sum_{i \in I_k} n_i$ , and

$$W(x) := \prod_{i \in I_k} (x - u_i)^{n_i}. \quad (2.1)$$

The notation  $H_{s-1}(x, S_\beta)$ , where  $S_\beta$  is given in (1.3), stands for the polynomial of degree less than or equal to  $s - 1$  that satisfies the following conditions:

$$H_{s-1}^{(j)}(u_i, S_\beta) = \beta_{ij}, \quad \forall i \in I_k, \forall j \in N_i. \quad (2.2)$$

For every  $f \in C[a, b]$ , we define

$$f_{H,\beta}(x) := f(x) - H_{s-1}(x, S_\beta). \quad (2.3)$$

The error function corresponding to  $f, g \in C[a, b]$  and the set of its extreme points in  $[a, b]$  will be denoted, respectively, as follows:

$$\begin{aligned} e_{f,g}(x) &:= |f(x) - g(x)|, \\ \text{crit}(e_{f,g}) &:= \{x \in [a, b] : |e_{f,g}(x)| = \|e_{f,g}\|_\infty\}. \end{aligned} \quad (2.4)$$

Let  $\pi_m^*$  denote the  $(m - s + 1)$ -dimensional subspace of  $\pi_m$  generated by the polynomials  $x^j W(x)$ ,  $j = 0, 1, 2, \dots, m - s$ , where  $W$  is given by (2.1). The following remark gives an explicit representation of the elements of  $\Phi_{m,\beta}$  (see (1.2)).

**REMARK 2.1.** A typical element  $\phi^*$  in the approximating class  $\Phi_{m,\beta}$  is of the form

$$\phi^*(x) = H_{s-1}(x, S_\beta) + q^*(x), \quad (2.5)$$

where  $q^* \in \pi_m^*$ . This shows that  $\phi^* \in \Phi_{m,\beta}$  is a best approximant to  $f$  from the class  $\Phi_{m,\beta}$  if and only if  $q^* \in \pi_m^*$  is a best approximant to  $f_{H,\beta}$  from the class  $\pi_m^*$ . In particular, if  $m = s - 1$  then  $H_{s-1}(x, S_\beta)$  will be the best approximation to  $f$ . For this obvious reason we assume that  $m \geq s$  in the rest of the paper.

In view of the above remark, **Problem 1.1** can be reformulated as follows.

**PROBLEM 2.2.** For a given function  $f \in C[a, b]$ , find a best approximation to  $f_{H,\beta}$  (see (2.3)) in the uniform norm from the class  $\pi_m^*$ .

**3. Characterization of best approximation.** This section deals with a necessary and sufficient condition for a solution of [Problem 2.2](#). We note that every  $q \in \pi_m^*$  can be expressed as

$$q(x) = W(x)R_q(x), \quad (3.1)$$

where  $R_q(x)$  is a polynomial of degree at most  $m-s$ . For an  $f \in C[a,b]$  and a  $q \in \pi_m^*$ , we set

$$E_{f_{H,\beta},q}(x) := \frac{f_{H,\beta}(x)}{W(x)} - R_q(x). \quad (3.2)$$

An alternate form of the characterization theorem [6, Theorem 3.1] that solves [Problem 1.1](#) may be stated as follows.

**THEOREM 3.1.** *Let  $f \in C[a,b]$  such that  $f_{H,\beta} \notin \pi_m^*$ . Then  $q^*$  is a best uniform approximant to  $f_{H,\beta}$  from the class  $\pi^*$  if and only if there exist  $N$  points  $\alpha_i \in \text{crit}(e_{f_{H,\beta},q^*})$  satisfying the following conditions:*

- (a)  $N = m-s+2$ ;
- (b)  $a \leq \alpha_1 < \alpha_2 < \dots < \alpha_N \leq b$ ;
- (c)  $\text{sgn}(E_{f_{H,\beta},q^*}(\alpha_i)) = (-1)^{i+1} \text{sgn}(E_{f_{H,\beta},q^*}(\alpha_1))$ , for all  $i = 2, 3, \dots, N$ .

Our method of proof is based on the following lemma which may be found in the standard texts of approximation theory, for example, [9, Lemma 7.1].

**LEMMA 3.2.** *Let  $Y$  be a linear subspace of  $C[a,b]$  and let  $h \in C[a,b]$ . Then  $g^* \in Y$  is a best uniform approximant to  $f$  in  $Y$  if and only if there does not exist any  $g \in Y$  such that*

$$\{h(x) - g^*(x)\}g(x) > 0 \quad (3.3)$$

for all  $x \in \text{crit}(e_{f,g^*})$  (see (2.3)).

**REMARK 3.3.** If we set  $Y = \pi_m^*$  and  $h = f_{H,\beta}$  in the above lemma, then the necessary and sufficient condition for  $q^* \in \pi_m^*$  to be a best approximation to  $f_{H,\beta}$  is that there does not exist any  $p \in \pi_m^*$  such that

$$E_{f_{H,\beta},q^*}(x)R_p(x) > 0 \quad (3.4)$$

for all  $x \in \text{crit}(e_{f_{H,\beta},q^*})$  where  $R_p(x)$  and  $E_{f_{H,\beta},q^*}$  are, respectively, given in (3.1) and (3.2). To justify this, it is enough to note that

$$\{f_{H,\beta}(x) - q^*(x)\}p(x) = E_{f_{H,\beta},q^*}(x)R_p(x)W^2(x). \quad (3.5)$$

**REMARK 3.4.** If  $f_{H,\beta} \notin \pi_m^*$ , then  $W(x)$  is a nonvanishing function on the compact set  $\text{crit}(e_{f_{H,\beta},p})$  regardless of the choice of  $p \in \pi_m^*$ . Consequently,  $f_{H,\beta}$  is continuous as well as nowhere zero on  $\text{crit}(e_{f_{H,\beta},p})$ .

**4. Proof of Theorem 3.1.** If  $q^*$  is not a best approximation to  $f_{H,\beta}$ , then by [Remark 3.3](#), there exists  $p \in \pi_m^*$  such that

$$E_{f_{H,\beta},q^*}(x)R_p(x) > 0 \quad (4.1)$$

as  $x$  ranges over  $\text{crit}(e_{f_{H,\beta},q^*})$ . We note that  $R_p$  being a polynomial of degree less than or equal to  $m-s$  (see (3.1)) changes sign at most at  $m-s$  places. Therefore, it follows

from (4.1) that  $E_{f_{H,\beta},q^*}(x)$  cannot change sign more than  $(m-s)$  times as  $x$  ranges over  $\text{crit}(e_{f_{H,\beta},q^*})$ . This contradicts [Theorem 3.1\(c\)](#) as  $N = m-s+2$ .

Conversely, assume that there are  $N$  points  $\alpha_i \in \text{crit}(e_{f_{H,\beta},q^*})$ ,  $i = 1, 2, \dots, N$ , satisfying [Theorem 3.1\(b\)](#) and [\(c\)](#) but  $N \leq m-s+1$ . For each  $i = 1, 2, 3, \dots, N-1$ , fix a point  $w_i \in (\alpha_i, \alpha_{i+1})$  such that

$$\begin{aligned} u_j &\notin [w_i, \alpha_{i+1}], \quad j = 1, 2, \dots, k, \\ (w_i, \alpha_{i+1}], \text{ crit}(e_{f_{H,\beta},q^*}) &\text{ are disjoint,} \\ \text{sgn}(E_{f_{H,\beta},q^*}(w_i)) &= \text{sgn}(E_{f_{H,\beta},q^*}(\alpha_{i+1})). \end{aligned} \quad (4.2)$$

The choice of  $w_i$ , as required above, directly follows from [Remark 3.4](#). Now we set

$$\tilde{p}(x) := \text{sgn}(E_{f_{H,\beta},q^*}(\alpha_1)) W(x) \prod_{i=1}^{N-1} (w_i - x). \quad (4.3)$$

Then  $\tilde{p} \in \pi_m^*$  with  $R_{\tilde{p}}(x) = \text{sgn}(E_{f_{H,\beta},q^*}(\alpha_1)) \prod_{i=1}^{N-1} (w_i - x)$ . We claim that

$$E_{f_{H,\beta},q^*}(\alpha) R_{\tilde{p}}(\alpha) > 0 \quad \forall \alpha \in \text{crit}(e_{f_{H,\beta},q^*}). \quad (4.4)$$

This can be seen by restricting  $\alpha$  to each set  $(\alpha_i, \alpha_{i+1}) \cap \text{crit}(e_{f_{H,\beta},q^*})$  for  $i = 0, 1, \dots, N-1$ , where  $\alpha_0 = a$ , and then using [Theorem 3.1\(c\)](#) along with (4.2). Hence by [Remark 3.3](#), we note that  $q^*$  cannot be a best uniform approximant to  $f_{H,\beta}$  from the class  $\pi_m^*$ . This completes the proof.  $\square$

**REMARK 4.1.** An immediate consequence of [Theorem 3.1](#) is that  $H_{s-1}(x, S_\beta) + q^*(x)$  is a best uniform approximant to  $f(x)$  from the class  $\Phi_{m,\beta}$ .

**5. Uniqueness.** We retain the setting of the previous sections in order to establish the uniqueness of the solution of [Problem 2.2](#). More precisely, we prove the following theorem.

**THEOREM 5.1.** *There is exactly one best uniform approximant  $p^*$  to  $f_{H,\beta}$  from  $\pi_m^*$ .*

If  $q^*$  is another best uniform approximant to  $f_{H,\beta}$  from  $\pi_m^*$ , then there exist  $N$  points  $\alpha_i \in \text{crit}(e_{f_{H,\beta},q^*})$ ,  $i = 1, 2, \dots, N$ , satisfying [Theorem 3.1\(a\)](#), [\(b\)](#), and [\(c\)](#). Using the properties of best approximant to  $f_{H,\beta}$ , we observe that  $|f_{H,\beta}(\alpha_i) - p^*(\alpha_i)| \leq \|e_{f_{H,\beta},p^*}\|_\infty = \|e_{f_{H,\beta},q^*}\|_\infty = |f_{H,\beta}(\alpha_i) - q^*(\alpha_i)|$  for each  $i = 1, 2, \dots, N$ , and consequently,

$$|E_{f_{H,\beta},p^*}(\alpha_i)| \leq |E_{f_{H,\beta},q^*}(\alpha_i)|. \quad (5.1)$$

We set  $D(x) := R_{p^*}(x) - R_{q^*}(x)$ . Then  $D \in \pi_{m-s}$  and

$$D(\alpha_i) = E_{f_{H,\beta},q^*}(\alpha_i) - E_{f_{H,\beta},p^*}(\alpha_i), \quad i = 1, 2, \dots, N. \quad (5.2)$$

Note that if  $D(\alpha_i) \neq 0$  for any  $i = 1, 2, \dots, N$ , then by (5.2),  $\text{sgn}(D(\alpha_i)) = \text{sgn}(E_{f_{H,\beta},q^*}(\alpha_i))$ . Thus, the polynomial  $D$  has either a double zero at  $\alpha_i$ , or it has a zero in  $(\alpha_i, \alpha_{i+1})$  implying that  $D \equiv 0$ . Hence  $R_{p^*}(x) = R_{q^*}(x)$ , and consequently,  $p^* = q^*$ .

**6. Convergence.** In this section, we discuss the convergence of the sequence of best uniform approximants  $\{q_k^*\}_{k=s-1}^\infty$  to  $f_{H,\beta}$  with the conditions that  $f$  is *sufficiently differentiable* and the set  $S_\beta$  (see [Problem 1.1](#)) is replaced by

$$S_f = \{f^{(j)}(u_i) : j \in N_i, i \in I_k\}. \quad (6.1)$$

In this case, we write  $f_{H,f}$  and  $\Phi_{m,f}$  instead of  $f_{H,\beta}$  and  $\Phi_{m,\beta}$  (see [\(2.3\)](#)).

**THEOREM 6.1.** *Assume that  $f \in C^{n^*}[a,b]$  with  $n^* = (\max_{i \in I_k} n_i) - 1$  and that the set  $S_\beta$  (see [\(1.3\)](#)) is replaced by  $S_f$ . If  $q_m^* \in \pi_m^*$  is the best approximant to  $f_{H,f}$  in the sense of [Theorem 3.1](#), then*

$$\lim_{m \rightarrow \infty} \|q_m^* - f_{H,f}\|_\infty = 0. \quad (6.2)$$

Consequently, the sequence  $\{q_m^* + H_{s-1}(\cdot, S_f)\}_{m=s-1}^\infty$  will converge uniformly to  $f$ .

The crux of the proof of this theorem is in an extension of a result based on the Weierstrass approximation theorem [10, page 160]. We state it in the next lemma without proof as it is a routine exercise.

**LEMMA 6.2.** *For any  $f \in C^r[a,b]$ , and for a given  $\varepsilon > 0$ , there exists a polynomial  $p$  such that*

$$\|f^{(j)} - p^{(j)}\|_\infty < \varepsilon \quad (6.3)$$

for all  $j = 0, 1, 2, \dots, r$ .

**PROOF OF THEOREM 6.1.** In the notations of [\(2.3\)](#) and [\(6.1\)](#), we can write

$$f_{H,f}(x) := f(x) - H_{s-1}(x, S_f). \quad (6.4)$$

The polynomial  $H_{s-1}(x, S_f)$  due to its interpolation properties may be expressed as

$$H_{s-1}(x, S_f) = \sum_{l \in I_k} \sum_{n \in N_l} f^{(n)}(u_l) L_{n,l}(x), \quad (6.5)$$

where  $L_{n,l}(x)$  are the fundamental polynomials of degree  $s-1$  satisfying the conditions

$$L_{n,l}^{(j)}(u_i) = \begin{cases} 1 & \text{for } l = i, n = j, \\ 0 & \text{otherwise.} \end{cases} \quad (6.6)$$

For a given  $\varepsilon > 0$ , we can fix a polynomial  $p$  of degree  $r > s$  such that (see [Lemma 6.2](#))

$$\|f^{(j)} - p^{(j)}\|_\infty < \frac{\varepsilon}{2\tau}, \quad j = 0, 1, \dots, n^*, \quad (6.7)$$

where  $\tau = \max\{1, \lambda\}$  with  $\lambda = \max_{x \in [a,b]} \sum_{l \in I_k} \sum_{n \in N_l} |L_{n,l}(x)|$ . From [\(6.1\)](#), [\(6.5\)](#), and [\(6.7\)](#) it follows that

$$\max_{x \in [a,b]} |H_{s-1}(x, S_f) - H_{s-1}(x, S_p)| < \frac{\varepsilon}{2}. \quad (6.8)$$

We set  $q(x) := p(x) - H_{s-1}(x, S_p)$ . Then  $q \in \pi_r$  and  $q^{(j)}(u_i) = 0$  for  $j \in N_i$  and  $i \in I_k$ . Hence,  $W$  as defined in [\(2.1\)](#) is a factor of the polynomial  $q$ . This shows that  $q \in \pi_r^*$ . Using [\(6.4\)](#), [\(6.5\)](#), and [\(6.8\)](#) it can be seen that

$$\|f_{H,f} - q\|_\infty < \varepsilon. \quad (6.9)$$

Now consider the best uniform approximant  $q_r^*$  to  $f_H$  from  $\pi_r^*$  (see [Theorem 3.1](#)) and note that

$$\varepsilon > \|f_{H,f} - q\|_\infty \geq \|f_{H,f} - q_r^*\|_\infty \geq \|f_{H,f} - q_m^*\|_\infty \quad (6.10)$$

for all  $m \geq r$ . The last inequality follows from the relation  $\pi_r^* \subseteq \pi_m^*$ . This proves the desired result.  $\square$

**ACKNOWLEDGMENT.** The author acknowledges with gratitude the research facilities available at King Fahd University of Petroleum and Minerals, Saudi Arabia, and the University of Alberta, Canada, during the preparation of this paper.

#### REFERENCES

- [1] E. W. Cheney, *Introduction to Approximation Theory*, McGraw-Hill, New York, 1966.
- [2] F. Deutsch, *On uniform approximation with interpolatory constraints*, J. Math. Anal. Appl. **24** (1968), 62–79.
- [3] F. R. Deutsch and P. H. Maserick, *Applications of the Hahn-Banach theorem in approximation theory*, SIAM Rev. **9** (1967), 516–530.
- [4] S. Karlin and W. J. Studden, *Tchebycheff Systems: With Applications in Analysis and Statistics*, Pure and Applied Mathematics, vol. 15, Interscience Publishers, New York, 1966.
- [5] P. Kirchberger, *Über Tchebychefsche annäherungsmethoden*, Math. Ann. **57** (1903), 509–540 (German).
- [6] H. L. Loeb, D. G. Moursund, L. L. Schumaker, and G. D. Taylor, *Uniform generalized weight function polynomial approximation with interpolation*, SIAM J. Numer. Anal. **6** (1969), 284–293.
- [7] D. G. Moursund, *Chebyshev approximation using a generalized weight function*, SIAM J. Numer. Anal. **3** (1966), 435–450.
- [8] S. Paszkowski, *Sur l'approximation uniforme avec des noeuds*, Ann. Polon. Math. **2** (1955), 118–135 (French).
- [9] ———, *On approximation with nodes*, Rozprawy Mat. **14** (1957), 63.
- [10] T. J. Rivlin, *The Chebyshev Polynomials*, Pure and Applied Mathematics, John Wiley & Sons, New York, 1974.

M. A. BOKHARI : DEPARTMENT OF MATHEMATICAL SCIENCES, KING FAHD UNIVERSITY OF PETROLEUM AND MINERALS, DHAHRAH 31261, SAUDI ARABIA

*E-mail address:* [mbokhari@kfupm.edu.sa](mailto:mbokhari@kfupm.edu.sa)

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)