

ON THE UNIVALENCY FOR CERTAIN SUBCLASS OF ANALYTIC FUNCTIONS INVOLVING RUSCHEWEYH DERIVATIVES

LIU MINGSHENG

Received 20 March 2001

Let H be the class of functions $f(z)$ of the form $f(z) = z + \sum_{k=2}^{+\infty} a_k z^k$, which are analytic in the unit disk $U = \{z; |z| < 1\}$. In this paper, we introduce a new subclass $B_\lambda(\mu, \alpha, \rho)$ of H and study its inclusion relations, the condition of univalence, and covering theorem. The results obtained include the related results of some authors as their special case. We also get some new results.

2000 Mathematics Subject Classification: 30C45.

1. Introduction. Let H be the class of functions of the form

$$f(z) = z + \sum_{k=2}^{+\infty} a_k z^k \quad (1.1)$$

which are analytic in the unit disk $U = \{z; |z| < 1\}$. Let S be the subclass of H consisting of univalent functions.

For the function $f(z) = \sum_{k=1}^{+\infty} a_k z^k$ and $g(z) = \sum_{k=1}^{+\infty} b_k z^k$, let $(f * g)(z)$ denote the Hadamard product or convolution of $f(z)$ and $g(z)$, defined by

$$(f * g)(z) = \sum_{k=1}^{+\infty} a_k b_k z^k. \quad (1.2)$$

Now define the function $\phi(a, c; z)$ by

$$\phi(a, c; z) = \sum_{k=0}^{+\infty} \frac{(a)_k}{(c)_k} z^{k+1}, \quad (c \neq 0, -1, -2, \dots, z \in U), \quad (1.3)$$

where

$$(\lambda)_k = \frac{\Gamma(\lambda+k)}{\Gamma(\lambda)} = \begin{cases} 1, & k = 0, \\ \lambda(\lambda+1) \cdots (\lambda+k-1), & k \in N = \{1, 2, \dots\}. \end{cases} \quad (1.4)$$

It follows from [4] that

$$z[\phi(c, c+1)]' = c\phi(c+1, c+1) - (c-1)\phi(c, c+1). \quad (1.5)$$

Carlson and Shaffer [2] defined a linear operator $L(a, c)$ on H by using the Hadamard product

$$L(a, c)f = \phi(a, c; z) * f(z), \quad f \in H. \quad (1.6)$$

It is known in [2] that $L(a, c)$ maps H into itself. If $a \neq 0, -1, -2, \dots$, then $L(a, c)$ has a continuous inverse $L(c, a)$. Clearly, $L(a, a)$ is the unit operator. Also, if $c > a > 0$, then $L(a, c)$ has the integral representation

$$L(a, c)f(z) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 u^{a-2}(1-u)^{c-a-1} f(uz) du. \quad (1.7)$$

Ruscheweyh [7] introduced an operator $D^\lambda : H \rightarrow H$ defined by the Hadamard product or convolution

$$D^\lambda f(z) = \frac{z}{(1-z)^{\lambda+1}} * f(z), \quad (\lambda > -1, z \in U), \quad (1.8)$$

which implies that

$$\begin{aligned} D^n f(z) &= \frac{z(z^{n-1}f(z))^{(n)}}{n!}, \quad (n \in N_0 = \{0, 1, 2, \dots\}), \\ D^\lambda f(z) &= L(\lambda+1, 1)f(z). \end{aligned} \quad (1.9)$$

Assume $\alpha > 0, \mu > 0, \lambda > -1, \rho < 1$, a function $p(z) = 1 + p_1z + p_2z^2 + \dots$ is said to be in the class P_ρ if and only if $p(z)$ is analytic in the unit disk U and $\operatorname{Re} p(z) > \rho$, $z \in U$; a function $f(z) \in H$ is said to be in the class $B_\lambda(\mu, \alpha, \rho)$ if and only if it satisfies

$$\operatorname{Re} \left[(1-\mu) \left(\frac{D^\lambda f(z)}{z} \right)^\alpha + \mu (D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] > \rho, \quad z \in U, \quad (1.10)$$

where the power are understood as principle values. Below we apply this agreement. It is obvious that the subclass $B_0(1, \alpha, 0)$ is the subclass of Bazilevič functions, which is the subclass of univalent functions S , let $B(\alpha, \rho) \equiv B_0(1, \alpha, \rho)$. The subclass $B_0(1, \alpha, \rho)$ ($0 \leq \rho < 1$) has been studied by Bazilevič [1], Singh [8], respectively. $B_0(0, \alpha, \rho)$ ($\rho < 1$) has been studied by Liu [5]. The subclass $B_0(\lambda, 1, \rho)$ ($0 \leq \rho < 1$) has been studied by Chichra [3], Ding et al. [4], respectively.

In this paper, we study the properties of $B_\lambda(\mu, \alpha, \rho)$. The results obtained generalize the related works of some authors. We also obtained some new univalent criterions.

2. Some lemmas

LEMMA 2.1 [4]. *Let $\rho < 1, 0 < u < 1, F(z) \in P_\rho$, then for $|z| = r < 1$,*

$$\operatorname{Re} [F(z) - F(uz)] \geq \frac{2(1-\rho)(u-1)r}{(1+r)(1+ur)}, \quad (2.1)$$

and the inequality is sharp.

LEMMA 2.2. *Let $c > 0$, $\mu > 0$, $\rho < 1$, $p(z) = 1 + p_1 z + p_2 z^2 + \dots$ be analytic in U . If*

$$\operatorname{Re}[p(z) + c\mu z p'(z)] > \rho, \quad z \in U, \quad (2.2)$$

then for $|z| = r < 1$,

$$\begin{aligned} \operatorname{Re}[p(z) + c\mu z p'(z)] &\geq 2\rho - 1 + \frac{2(1-\rho)}{\mu(1+r)} + 2(1-\rho)\left(1 - \frac{1}{\mu}\right) \frac{1}{c\mu} \int_0^1 \frac{u^{1/c\mu-1}}{1+ur} du, \\ \operatorname{Re}[p(z) + c\mu z p'(z)] &\geq 2\rho - 1 + \frac{1-\rho}{\mu} + 2(1-\rho)\left(1 - \frac{1}{\mu}\right) \frac{1}{c\mu} \int_0^1 \frac{u^{1/c\mu-1}}{1+u} du, \end{aligned} \quad (2.3)$$

and these results are sharp.

PROOF. Set $F(z) = p(z) + c\mu z p'(z)$, then it follows from (2.2) that $F(z) \in P_\rho$, and

$$zF(z) = (1 - c\mu)[zp(z)] + c\mu z[zp(z)]' = L\left(\frac{1}{c\mu} + 1, \frac{1}{c\mu}\right)[zp(z)] \quad (2.4)$$

that is,

$$zp(z) = L\left(\frac{1}{c\mu}, \frac{1}{c\mu} + 1\right)[zF(z)] = \frac{1}{c\mu} \int_0^1 u^{1/c\mu-1} zF(uz) du. \quad (2.5)$$

Let $b = 1/c\mu$, then

$$p(z) = b \int_0^1 u^{b-1} F(uz) du. \quad (2.6)$$

According to (1.5) and (2.5), we get

$$\begin{aligned} z[zp(z)]' &= [z(\phi(b, b+1; z))'] * [zF(z)] \\ &= bL(b+1, b+1)[zF(z)] - (b-1)L(b, b+1)[zF(z)] \\ &= bzF(z) - b(b-1)z \int_0^1 u^{b-1} F(uz) du. \end{aligned} \quad (2.7)$$

On the other hand, we have

$$[zp(z)]' = p(z) + zp'(z). \quad (2.8)$$

Thus

$$\begin{aligned} p(z) + c\mu z p'(z) &= (1 - c)p(z) + c[zp(z)]' \\ &= bcF(z) + b(1 - c) \int_0^1 u^{b-1} F(uz) du - bc(b-1)z \int_0^1 u^{b-1} F(uz) du \\ &= bcF(z) + b(1 - bc) \int_0^1 u^{b-1} F(uz) du. \end{aligned} \quad (2.9)$$

If $\mu > 1$, then $0 < bc = 1/\mu < 1$, and

$$\begin{aligned} \operatorname{Re}[p(z) + czp'(z)] &= bc \operatorname{Re}[F(z)] + b(1-bc) \int_0^1 u^{b-1} \operatorname{Re}[F(uz)] du \\ &\geq bc \cdot \frac{1-(1-2\rho)r}{1+r} + b(1-bc) \int_0^1 u^{b-1} \frac{1-(1-2\rho)ur}{1+ur} du \quad (2.10) \\ &= 2\rho-1 + \frac{2(1-\rho)}{\mu(1+r)} + 2(1-\rho) \left(1 - \frac{1}{\mu}\right) \frac{1}{c\mu} \int_0^1 \frac{u^{1/c\mu-1}}{1+ur} du. \end{aligned}$$

If $0 < \mu \leq 1$, then $bc = 1/\mu \geq 1$, so that it follows from [Lemma 2.1](#) and [\(2.9\)](#) that

$$\begin{aligned} \operatorname{Re}[p(z) + czp'(z)] &= \operatorname{Re} \left[bcF(z) - b(bc-1) \int_0^1 u^{b-1} F(uz) du \right] \\ &= \operatorname{Re} F(z) + b(bc-1) \int_0^1 u^{b-1} \operatorname{Re}[F(z) - F(uz)] du \\ &\geq \frac{1-(1-2\rho)r}{1+r} + b(bc-1) \int_0^1 u^{b-1} \frac{2(1-\rho)(u-1)r}{(1+r)(1+ur)} du \quad (2.11) \\ &= 2\rho-1 + \frac{2(1-\rho)}{\mu(1+r)} + 2(1-\rho) \left(1 - \frac{1}{\mu}\right) \frac{1}{c\mu} \int_0^1 \frac{u^{1/c\mu-1}}{1+ur} du. \end{aligned}$$

Since the function

$$2\rho-1 + \frac{2(1-\rho)}{\mu(1+r)} + 2(1-\rho) \left(1 - \frac{1}{\mu}\right) \frac{1}{c\mu} \int_0^1 \frac{u^{1/c\mu-1}}{1+ur} du \quad (2.12)$$

is decreasing with respect to r , therefore

$$\operatorname{Re}[p(z) + czp'(z)] \geq 2\rho-1 + \frac{1-\rho}{\mu} + 2(1-\rho) \left(1 - \frac{1}{\mu}\right) \frac{1}{c\mu} \int_0^1 \frac{u^{1/c\mu-1}}{1+u} du. \quad (2.13)$$

Note that

$$p_{\mu,c,\rho}(z) = \frac{1}{c\mu} \int_0^1 u^{1/c\mu-1} \frac{1+(1-2\rho)uz}{1-uz} du, \quad (2.14)$$

satisfies [\(2.2\)](#), we obtain that the inequalities [\(2.3\)](#) are sharp. \square

LEMMA 2.3 [6]. *Let $p(z) = 1 + p_1 z + \dots \in P_\rho$, then $|p_k| \leq 2 - 2\rho$, $k = 1, 2, \dots$*

3. Main results

THEOREM 3.1. *Let $\alpha > 0$, $\beta > 0$, $\lambda \geq 0$, $f(z) \in H$ and*

$$\operatorname{Re} \left[(D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] > \frac{\lambda}{\beta+\lambda}, \quad z \in U, \quad (3.1)$$

then $f(z)$ is univalent in U , that is, $B_\lambda(1, \alpha, \lambda/(\beta+\lambda)) \subset B(\alpha, 0)$.

PROOF. Since $\lambda/(\beta+\lambda) \geq 0$, we have

$$B\left(\alpha, \frac{\lambda}{\beta+\lambda}\right) \subset B(\alpha, 0). \quad (3.2)$$

Let $T = D^\lambda f(z)$, then T is a linear and homeomorphism from $B_\lambda(1, \alpha, \lambda/(\beta + \lambda))$ onto $B(\alpha, \lambda/(\beta + \lambda))$, therefore it follows from (3.2) that

$$B_\lambda\left(1, \alpha, \frac{\lambda}{\beta + \lambda}\right) = T^{-1}B\left(\alpha, \frac{\lambda}{\beta + \lambda}\right) \subset T^{-1}B(\alpha, 0) = B_0(1, \alpha, 0) \equiv B(\alpha, 0). \quad (3.3)$$

Hence the proof is completed. \square

Since $\lim_{\beta \rightarrow \infty} \lambda/(\beta + \lambda) = 0$, the following corollary follows from [Theorem 3.1](#).

COROLLARY 3.2. *Let $\alpha > 0$, $0 < \rho < 1$, $\lambda \geq 0$, then*

$$B_\lambda(1, \alpha, \rho) \subset B(\alpha, 0) \subset S. \quad (3.4)$$

THEOREM 3.3. *Let $\mu_2 \geq \mu_1 > 0$, $1 > \rho_2 \geq \rho_1$, then*

$$B_\lambda(\mu_2, \alpha, \rho_2) \subset B_\lambda(\mu_1, \alpha, \rho_1). \quad (3.5)$$

PROOF. First, it is obvious that

$$B_\lambda(\mu_2, \alpha, \rho_2) \subset B_\lambda(\mu_2, \alpha, \rho_1). \quad (3.6)$$

Therefore we only need to verify that

$$B_\lambda(\mu_2, \alpha, \rho_1) \subset B_\lambda(\mu_1, \alpha, \rho_1). \quad (3.7)$$

Let $p(z) = [D^\lambda f(z)/z]^\alpha$ for $f \in B_\lambda(\mu_2, \alpha, \rho_1)$, where the power are understood as principle values, then $p(z) = 1 + (1 + \lambda)\alpha a_2 z + \dots$ is analytic in U and

$$[D^\lambda f(z)]^\alpha = z^\alpha p(z). \quad (3.8)$$

By taking the derivatives in the both sides of (3.8), we obtain

$$(1 - \mu_2) \left(\frac{D^\lambda f(z)}{z} \right)^\alpha + \mu_2 (D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} = p(z) + \frac{\mu_2}{\alpha} z p'(z). \quad (3.9)$$

Since $f \in B_\lambda(\mu_2, \alpha, \rho_1)$, we have

$$\operatorname{Re} \left[p(z) + \frac{\mu_2}{\alpha} z p'(z) \right] > \rho_1, \quad z \in U. \quad (3.10)$$

According to [Lemma 2.2](#), we obtain

$$\begin{aligned} \operatorname{Re} \left[p(z) + \frac{\mu_1}{\alpha} z p'(z) \right] &\geq 2\rho_1 - 1 + \frac{2(1 - \rho_1)}{\mu(1 + r)} + 2(1 - \rho_1) \left(1 - \frac{1}{\mu}\right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1 + ur} du \\ &\geq 2\rho_1 - 1 + \frac{2(1 - \rho_1)}{\mu} + 2(1 - \rho_1) \left(1 - \frac{1}{\mu}\right) \cdot \frac{1}{2} > \rho_1 \end{aligned} \quad (3.11)$$

for $\mu = \mu_2/\mu_1 \geq 1$. Hence it follows from (3.6) and (3.11) that

$$\operatorname{Re} \left[(1 - \mu_1) \left(\frac{D^\lambda f(z)}{z} \right)^\alpha + \mu_1 (D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] = \operatorname{Re} \left[p(z) + \frac{\mu_1}{\alpha} z p'(z) \right] > \rho_1, \quad (3.12)$$

that is,

$$f \in B_\lambda(\mu_2, \alpha, \rho_1). \quad (3.13)$$

Hence

$$B_\lambda(\mu_2, \alpha, \rho_1) \subset B_\lambda(\mu_1, \alpha, \rho_1). \quad (3.14)$$

□

According to **Theorem 3.3** and **Corollary 3.2**, we have the following corollary.

COROLLARY 3.4. *Let $\lambda \geq 0$, $\mu \geq 1$, $0 < \rho < 1$, then*

$$B_\lambda(\mu, \alpha, \rho) \subset B_\lambda(1, \alpha, \rho) \subset S. \quad (3.15)$$

THEOREM 3.5. *Let $\alpha > 0$, $\lambda \geq 0$, $\mu > 0$, $\rho < 1$. If $f \in B_\lambda(\mu, \alpha, \rho)$, then for $|z| = r < 1$,*

$$\begin{aligned} \operatorname{Re} \left[(D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] &\geq 2\rho - 1 + \frac{2(1-\rho)}{\mu(1+r)} + 2(1-\rho) \left(1 - \frac{1}{\mu}\right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+ur} du, \\ \operatorname{Re} \left[(D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] &\geq 2\rho - 1 + \frac{1-\rho}{\mu} + 2(1-\rho) \left(1 - \frac{1}{\mu}\right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+u} du, \end{aligned} \quad (3.16)$$

and these results are sharp.

PROOF. Let $p(z) = [D^\lambda f(z)/z]^\alpha$ for $f \in B_\lambda(\mu, \alpha, \rho)$, where the power are understood as principle values, then

$$p(z) = 1 + (1 + \lambda) \alpha a_2 z + \dots \quad (3.17)$$

is analytic in U and

$$[D^\lambda f(z)]^\alpha = z^\alpha p(z). \quad (3.18)$$

By taking the derivatives in the both sides of (3.18), we obtain

$$(1 - \mu) \left(\frac{D^\lambda f(z)}{z} \right)^\alpha + \mu (D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} = p(z) + \frac{\mu}{\alpha} z p'(z). \quad (3.19)$$

Since $f \in B_\lambda(\mu, \alpha, \rho)$, we have

$$\operatorname{Re} \left[p(z) + \frac{\mu}{\alpha} z p'(z) \right] > \rho, \quad z \in U. \quad (3.20)$$

According to [Lemma 2.2](#), we obtain

$$\begin{aligned}
 \operatorname{Re} \left[(D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] &= \operatorname{Re} \left[p(z) + \frac{1}{\alpha} z p'(z) \right] \\
 &\geq 2\rho - 1 + \frac{2(1-\rho)}{\mu(1+r)} + 2(1-\rho) \left(1 - \frac{1}{\mu} \right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+ur} du, \\
 \operatorname{Re} \left[(D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] &= \operatorname{Re} \left[p(z) + \frac{1}{\alpha} z p'(z) \right] \\
 &\geq 2\rho - 1 + \frac{1-\rho}{\mu} + 2(1-\rho) \left(1 - \frac{1}{\mu} \right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+u} du. \tag{3.21}
 \end{aligned}$$

Note that,

$$f_{\lambda, \mu, \alpha, \rho}(z) = L(1, \lambda+1) \left\{ z \left[\frac{\alpha}{\mu} \int_0^1 u^{\alpha/\mu-1} \frac{1+(1-2\rho)uz}{1-uz} du \right]^{1/\alpha} \right\} \in B_\lambda(\mu, \alpha, \rho), \tag{3.22}$$

we obtain that inequalities [\(3.16\)](#) are sharp. \square

REMARK 3.6. Setting $\lambda = 0$, $\alpha = 1$ in [Theorem 3.5](#), we get [[4](#), Theorem 1(ii)].

THEOREM 3.7. Let $\alpha > 0$, $\mu > 0$, $\lambda \geq 0$, $\rho_0 \leq \rho < 1$, then $B_\lambda(\mu, \alpha, \rho) \subset B_\lambda(1, \alpha, \rho_1) \subset S$, where

$$\rho_0 = 1 - \frac{1}{2 - 1/\mu - 2(1-1/\mu)(\alpha/\mu) \int_0^1 (u^{\alpha/\mu-1}/(1+u)) du}, \tag{3.23}$$

and the constant ρ_0 cannot be replaced by any smaller one.

PROOF. Let $f(z) \in B_\lambda(\mu, \alpha, \rho)$, then it follows from [Theorem 3.5](#) that

$$\operatorname{Re} \left[(D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] > \rho_1, \quad z \in U, \tag{3.24}$$

where

$$\begin{aligned}
 \rho_1 &= 2\rho - 1 + \frac{1-\rho}{\mu} + 2(1-\rho) \left(1 - \frac{1}{\mu} \right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+u} du \\
 &= 1 - (1-\rho) \left[2 - \frac{1}{\mu} - 2 \left(1 - \frac{1}{\mu} \right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+u} du \right]. \tag{3.25}
 \end{aligned}$$

Since

$$\frac{1}{2} < \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+u} du = \frac{1}{2} + \int_0^1 \frac{u^{\alpha/\mu}}{(1+u)^2} du < 1, \tag{3.26}$$

so that

$$\max \left\{ 1, \frac{1}{\mu} \right\} > 2 - \frac{1}{\mu} - 2 \left(1 - \frac{1}{\mu} \right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+u} du > \min \left\{ 1, \frac{1}{\mu} \right\} > 0. \tag{3.27}$$

Therefore from $\rho_0 \leq \rho < 1$, we have

$$\rho_1 > 1 - (1 - \rho_0) \left[2 - \frac{1}{\mu} - 2 \left(1 - \frac{1}{\mu}\right) \frac{\alpha}{\mu} \int_0^1 \frac{u^{\alpha/\mu-1}}{1+u} du \right] = 0. \quad (3.28)$$

Hence it follows from (1.10) and [Corollary 3.2](#) that $f(z) \in B_\lambda(1, \alpha, \rho_1) \subset S$ and $f(z)$ is univalent in U , hence

$$B_\lambda(\mu, \alpha, \rho) \subset B_\lambda(1, \alpha, \rho_1) \subset S \quad (3.29)$$

and the constant ρ_0 cannot be replaced by any smaller one from [Theorem 3.5](#). \square

REMARK 3.8. Setting $\lambda = 0$, $\alpha = 1$ in [Theorem 3.7](#), we get [4, Theorem 2]; setting $\lambda = 0$, $\mu = 1$ in [Theorem 3.7](#), we get the result of [1].

Setting $\lambda = 0$, $\mu = \alpha > 0$ in [Theorem 3.7](#), we have the following corollary.

COROLLARY 3.9. *If $f(z) \in H$, and*

$$\operatorname{Re} \left[(1 - \alpha) \left(\frac{f(z)}{z} \right)^\alpha + \alpha f'(z) \left(\frac{f(z)}{z} \right)^{\alpha-1} \right] > \rho_\alpha = \frac{(\alpha - 1)(1 - 2 \ln 2)}{\alpha + (\alpha - 1)(1 - 2 \ln 2)}, \quad z \in U, \quad (3.30)$$

then $f(z)$ is univalent in U , and the result is sharp.

REMARK 3.10. We note that $\rho_\alpha < 0$ for $\alpha > 1$.

Setting $\mu = \alpha = 3$, $\lambda = 1$ in [Theorem 3.7](#), we have the following corollary.

COROLLARY 3.11. *If $f(z) \in H$, and*

$$\operatorname{Re} \left\{ z \left[(f'(z))^3 \right]' + (f'(z))^3 \right\} > \frac{2 - 4 \ln 2}{5 - 4 \ln 2} \approx -0.34, \quad z \in U, \quad (3.31)$$

then $f(z)$ is univalent in U .

THEOREM 3.12. *Let $f(z) = z + \sum_{k=2}^{+\infty} a_k z^k \in B_\lambda(\mu, \alpha, \rho)$, then*

$$|a_2| \leq \frac{2 - 2\rho}{(1 + \lambda)(\alpha + \mu)}, \quad (3.32)$$

and the inequality is sharp, with the extremal function defined $f_{\lambda, \mu, \alpha, \rho}(z)$ by (3.22).

PROOF. Since $f(z) = z + \sum_{k=2}^{+\infty} a_k z^k \in B_\lambda(\mu, \alpha, \rho)$, we obtain

$$\begin{aligned} \operatorname{Re} \left[(1 - \mu) \left(\frac{D^\lambda f(z)}{z} \right)^\alpha + \mu (D^\lambda f(z))' \left(\frac{D^\lambda f(z)}{z} \right)^{\alpha-1} \right] \\ = \operatorname{Re} [1 + (1 + \lambda)(\alpha + \mu)z + \dots] > \rho. \end{aligned} \quad (3.33)$$

Therefore, it follows from [Lemma 2.3](#) that

$$| (1 + \lambda)(\alpha + \mu)a_2 | \leq 2 - 2\rho, \quad (3.34)$$

or

$$|a_2| \leq \frac{2-2\rho}{(1+\lambda)(\alpha+\mu)}. \quad (3.35)$$

Note that $f_{\lambda,\mu,\alpha,\rho}(z) = z + ((2-2\rho)/(1+\lambda)(\alpha+\mu))z^2 + \dots \in B_\lambda(\mu, \alpha, \rho)$, we obtain that inequality (3.32) is sharp. \square

REMARK 3.13. Setting $\lambda = 0, \mu = 1$ in [Theorem 3.12](#), we get [8, Theorem 6].

THEOREM 3.14 (covering theorem). *Let $\alpha > 0, \mu > 0, \lambda \geq 0, \rho_0 \leq \rho < 1, f(z) \in B_\lambda(\mu, \alpha, \rho)$, then the unit disk U is mapped on a domain that contain the disk $|w| < r_1$, where r_1 defined by (3.23) and*

$$r_1 = \frac{(1+\lambda)(\alpha+\mu)}{2-2\rho+2(1+\lambda)(\alpha+\mu)}. \quad (3.36)$$

PROOF. Let w_0 be any complex number such that $f(z) \neq w_0$ for $z \in U$, then $w_0 \neq 0$ and

$$\frac{w_0 f(z)}{w_0 - f(z)} = z + \left(a_2 + \frac{1}{w_0} \right) z^2 + \dots, \quad (3.37)$$

is univalent in U by [Theorem 3.7](#), so

$$\left| a_2 + \frac{1}{w_0} \right| \leq 2. \quad (3.38)$$

Therefore according to [Theorem 3.12](#), we obtain

$$|w_0| \geq \frac{(1+\lambda)(\alpha+\mu)}{2-2\rho+2(1+\lambda)(\alpha+\mu)} = r_1. \quad (3.39)$$

Hence we have completed the proof. \square

Setting $\lambda = 0$ and $\mu = 1$ in [Theorem 3.14](#), we have the following corollary.

COROLLARY 3.15 (covering theorem). *Let $f(z) \in B_0(1, \alpha, \rho)$ with $\mu > 0, 0 \leq \rho < 1$, then the unit disk U is mapped on a domain that contain the disk $|w| < (1+\alpha)/(4-2\rho+2\alpha)$.*

ACKNOWLEDGMENT. This research was supported by the National Natural Science Foundation of China grant no. 19971029.

REFERENCES

- [1] I. E. Bazilevič, *On a case of integrability in quadratures of the Loewner-Kufarev equation*, Mat. Sb. (N.S.) **37**(79) (1955), 471–476.
- [2] B. C. Carlson and D. B. Shaffer, *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal. **15** (1984), no. 4, 737–745.
- [3] P. N. Chichra, *New subclasses of the class of close-to-convex functions*, Proc. Amer. Math. Soc. **62** (1976), no. 1, 37–43.
- [4] S. S. Ding, Y. Ling, and G. J. Bao, *Some properties of a class of analytic functions*, J. Math. Anal. Appl. **195** (1995), no. 1, 71–81.

- [5] M. S. Liu, *The radius of univalence for certain class of analytic functions*, Boundary Value Problems, Integral Equations and Related Problems (Beijing/Chengde, 1999) (L. Jianke and W. Guochun, eds.), World Scientific Publishing, New Jersey, 2000, pp. 122-128.
- [6] C. Pommerenke, *Univalent Functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [7] S. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc. **49** (1975), 109-115.
- [8] R. Singh, *On Bazilevič functions*, Proc. Amer. Math. Soc. **38** (1973), 261-271.

LIU MINGSHENG: DEPARTMENT OF MATHEMATICS, SOUTH CHINA NORMAL UNIVERSITY, GUANGZHOU 510631, CHINA

E-mail address: liumsh@scnu.edu.cn

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Edson Denis Leonel, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru