

CONSTRUCTIONS OF POSITIVE COMMUTATIVE SEMIGROUPS ON THE PLANE, II

REUBEN W. FARLEY

Department of Mathematical Sciences
Virginia Commonwealth University
Richmond, Virginia 23284 U.S.A.

(Received October 2, 1984 and in revised form December 10, 1984)

ABSTRACT. A positive semigroup is a topological semigroup containing a subsemigroup N isomorphic to the multiplicative semigroup of nonnegative real numbers, embedded as a closed subset of E^2 in such a way that 1 is an identity and 0 is a zero. Using results in Farley [1] it can be shown that positive commutative semigroups on the plane constructed by the techniques given in Farley [2] cannot contain an infinite number of two dimensional groups. In this work an example of such a semigroup will be constructed which does, however, contain an infinite number of one dimensional groups. Also, some preliminary results are given here concerning what types of semilattices of idempotent elements are possible for positive commutative semigroups on E^2 . In particular, we will show that there is a unique positive commutative semigroup on E^2 which is the union of connected groups and which contains five idempotent elements. Also, we will show that such semigroups having nine idempotent elements are not unique by constructing an example of such a semigroup with nine idempotent elements whose semilattice of idempotent elements is not "symmetric" and hence which is not isomorphic to the semigroup with nine idempotent elements constructed in Farley [2].

KEY WORDS AND PHRASES. *Positive semigroup, Clifford semigroup, semilattice of idempotent elements.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 22A15.

1. INTRODUCTION.

A topological semigroup is a Hausdorff space together with a continuous, associative multiplication. The author has defined a positive semigroup to be a topological semigroup containing a subsemigroup N isomorphic to the multiplicative semigroup of nonnegative real numbers, embedded as a closed subset of E^2 so that 1 is an identity and 0 is a zero [1]. Such semigroups which meet the additional requirement of being the union of groups are called positive Clifford semigroups [1].

In Farley [2] a method was given for constructing positive commutative semigroups on the plane as the union of $\{0\}$, an arbitrarily large number of two dimensional groups, and one dimensional groups which are bounding rays of the two dimensional groups. Since such semigroups are positive commutative Clifford semigroups on E^2 , they cannot contain an infinite number of two dimensional groups as shown in Farley [1]. However, in this work an example will be constructed of a positive commutative semigroup on E^2 which contains a sector of one dimensional groups, and thus contains an infinite number of one dimensional groups.

While the problem of attempting to discover what types of semilattices of idempotent elements are possible for positive commutative Clifford semigroups on E^2 appears to be difficult to answer in general, some preliminary results are given in this work. Using the techniques employed in Farley [1] we will show that there is a unique (up to isomorphism) positive commutative Clifford semigroup on E^2 which is the union of connected groups and which contains five idempotent elements. This semigroup, along with its semilattice of idempotent elements will be described herein. Also, we will show that such semigroups having nine idempotent elements are not unique by constructing an example of such a semigroup having nine idempotent elements whose semilattice of idempotent elements is, in some sense, not symmetric, and hence is not isomorphic to the semigroup with nine idempotent elements constructed in Farley [2].

2. PRELIMINARIES.

An isomorphism between two topological semigroups is a function which is both an algebraic isomorphism and a homeomorphism. Unless otherwise indicated, R will denote a semigroup isomorphic to the multiplicative semigroup of real numbers, N will denote the nonnegative members of R , and P will denote the positive members of R . For a topological space X , we will let $\Delta = \{(x,x):x \in X\}$. If R is an equivalence relation on X , X/R will denote the set of equivalence classes of X modulo R . A min thread is a closed interval $[a,b]$ together with the usual topology and multiplication defined by $x \cdot y = \min\{x,y\}$.

3. CONSTRUCTION OF EXAMPLES.

We construct first an example of a positive commutative semigroup on E^2 which contains an infinite number of one dimensional groups.

EXAMPLE 1. Let us consider two copies of $N \times N$ denoting them as $N \times N$ and $M \times M$. Also, let us form a sector of one dimensional groups by taking $N \cdot \{\text{min thread}\}$ and shrinking the base to a point. Let us call this sector N' , and let the idempotent elements on its bounding rays be denoted e_2 and e_3 , with $e_2 e_3 = e_3$. Let $(a,b)_n$ denote an element of $N \times N$, let $(a,b)_m$ denote an element of $M \times M$, and let (p, e_α) denote an element of N' , where e_α is an idempotent element. Let us define a multiplication on $S' = [(N \times N) \cup (M \times M) \cup N']$ in the following manner. Let multiplication be coordinatewise in $N \times N$, $M \times M$, and N' . Let us define $(a,b)_n \cdot (p, e_\alpha) = (bp, e_\alpha) = (p, e_\alpha) \cdot (a,b)_n$, $(p, e_\alpha) \cdot (p', e_\beta) = (pp', e_\alpha e_\beta) = (p', e_\beta) \cdot (p, e_\alpha)$, $(a,b)_n \cdot (c,d)_m = (ac, bd)_m = (c,d)_m \cdot (a,b)_n$, and $(p, e_\alpha) \cdot (x,y)_m = (0, py)_m = (x, y)_m \cdot (p, e_\alpha)$. This multiplication is easily checked to be associative, and its continuity follows from the continuity of multiplication in $N \times N$, $M \times M$, and N' . So, with the usual topology, S' is a topological semigroup. Let us define a relation R on S' in the following way. Let $\Delta \subseteq R$, and let R be symmetric by definition. Let $[(a,b)_m, (c,d)_n] \cdot R$ if and only

if $a = c$, and $b = 0 = d$. Let $[(x,y)]_m, [(p,e_3)] \in R$ if and only if $x = 0$ and $y = p$. Let $[(s,t)]_n, [(p,e_2)] \in R$ if and only if $s = 0$, and $t = p$. Finally, let $[(0,e_\alpha)], [(0,0)]_m \in R$ and $[(0,e_\alpha)], [(0,0)]_n \in R$. Then, R is clearly an equivalence relation and it is not difficult to check that R is, in fact, a closed congruence. Thus by [2] $S = S'/R$ is a topological semigroup on E^2 .

It should be noted that an example of this type cannot be constructed using just one copy of $N \times N$ and sector of one dimensional groups since $e_1 e_2$ would have to be 0 where $e_1 = (1,0)_n$. So, this example is more or less minimal.

Upon investigating what types of semilattices of idempotent elements are possible for positive Clifford semigroups on the plane, we first note that the semigroup containing five idempotent elements constructed by the method employed in Example 1 or Example 3 of Farley [2] has the following semilattice of idempotent elements, where 1 denotes the element $(1,1)_N$, f denotes $(1,1)_J$, e_1 denotes $(1,0)_N = (1,0)_J$, e_2 denotes $(0,1)_N = (0,1)_J$, and 0 denotes $(0,0)_N = (0,0)_J$:




Figure 1.




Figure 2.

For positive Clifford semigroups on E^2 which are the union of connected groups, we will show that the semigroup is unique (up to isomorphism) among those having exactly five idempotent elements.

THEOREM 1. The positive commutative Clifford semigroup S on E^2 which contains exactly five idempotent elements and is the union of connected groups is unique.

PROOF. Let $H(1)$ denote the group having the identity element 1, and let $CL[H(1)]$ denote the closure of $H(1)$. Then $CL[H(1)]$ is isomorphic to $N \times N$ as shown in Farley [1], since $CL[H(1)]$ is two dimensional by Horne [3]. Let e_1 and e_2 denote the idempotent elements on Pe_1 and Pe_2 , the bounding rays of $H(1)$, let f be the remaining idempotent element which is in $E^2 \setminus CL[H(1)]$, and let $H(f)$ be the group with identity element f . Then, the only possibilities for the semilattice of idempotent elements are the one given above (Figure 1) and the other given in Figure 2.

We now show that this semilattice is, in fact, impossible. Since S has exactly five idempotent elements and is the union of connected groups, $H(f) = E^2 \setminus CL[H(1)]$. Now, suppose $e_1 f = 0$. Let $\{X_n\}$ be a sequence in $H(f)$ such that $\{X_n\} \rightarrow e_1$. Then $\{e_1 X_n\} \rightarrow e_1$. But, $e_1 X = e_1(fX)$, since f is the identity element in $H(f)$. So, $\{e_1(fX_n)\} \rightarrow e_1$. However, each term of the sequence $\{e_1(fX_n)\}$ is 0, under the assumption that $e_1 f = 0$. This implies $\{e_1(fX_n)\} \rightarrow 0$ which is a contradiction.

A positive Clifford semigroup having nine idempotent elements as constructed by the method employed in Example 3 of Farley [2] has the following semilattice of idempotent elements, where 1 denotes $(1,1)_N$, j denotes $(1,1)_J$, m denotes $(1,1)_m$, k denotes $(1,1)_k$, e denotes $(1,0)_N = (1,0)_J$, g denotes $(0,1)_J = (1,0)_m$, h denotes $(0,1)_m = (1,0)_k$, f denotes $(0,1)_k = (0,1)_N$ and 0 denotes $(0,0)_m = (0,0)_J = (0,0)_N = (0,0)_k$: (Fig.3):




Figure 3.




Figure 4.

We now construct an example of a positive Clifford semigroup on E^2 whose semilattice of nine idempotent elements is not isomorphic to the one above, and is, in some sense, not symmetric.

EXAMPLE 2. Let us consider four copies of $N \times N$. Let us denote these copies as $N \times N$ (which will be \bar{G}), $J \times J$, $M \times M$, and $K \times K$. Let us define a multiplication on $S' = [(N \times N) \cup (J \times J) \cup (M \times M) \cup (K \times K)]$ in the following manner. Let the multiplication be coordinatewise in each copy of $N \times N$, and let the multiplication be commutative on all of S' . Let $(a,b)_j \cdot (c,d)_m = (ac,bd)_m$, $(a,b)_j \cdot (c,d)_k = (ac,bcd)_m$, $(a,b)_k \cdot (c,d)_m = (ac,abd)_m$, $(a,b)_n \cdot (c,d)_j = (ac,abd)_j$, $(a,b)_n \cdot (c,d)_m = (ac,abd)_m$, and $(a,b)_n \cdot (c,d)_k = (ac,abd)_k$, where $(a,b)_n \in (N \times N)$, $(a,b)_m \in (M \times M)$, $(a,b)_j \in (J \times J)$, and $(a,b)_k \in (K \times K)$. This multiplication is easily verified to be continuous and associative. So, with the usual topology, S' is a topological semigroup. Let us define a relation R on S' as follows. Let $\Delta \subseteq R$, and let R be symmetric by definition. Let $[(a,b)_j, (c,d)_n] \in R$ if and only if $a = c$ and $b = 0 = d$. Let $[(a,b)_j, (c,d)_m] \in R$ if and only if $b = d$ and $a = 0 = c$. Let $[(a,b)_m, (c,d)_n] \in R$ if and only if $a = c$ and $b = 0 = d$. Finally, let $[(a,b)_k, (c,d)_n] \in R$ if and only if $b = d$ and $a = 0 = c$. Then, R is easily checked to be a closed congruence, and consequently S'/R is a topological semigroup on E^2 . Let us denote by e the element $(1,0)_n = (1,0)_j$, by g the element $(0,1)_j = (0,1)_m$, by h the element $(1,0)_m = (1,0)_k$, and by f the element $(0,1)_k = (0,1)_n$. Also, let us denote by l the element $(1,1)_n$, by j the element $(1,1)_j$, by m the element $(1,1)_m$, by k the element $(1,1)_k$, and by 0 the element $(0,0)_n = (0,0)_j = (0,0)_m = (0,0)_k$. Then, the semigroup $S = S'/R$ has the semilattice of idempotent elements (Fig.4).

REFERENCES

1. FARLEY, R.W. Positive Clifford Semigroups on the Plane, Trans. Amer. Math. Soc. 151 (1970), 353-369.
2. FARLEY, R.W. Constructions of Positive Commutative Semigroups on the Plane, Bull. Inst. Math. Acad. Sinica 7 (1979), no. 4, 357-362.
3. HORNE, J.G. Real Commutative Semigroups on the Plane, Pacific J. Math. 11 (1961), 981-997.
4. HORNE, J.G. Real Commutative Semigroups on the Plane, II, Trans. Amer. Math. Soc. 104 (1962), 17-23.
5. FARLEY, R.W. An Isomorphism Theorem for Positive Commutative Semigroups on the Plane, Mat. Casopis Sloven Akad. Vied. 23 (1973), no. 2, 136-146.
6. HOFMANN, K.H., MOSTERT, P.S. Elements of Compact Semigroups, Merrill, Columbus, Ohio, 1966.
7. CLIFFORD, A.H., PRESTON, G.B. The Algebraic Theory of Semigroups, In Mathematical Surveys, Vol 1, no. 7, Amer. Math. Soc., Providence, R.I., 1961.
8. MOSTERT, P.S. Plane Semigroups, Trans. Amer. Math. Soc. 103 (1962), 320-328.
9. MOSTERT, P.S., and SHIELDS, A.L. Semigroups with Identity on a Manifold, Trans. Amer. Math. Soc. 91 (1959), 380-389.
10. PONTRJAGIN, L. Topological Groups, Gittl, Moscow, 1938; English transl., Princeton Math. Series, Vol. 2, Princeton University Press, Princeton, N.J., 1939.

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mskklai@cityu.edu.hk