

ON THE DISTRIBUTIONAL STIELTJES TRANSFORMATION

D. NIKOLIC — DESPOTOVIC and A. TAKACI

Institute of Mathematics
Novi Sad, Yugoslavia

(Received January 10, 1984 and in revised form June 24, 1985)

ABSTRACT. This paper is concerned with some general theorems on the distributional Stieltjes transformation. Some Abelian theorems are proved.

KEY WORDS AND PHRASES. *Stieltjes transforms of distributional, asymptotic behavior, Abelian theorems.*

1980 AMS SUBJECT CLASSIFICATION CODE. 44F12, 44A15.

1. REGULARLY VARYING FUNCTIONS

Throughout the paper, r will denote a positive continuous function on an interval (X, ∞) , $X \geq 0$, such that the limit

$$\lim_{t \rightarrow \infty} \frac{r(pt)}{r(t)}$$

exists for every $p > 0$. Such functions are called regularly varying functions (r.v.f.) at infinity and it is well known ([7]) that they are of the form $r(t) = t^a L(t)$ for some $a \in \mathbb{R}$ (called the order or index of r) and some slowly varying function (s.v.f.) L . This means that the function $L : (X, \infty) \rightarrow (0, \infty)$ is continuous and that

$$\lim_{t \rightarrow \infty} \frac{L(pt)}{L(t)} = 1$$

for every $p > 0$.

2. QUASIASYMPTOTIC BEHAVIOUR AT INFINITY RELATED TO r

The quasiasymptotic behaviour (q.a.b.) at infinity of tempered distributions with support in $[0, \infty)$ (denoted by S'_+) was defined by Zavjalov (see, for instance, [2]). In this paper we use a somewhat more general concept of q, a, b , related to a r.v.f. as defined and analysed in [10].

Definition 1. Let $T \in S'_+$ and r be some r.v.f. . The distribution T has q.a.b. at infinity related to r if there exists the limit in the sense of S' :

$$\lim_{t \rightarrow \infty} \frac{T(kt)}{r(k)} = g(t)$$

provided that $g \neq 0$.

If the order of r is a , then $g(t) = A f_{a+1}(t)$ for some $A \neq 0$ (from now on we take $A = 1$ for simplicity), where

$$f_{a+1}(t) = H(t)t^a/\Gamma(a+1) \text{ for } a \geq 0 \text{ and } f_{a+1}(t) = D^n f_{a+n+1}(t)$$

for $a < 0$ and $n+a > 0$, $n \in \mathbb{N}$. As usual, H is the characteristic function of the interval $(0, \infty)$, and D stands for the distributional derivative.

It is easy to see that a continuous function on $[0, \infty)$ having ordinary asymptotic behaviour of order $a > -1$ related to r has also q.a.b. of the same order and conversely. However for $a \leq -1$ this may not be true. This follows from the following

Structural Theorem. ([10]) A distribution $T \in S'_+$ has q.a.b. at infinity related to a r.v.f. r of order a iff there exist a natural number n , $n+a > 0$, and a continuous function F on \mathbb{R} such that

$$F = T * f_n \quad \text{and} \quad F(t) \sim \frac{1}{\Gamma(n+a+1)} t^n r(t) \quad \text{as } t \rightarrow \infty.$$

The proof of this important theorem is analogous to the one of Theorem I in [2], p. 373.

3. EQUIVALENCE AT INFINITY

The other "asymptotic behaviour" of distributions at infinity given in the following definition was used in [3], [1] and [6]; however, this notion goes back to Sebastiao e Silva ([8]).

Definition 2. A distribution $T \in S'_+$ is equivalent at infinity to $r(t) = t^a L(t)$, $a \in \mathbb{Z}_-$, if for some X' , $X' \geq X$, and some nonnegative integer n , $n+a > 0$, there exists a continuous function F on $[X', \infty)$ such that $T = D^n F$ on (X', ∞) and

$$F(t) \sim t^n r(t)/(a+1)(a+2)\dots(a+n) \quad (3.1)$$

in the ordinary sense as $t \rightarrow \infty$.

It seems to be of interest to compare these two asymptotics; for our purposes it is enough to prove

Lemma 1. Let $T \in S'_+$ be equivalent at infinity to $r(t) = t^a L(t)$ for $a > -1$. Then it has q.a.b. of order a related to r .

Proof. We can write $T = B + D^n F(t)$, where the supports of B and F are, respectively, in $[0, X']$ and $[X', \infty)$, $X' \geq 1$. Let us prove that

$$\lim_{k \rightarrow \infty} \frac{B(kt)}{k^a L(k)} \stackrel{S'}{\rightarrow} 0$$

In fact, for every $\epsilon > 0$ there exists a number $n_1 \in \mathbb{N}_0$ ($\mathbb{N}_0 = \mathbb{N} \cup \{0\}$) and a continuous function F_1 on \mathbb{R} such that $D^{n_1} F_1 = B$ and $\text{supp } F_1 \subset [-\epsilon, X'+\epsilon]$. For $\phi \in S$ we have

$$\begin{aligned} & \left\langle \frac{B(kt)}{k^a L(k)}, \phi(t) \right\rangle = \left\langle \frac{F_1(t)}{k^{n_1+a+1} L(k)}, (-1)^{n_1} \phi^{(n_1)}(t/k) \right\rangle = \\ & = \frac{(-1)^{n_1}}{k^{n_1+a+1} L(k)} \int_{-\epsilon}^{X'+\epsilon} F_1(t) \phi^{(n_1)}(t/k) dt \rightarrow 0 \end{aligned} \quad (3.1)$$

since $n_1+a+1 \geq a+1 > 0$. By supposition F satisfies (3.1), so by the Structural theorem it has q.a.b. of order a related to r .

4. STIELTJES TRANSFORM OF DISTRIBUTIONS

For the sake of completeness we rewrite the definition of the distributional Stieltjes transform given in [4]. Let $I'(z)$, $z \in C$, denote the subspace of distributions $t \in S'_+$ such that $T = D^n G$ for some $n \in N$ and some locally integrable function G on R with support in $[0, \infty)$ and

$$\int |G(t)| t^{-(z+n+1)} dt < \infty.$$

From now on we take $z \in R$ and $z > -1$, though a complex setting is also possible (see [4] or [1]). Obviously $I'(z) \subset S'_+$ and $I'(z_1) \subset I'(z_2)$ for $-1 < z_1 < z_2$.

Definition 3. The Stieltjes transform of index z of a distribution $T \in I'(z)$ is the complex valued function

$$S_z\{T\}(s) = \langle T(t), \frac{h(t)}{(t+s)^{z+1}} \rangle, \quad s \in C \setminus (-\infty, 0], \quad (4.1)$$

where h is an infinitely differentiable function on R such that $h(t) = 1$ in some neighbourhood of $[0, \infty)$ and $h(t) = 0$ in some interval $(-\infty, -\varepsilon)$, $\varepsilon > 0$.

It is easy to see that (4.1) does not depend on the function h , so it is usually omitted. It is proved in [5] that $S_z\{T\}(s)$ is a holomorphic function of the complex variable s in the domain $C \setminus (-\infty, 0]$ provided that $T \in I'(z)$. We shall need the following equality ([5], p. 140)

$$S_{z+n}\{T\}(s) = \frac{1}{(z+1)(z+2)\dots(z+n)} S_z\{D^n T\}(s) \quad (4.2)$$

for $T \in I'(z)$ and $n \in N$. Observe that $T \in I'(z)$ implies $T \in I'(z+n)$ and $D^n T \in I'(z)$.

5. ABELIAN THEOREMS

The initial value type Abelian theorems for the distributional Stieltjes transform seem to have a satisfactory form. So, we prove only final value type ones. We use first the following result from [6]:

Theorem 1. Let us suppose that $T \in I'(z)$ is equivalent at infinity to a regularly varying function $r(t) = t^a L(t)$ of order $a > -1$. Then

$$S_z\{T\}(s) \sim B(a+1, z-a) L(s) s^{a-z} \text{ as } s \rightarrow \infty, \quad s \in R, \quad (5.1)$$

provided that $z > a > -1$.

As usual, $B(p, q)$ stands for the beta function. In view of Lemma 1 we see that this Theorem can be rewritten as

Theorem 1'. Let us suppose that $T \in S'_+$ has q.a.b. of order $a > -1$ related to the r.v.f. $r(t) = t^a L(t)$. Then (5.1) holds if $z > a > -1$.

If T in these two theorems is a continuous function on $[0, \infty)$, then $T(t) \sim t^a L(t)$ as $t \rightarrow \infty$ in the ordinary sense. Essentially, we need such

a "functional" (i.e. not "distributional") version of them in the following

Abelian Theorem. Let $T \in S_+^1$ have q.a.b. of order a related to a r.v.f. $r(t) = t^a L(t)$. Then

i) $T \in I'(z)$ for $z > \max(-1, a)$

and

ii) $S_z\{T\}(s) \sim \frac{\Gamma(z-a)}{\Gamma(z+1)} L(s) s^{a-z}$ as $s \rightarrow \infty$,
staying on the real line.

Remark. Such a statement was proved in [3] for $r(t) = t^a$, $a > -1$ and in [4] for $r(t) = t^a \log^j t$, $a > -1$. Further on, r.v.f. were used in [6] (again for $a > -1$). In all these papers the equivalence at infinity was used. The q.a.b. was used in [9] for $r(t) = a^a$ (a - arbitrary real number) and now for any r.v.f. In [1] the results from [3] are given in a complex setting; it might be of interest to prove an analogous statement for our Abelian theorem.

Proof of the Abelian theorem. Part i) follows from the Structural theorem and the estimate $L(t) \leq C_\varepsilon t^\varepsilon$ for $t \geq t_0 = t_0(\varepsilon)$ (ε in $(0,1)$). For ii), we take $n > -a$ and F as in the structural theorem; then

$$F(t) \sim C_n t^{n+a} L(t) \text{ as } t \rightarrow \infty$$

for $C_n = 1/\Gamma(n+a+1)$. By Theorem 1' we get

$$S_{z+n}\{F\}(s) \sim C_n B(n+a+1, z+n-(n+a)) L(s) s^{a-z} \text{ as } s \rightarrow \infty,$$

and from (4.2) we have

$$\begin{aligned} S_z\{T\}(s) &\sim (z+1)(z+2)\dots(z+n) S_{z+n}\{F\}(s), \text{ so} \\ S_z\{T\}(s) &\sim C_n \frac{\Gamma(n+a+1) \Gamma(z-a)}{\Gamma(z+1)} L(s) s^{a-z}. \end{aligned} \tag{5.2}$$

This gives the statement ii).

Example. The equivalence at infinity with the distribution

$$T = A(a, j) F_p(t^a \log_+^j t), \quad a \in \mathbb{R}, j \in \mathbb{N}_0 \tag{5.3}$$

for appropriate constant $A(a, j)$ was analysed in [4]: F_p stands for the finite part. Obviously, T is equivalent at infinity to $t^a \log^j t$ for $a \notin \mathbb{Z}_+$; we take $A(a, j) = 1$ then. On the other hand, T has q.a.b. of order a related to $t^a \log^j t$ for $a \notin \mathbb{Z}_-$ and related to $A(a, j)t^a \log^{j+1} t$ for $a \in \mathbb{Z}_-$; we take $A(a, j) = (-1)^{-a-1}/((-a-1)!(j+1))$ then. Computing the Stieltjes transform of T we see that it behaves at infinity as the Abelian theorem predicts (see [4], formulae (2.3) and (2.4)).

Now let $-2 < a < -1$. Then the distribution $S = T + \delta$ has q.a.b. of order -1 related to $1/t$ and is equivalent at infinity with $t^a \log^j t$. But for $z > -1$

$$S_z\{S\} = S_z\{T\} + S_z\{\delta\} \sim C_{a,j} \log^{j+1} s s^{a-z} + s^{-(z+1)} \sim s^{-(z+1)}$$

when $s \rightarrow \infty$. This trivial example (which can be generalized easily) shows

again that q.a.b. is more appropriate for final value type Abelian theorems for Stieltjes transformation than equivalence at infinity, though the latter seems more "natural".

REFERENCES

1. CARMICHAEL, R.D. and MILTON, E.O. Abelian theorems for the distributional Stieltjes transform, J. Math. Anal. Appl., Vol. 72, No. 1, (1979), 195 - 205.
2. DROŽŽINOV, JU.N. and ZAVIJALOV, B.I. Quasiasymptotic behaviour of generalized functions and Tauberian theorems in complex domain (in Russian), Mat. Sbornik, Tom 102 (144), No. 3, (1977), 372 - 390.
3. LAVOINE, J. and MISRA, O.P. Théorèmes abéliens pour la transformation de Stieltjes des distributions, C.R. Acad. Sci. Paris 270 (1974), 99 - 102.
4. LAVOINE, J. and MISRA, O.P. Abelian theorems for the distributional Stieltjes transformation, Math. Proc. Phil. Soc. 86 (1979), 287 - 293.
5. LAVOINE, J. and MISRA, O.P. Sur la transformation de Stieltjes des distributions et son inversion au moyen de la transformation de Laplace, C.R. Acad. Sci. Paris, 290 (1980), 139 - 142.
6. MARIĆ, V., SKENDŽIĆ, M. and TAKAČI, A. On Stieltjes transform of distributions behaving as regularly varying functions, (to appear).
7. SINETA, E. Regularly varying functions, Springer Verlag Berlin, Heidelberg, New York, Lecture Notes in Math. 508 (1976).
8. SILVA, J.S.e. Integrals and order of growth of distributions, Inst. Cubelkijan de Ciencia, Lisboa 35 (1964), 7124.
9. TAKAČI, A. A note on the distributional Stieltjes transformation, Math. Proc. Phil. Soc., 94 (1983), 523 - 527.
10. VLADIMIROV, V.S. and ZAVIJALOV, B.I. Tauberian theorems in the quantum field theory, (in Russian), Itogi Nauki i Tehniki, Serija Sovremennie problemi matematiki, Tom 15 (1980), Moskva.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Center for Applied Dynamics Research, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk