

## THE LEBESGUE INTEGRAL AS A RIEMANN INTEGRAL

ENRIQUE A. GONZALEZ-VELASCO

Department of Mathematics  
University of Lowell  
Lowell, Massachusetts 01854 U.S.A.

(Received November 21, 1985 and in revised form December 30, 1986)

**ABSTRACT.** The object of this paper is to develop a very direct theory of the Lebesgue integral that is easily accessible to any audience familiar with the Riemann integral.

**KEY WORDS AND PHRASES.** *Lebesgue integration, Riemann integration.*

1980 AMS SUBJECT CLASSIFICATION CODE. 2801.

### 1. INTRODUCTION.

The problem of making the theory of Lebesgue integration accessible to a wide audience, including students of mathematics, science and engineering, is not easily solved. And yet, the Lebesgue integral is an indispensable tool in many branches of mathematics and its applications, such as Fourier analysis and signal processing.

The inaccessibility of the theory is mainly due to the fact that the usual presentations, both those that are and those that are not based on measure theory, are lengthy and indirect and rely on a number of new basic concepts that must be assimilated first. For instance, in Royden [1] the general definition of the Lebesgue integral is preceded by definitions for simple functions, then for bounded functions over a set of finite measure and then for nonnegative functions, and in each particular case some amount of theory is elaborated. In Apostol [2] the general definition is preceded by definitions for step functions and then for upper functions, and again a certain amount of theory is developed in each particular case. Furthermore, these classic presentations represent a new start, with all knowledge of Riemann integration to be abandoned in favor of the new Lebesgue theory. To avoid this, McShane [3] has recently developed a unified theory of Riemann and Lebesgue integration that is, hopefully, elementary enough for presentation to a wide audience. While sharing McShane's concern and agreeing that he succeeds in constructing an appealing unified theory, this author believes that, regarding simplicity and accessibility, the answer lies elsewhere.

There is an entirely different approach to the definition of the Lebesgue integral, due to Mikusiński [4], which is very direct and does not need the previous development of either measure theory or the Riemann integral. It is based, instead, on a series representation of the function to be integrated, but, since this series

representation need not be unique, it requires a somewhat lengthy proof of consistency. In addition, it is not readily apparent how the definition of the Lebesgue integral in any of the approaches mentioned so far can be immediately used for the quick computation of simple integrals.

Our task is, then, to construct a simple theory that builds on concepts familiar to any audience and that is based on a very direct and general definition, capable of dealing with the computation of simple integrals. To this end, the best course of action is to define the Lebesgue integral as a (possibly improper) Riemann integral. This is not only brief but allows for complete generality at once and, moreover, we know how to compute Riemann integrals (there is no need to develop a theory of improper Riemann integrals as a prerequisite because, beyond the definition, all that is needed is the fact that such an integral is a linear and increasing function of the integrand). Such a theory is presented in Rey Pastor [5], but only for functions defined on a set of finite measure and his development makes use of the Riemann-Stieltjes integral. The main object of this paper is to reconstruct this theory from the beginning (with one minor modification) for functions defined over sets of arbitrary measure and, for additional simplicity, avoiding all use of the Riemann-Stieltjes integral. This has necessitated a complete set of new proofs and a collection of new auxiliary results (Propositions 2, 3 and 4 and Lemmas 3, 4 and 5 below). In addition, we have included the Levi monotone convergence theorem and have adopted an entirely different approach to the Lebesgue dominated convergence theorem. Instead of basing its proof on the additivity of the integral (not an easy fact to establish), our approach highlights the role of uniform convergence. Lemmas 3 and 4 show that the contribution of the dominated convergence is just to enable the uniform convergence to do the job via Egorov's theorem and Proposition 4.

## 2. THE LEBESGUE MEASURE AND MEASURABLE FUNCTIONS.

For easy reference, and to specify how much Lebesgue measure theory is necessary for our presentation, we collect in this section those results that will be used in the rest of the article. For proofs see [1] or [6]. By a rectangle in  $\mathbb{R}^n$  we shall mean the product space of  $n$  bounded intervals, open, closed, or neither. The volume of such a rectangle  $R$ , denoted by  $v(R)$ , is the product of the lengths of its component intervals.

DEFINITION 1. The outer measure of a set  $E \subset \mathbb{R}^n$  is

$$m(E) \stackrel{\text{def}}{=} \inf \sum v(R_i) \quad (2.1)$$

where the infimum is taken over all finite or countable collections of open rectangles  $\{R_i\}$  such that  $E \subset \bigcup R_i$ .

It turns out that it is not always true that the outer measure of a finite union of disjoint sets is the sum of their outer measures. But this will always be true if we restrict ourselves to sets of the following type.

DEFINITION 2. A set  $E \subset \mathbb{R}^n$  is called measurable if and only if for any set  $S \subset \mathbb{R}^n$   $m(S) = m(S \cap E) + m(S \cap E^c)$ , where  $E^c$  is the complement of  $E$  in  $\mathbb{R}^n$ . If  $E$  is measurable  $m(E)$  is called the Lebesgue measure of  $E$ .

Notice that  $\mathbb{R}^n$  and the empty set are measurable with Lebesgue measures  $\infty$

and zero respectively. It is also clear that a set is measurable if and only if its complement is measurable and that if  $E$  and  $F$  are measurable with  $F \subset E$  then  $m(F) \leq m(E)$ .

THEOREM 1.

- (1) If  $E$  is a finite or countable set then  $m(E) = 0$ .
- (2) If  $\{E_i\}$  is a finite or countable collection of measurable sets then their union  $\bigcup E_i$  is measurable and  $m(\bigcup E_i) \leq \sum m(E_i)$ . If, in addition, the  $E_i$  are disjoint then equality holds.
- (3) If  $\{E_i\}$  is a finite or countable collection of measurable sets then their intersection  $\bigcap E_i$  is measurable. If, in addition,  $m(E_i) < \infty$  and  $E_{i+1} \subset E_i$  for all  $i$  then  $m(E_i) \rightarrow m(\bigcap E_i)$  as  $i \rightarrow \infty$ .
- (4) If  $E$  and  $F$  are measurable then  $E - F \stackrel{\text{def}}{=} \{x \in E : x \notin F\}$  is measurable. If, in addition,  $m(E) < \infty$  and  $F \subset E$  then  $m(E - F) = m(E) - m(F)$ .
- (5) Every half space in  $\mathbb{R}^n$  is measurable.
- (6) Every rectangle in  $\mathbb{R}^n$  is measurable and its Lebesgue measure is its volume.
- (7) Open and closed sets are measurable.

DEFINITION 3. A function  $f: E \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is measurable if and only if  $E$  is measurable and for any  $y \in \mathbb{R}$  the set  $\{x \in E : f(x) > y\}$  is measurable.

THEOREM 2. The following statements are equivalent:

- (1) For each  $y \in \mathbb{R}$  the set  $\{x \in E : f(x) > y\}$  is measurable.
- (2) For each  $y \in \mathbb{R}$  the set  $\{x \in E : f(x) \geq y\}$  is measurable.
- (3) For each  $y \in \mathbb{R}$  the set  $\{x \in E : f(x) < y\}$  is measurable.
- (4) For each  $y \in \mathbb{R}$  the set  $\{x \in E : f(x) \leq y\}$  is measurable.

EXERCISE 1. If  $E$  is measurable and  $f:E \rightarrow \mathbb{R}$  is continuous then  $f$  is measurable.

THEOREM 3. Let  $E \subset \mathbb{R}^n$  be a measurable set.

- (1) If  $f, g: E \rightarrow \mathbb{R}$  are measurable then  $f + g$  is measurable.
- (2) If  $\{f_N\}$  is a sequence of measurable functions on  $E$  and if  $f_N \rightarrow f$  on  $E$  then  $f$  is measurable.

THEOREM 4. (Egorov's theorem) Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $\{f_N\}$  be a sequence of measurable functions on  $E$  such that  $f_N \rightarrow f$  on  $E$ . For any  $\delta > 0$  there is a subset  $G$  of  $E$  with  $m(G) < \delta$  and such that  $f_N \rightarrow f$  uniformly on  $E - G$ .

### 3. THE LEBESGUE INTEGRAL.

The Lebesgue approach to define the integral of a function  $f: E \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is to partition its range, not its domain as in the Riemann theory. To be specific, consider the case of a measurable set  $E$  with  $m(E) < \infty$  and a measurable function  $f: E \rightarrow \mathbb{R}$  whose values are between zero and  $M > 0$ . If  $0 = y_0 < y_1 < \dots < y_k = M$  is a partition of  $[0, M]$  and if we define  $S_i = \{x \in E : f(x) > y_i\}$  for  $i = 0, \dots, k$ , then  $f$  is said to be integrable on  $E$  if and only if

$$\sup \sum_{i=1}^k m(S_i)(y_i - y_{i-1}) = \inf \sum_{i=1}^k m(S_{i-1})(y_i - y_{i-1}), \quad (3.1)$$

where the supremum and infimum are taken over all partitions of  $[0, M]$ , and this

common value will be denoted by  $\int_E f$ .

EXERCISE 2. Define a set  $E_i = \{x \in E : y_{i-1} < f(x) \leq y_i\}$  for each  $i$ . Prove that

$$\sum_{i=1}^k m(E_i)y_{i-1} = \sum_{i=1}^k m(S_i)(y_i - y_{i-1}) \quad (3.2)$$

and

$$\sum_{i=1}^k m(E_i)y_i = \sum_{i=1}^k m(S_{i-1})(y_i - y_{i-1}). \quad (3.3)$$

Now, if we define a function  $\mu_f : [0, M] \rightarrow \mathbb{R}$  by  $\mu_f(y) = m\{x \in E : f(x) > y\}$  then  $\mu_f$  is clearly bounded and decreasing and, thus, Riemann integrable on  $[0, M]$ .

PROPOSITION 1. Let  $E \subset \mathbb{R}^n$  be a measurable set with  $m(E) < \infty$ . Every bounded, nonnegative measurable function  $f : E \rightarrow \mathbb{R}$  is integrable on  $E$  and

$$\int_E f = R \int_0^M \mu_f, \quad (3.4)$$

where the  $R$  indicates Riemann integral.

PROOF. Notice that  $m(S_i) = \mu_f(y_i)$  and then, since  $\mu_f$  is decreasing,

$$\sum_{i=1}^k m(S_i)(y_i - y_{i-1}) \quad \text{and} \quad \sum_{i=1}^k m(S_{i-1})(y_i - y_{i-1}) \quad (3.5)$$

are, respectively, the lower and upper Riemann sums of  $\mu_f$  with respect to the partition  $0 = y_0 < y_1 < \dots < y_k = M$  of  $[0, M]$ . The result follows because  $\mu_f$  is Riemann integrable on  $[0, M]$ , Q.E.D.

If  $f$  is nonpositive instead of nonnegative we would like its integral on  $E$  to equal the opposite of the integral of  $-f$ , that is, we want

$$\int_E f = -R \int_0^M \mu_{-f} = R \int_M^0 \mu_{-f} = R \int_{-M}^0 -\mu_{-f}(-y) dy. \quad (3.6)$$

This suggests that we define  $\mu_f(y) = -\mu_{-f}(-y)$  if  $y < 0$ , and then the entire preceding discussion motivates the following general definitions.

DEFINITION 4. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $f : E \rightarrow \mathbb{R}$  be a measurable function. The measure function of  $f$  on  $E$  is the function  $\mu_f$  defined by

$$\mu_f(y) = \begin{cases} m\{x \in E : f(x) > y\} & \text{if } y > 0 \\ -m\{x \in E : f(x) < y\} & \text{if } y < 0 \end{cases} \quad (3.7)$$

Notice that  $\mu_f \leq 0$  on  $(-\infty, 0)$  and  $\mu_f \geq 0$  on  $(0, \infty)$ , and that  $y_1 \leq y_2 \Rightarrow \mu_f(y_1) \geq \mu_f(y_2)$  on each of these intervals. The measure function can have infinite values if  $m(E) = \infty$ , but, if it does not, it is Riemann integrable on every bounded subinterval of  $(-\infty, 0)$  or  $(0, \infty)$ , and then the improper integrals

$$R \int_{-\infty}^0 \mu_f \quad \text{and} \quad R \int_0^{\infty} \mu_f \quad (3.8)$$

exist (they may be infinite). For convenience, if  $\mu_f$  has infinite values on  $(-\infty, 0)$  or on  $(0, \infty)$  we define

$$R \int_{-\infty}^0 \mu_f = -\infty \quad \text{or} \quad R \int_0^{\infty} \mu_f = \infty. \quad (3.9)$$

DEFINITION 5. Let  $E \subset \mathbb{R}^n$  be a measurable set,  $f:E \rightarrow \mathbb{R}$  a measurable function, and  $\mu_f$  the measure function of  $f$  on  $E$ . Then the (Lebesgue) integral of  $f$  on  $E$  is defined to be

$$\int_E f = R \int_{-\infty}^{\infty} \mu_f \quad (3.10)$$

if the right-hand side exists (that is if the two improper Riemann integrals over  $(-\infty, 0)$  and  $(0, \infty)$  exist). If, in addition, the right-hand side is finite then  $f$  is said to be (Lebesgue) integrable on  $E$  and we write  $f \in L(E)$ .

EXAMPLE 1. If  $E = \{x \in \mathbb{R} : 0 < x < 1\}$  and  $f(x) = 1/\sqrt{x}$  we have  $\mu_f(y) = 0$  if  $y < 0$ ,  $\mu_f(y) = 1$  if  $0 < y < 1$  and  $\mu_f(y) = y^{-2}$  if  $y \geq 1$ , and then

$$\int_E f = R \int_0^1 dy + R \int_1^{\infty} y^{-2} dy = 2. \quad (3.11)$$

EXAMPLE 2. If  $E = \{x \in \mathbb{R} : 0 < x < 1\}$  and  $f(x) = 1/x$  then  $\mu_f(y) = 0$  if  $y < 0$ ,  $\mu_f(y) = 1$  if  $0 < y < 1$  and  $\mu_f(y) = 1/y$  if  $y \geq 1$ . We have

$$\int_E f = R \int_0^1 dy + R \int_1^{\infty} \frac{1}{y} dy = \infty. \quad (3.12)$$

Examples 1 and 2 show that the product of integrable functions need not be integrable.

EXAMPLE 3. If  $E = \{x \in \mathbb{R} : 0 < x < 1\}$  and  $f(x) = 1$  when  $x$  is rational and  $f(x) = 0$  when  $x$  is irrational, then  $\mu_f(y) = 0$  for all  $y \neq 0$  and  $\int_E f = 0$ .

EXERCISE 3. Show that if  $f \geq 0$  and  $\int_E f = 0$  then  $m\{x \in E : f(x) \neq 0\} = 0$ .

#### 4. BASIC PROPERTIES OF THE LEBESGUE INTEGRAL.

First we establish the basic properties of the measure function. We shall write  $\mu_f^E$  instead of  $\mu_f$  when we need to specify the domain  $E$  of  $f$ .

LEMMA 1. Let  $E \subset \mathbb{R}^n$  be a measurable set, let  $f, g: E \rightarrow \mathbb{R}$  be measurable functions and let  $c \in \mathbb{R}$ . Then

$$(1) \quad f \leq g \Rightarrow \mu_f \leq \mu_g.$$

$$(2) \quad cf \text{ is measurable and } \mu_{cf}(y) = \text{sgn}(c)\mu_f(y/c) \text{ for all } y \neq 0.$$

PROOF.

(1) If  $y > 0$ ,  $\{x \in E : f(x) > y\} \subset \{x \in E : g(x) > y\} \Rightarrow \mu_f(y) \leq \mu_g(y)$ . If  $y < 0$ ,  $\{x \in E : g(x) < y\} \subset \{x \in E : f(x) < y\} \Rightarrow -\mu_g(y) \leq -\mu_f(y) \Rightarrow \mu_f(y) \leq \mu_g(y)$ .

(2) We shall consider the case  $c < 0$ ,  $y > 0$ , and the others are similar.

$$\begin{aligned} \mu_{cf}(y) &= m\{x \in E : cf(x) > y\} \\ &= m\{x \in E : f(x) < y/c\} \\ &= -\mu_f(y/c) \\ &= \text{sgn}(c)\mu_f(y/c) \end{aligned} \quad (4.1)$$

Q.E.D.

LEMMA 2. If  $\{E_i\}$  is a finite or countable collection of disjoint measurable sets in  $\mathbb{R}^n$ , if  $E = \bigcup E_i$  and if  $f:E \rightarrow \mathbb{R}$  is a measurable function, then  $\mu_f^E = \sum \mu_f^{E_i}$ .

PROOF. If  $y > 0$ ,  $\mu_f^E(y) = m\{x \in E : f(x) > y\} = m(\bigcup\{x \in E_i : f(x) > y\}) = \sum m\{x \in E_i : f(x) > y\} = \sum \mu_f^{E_i}$ . Similarly if  $y < 0$ , Q.E.D.

COROLLARY 1. If  $E, F \subset \mathbb{R}^n$  are measurable sets with  $F \subset E$  and if  $f: E \rightarrow \mathbb{R}$  is a measurable function, then  $\mu_f^F \leq \mu_f^E$  on  $(0, \infty)$  and  $\mu_f^F \geq \mu_f^E$  on  $(-\infty, 0)$ .

PROOF.  $E = F \cup (E - F) \Rightarrow \mu_f^E = \mu_f^F + \mu_f^{E-F}$ . Then  $\mu_f^{E-F} \geq 0$  on  $(0, \infty) \Rightarrow \mu_f^F \leq \mu_f^E$  on  $(0, \infty)$  and  $\mu_f^{E-F} \leq 0$  on  $(-\infty, 0) \Rightarrow \mu_f^F \geq \mu_f^E$  on  $(-\infty, 0)$ , Q.E.D.

THEOREM 5. Let  $E, F \subset \mathbb{R}^n$  be measurable sets, let  $f, g: E \rightarrow \mathbb{R}$  be measurable functions and let  $c \in \mathbb{R}$ . Then, if the integrals below exist,

(1)  $f \leq g \Rightarrow \int_E f \leq \int_E g$  and if, in addition,  $f \geq 0$  then  $g \in L(E) \Rightarrow f \in L(E)$ .

(2)  $\int_E cf = c \int_E f$  and then  $f \in L(E) \Rightarrow cf \in L(E)$ .

(3) If  $F \subset E$  then  $f \in L(E) \Rightarrow f \in L(F)$  and if, in addition,  $f \geq 0$  then  $\int_F f \leq \int_E f$ .

(4) If  $c > 0$  and if  $f^c \stackrel{\text{def}}{=} \min\{f, c\}$  then  $\int_E f^c \rightarrow \int_E f$  as  $c \rightarrow \infty$ .

PROOF. (1) is obvious from Lemma 1(1).

(2) Assume that  $c < 0$  (proceed analogously if  $c > 0$ ). By Lemma 1(2),

$$R \int_0^b \mu_{cf}(y) dy = R \int_0^b -\mu_f(y/c) dy = c R \int_{b/c}^0 \mu_f(u) du \quad (4.2)$$

and taking the limit as  $b \rightarrow \infty$ ,

$$R \int_0^\infty \mu_{cf} dy = c R \int_{-\infty}^0 \mu_f du. \quad (4.3)$$

Similarly for the integral from  $-\infty$  to 0. Adding the two proves (2).

(3) By Corollary 1,

$$R \int_{-\infty}^0 \mu_f^E \leq R \int_{-\infty}^0 \mu_f^F \leq 0 \leq R \int_0^\infty \mu_f^F \leq R \int_0^\infty \mu_f^E. \quad (4.4)$$

If the outer integrals are finite, so are the inner integrals. Thus,  $f \in L(E) \Rightarrow f \in L(E)$ . The last assertion follows from the last two inequalities on the right.

$$(4) \quad \int_E f^c = R \int_{-\infty}^c \mu_f^c = R \int_{-\infty}^c \mu_f \rightarrow R \int_{-\infty}^\infty \mu_f = \int_E f \quad (4.5)$$

as  $c \rightarrow \infty$ , Q.E.D.

THEOREM 6. Let  $E_1, \dots, E_N$  be a finite collection of disjoint measurable sets in  $\mathbb{R}^n$ , let  $E = \bigcup E_i$  and let  $f: E \rightarrow \mathbb{R}$  be a measurable function. Then

$$\int_E f = \sum \int_{E_i} f. \quad (4.6)$$

Furthermore, if  $f$  has a constant value  $c_i$  on each  $E_i$ , then

$$\int_E f = \sum c_i m(E_i). \quad (4.7)$$

PROOF. The first assertion follows from Lemma 2 and Definition 5. Now, if  $c_i > 0$  then  $\mu_f^{E_i}(y) = m(E_i)$  when  $0 < y < c_i$  (here  $m(E_i)$  can be  $\infty$ ) and  $\mu_f^{E_i}(y) = 0$  otherwise. Then

$$\int_{E_i} f = R \int_0^{c_i} \mu_f^{E_i} = c_i m(E_i). \quad (4.8)$$

Similarly if  $c_i < 0$ , Q.E.D.

EXERCISE 4. If there are constants  $a, b \in \mathbb{R}$  such that  $a \leq f(x) \leq b$  for all  $x \in E$ , then  $a m(E) \leq \int_E f \leq b m(E)$  and  $m(E) < \infty \Rightarrow f \in L(E)$ .

EXERCISE 5. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $f, g: E \rightarrow \mathbb{R}$  be measurable functions. If  $f = g$  on  $E$  except, possibly, on a subset of  $E$  of measure zero, then  $\int_E f = \int_E g$ .

THEOREM 7. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $f: E \rightarrow \mathbb{R}$  be a measurable function. If  $\int_E f$  exists, then

$$|\int_E f| \leq \int_E |f| \quad (4.9)$$

Also,  $f \in L(E) \Rightarrow |f| \in L(E)$ .

PROOF. If  $y > 0$  the identity  $\{x \in E : |f(x)| > y\} = \{x \in E : f(x) > y\} \cup \{x \in E : f(x) < -y\}$  shows that  $|f|$  is measurable and that  $\mu_{|f|}(y) = \mu_f(y) - \mu_f(-y)$ . Now,

$$R \int_0^\infty \mu_f(-y) dy = -R \int_0^{-\infty} \mu_f(u) du = R \int_{-\infty}^0 \mu_f(y) dy, \quad (4.10)$$

so that

$$R \int_0^\infty \mu_{|f|} = R \int_0^\infty (\mu_f(y) - \mu_f(-y)) dy = R \int_0^\infty \mu_f - R \int_{-\infty}^0 \mu_f. \quad (4.11)$$

The right-hand side is always well defined as a real number or  $\infty$  since  $\mu_f \leq 0$  on  $(-\infty, 0)$  and  $\mu_f \geq 0$  on  $(0, \infty)$ , and it is finite if and only if both integrals are finite, that is  $f \in L(E) \Rightarrow |f| \in L(E)$ . In either case, the inequality  $|a + b| \leq a - b$  holds when  $a \geq 0$  and  $b \leq 0$ , when  $a \geq 0$  and  $b = -\infty$ , and when  $a = \infty$  and  $b \leq 0$ . Thus,

$$|\int_E f| = |R \int_0^\infty \mu_f + R \int_{-\infty}^0 \mu_f| \leq \int_E |f|, \quad (4.12)$$

Q.E.D.

EXERCISE 6. Let  $R$  be a rectangle in  $\mathbb{R}^n$ . If  $f: R \rightarrow \mathbb{R}$  is Riemann integrable then  $f \in L(E)$  and  $\int_E f = R \int_R f$ .

##### 5. THE ADDITIVITY OF THE LEBESGUE INTEGRAL.

This is one of the weak points of the Lebesgue theory and it will not be straightforward to establish this fact. We start by considering two particular cases: those of piecewise continuous functions and nonnegative bounded functions.

PROPOSITION 2. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $f, g: E \rightarrow \mathbb{R}$  be measurable functions each of which has a finite number of values. Then

$$\int_E (f + g) = \int_E f + \int_E g. \quad (5.1)$$

PROOF. If  $f$  has only a finite number of values  $c_1, \dots, c_N$  on disjoint subsets  $F_1, \dots, F_N$  of  $E$  and if  $g$  has only a finite number of values  $c_1, \dots, c_K$  on disjoint subsets  $G_1, \dots, G_K$  of  $E$ , then  $f + g$  is the function whose values are  $c_i + c_j$  on  $F_i \cap G_j$ . Noticing that  $F_i$  is the disjoint union  $\sum_{j=1}^K F_i \cap G_j$  and using Theorem 6 we have

$$\int_E (f + g) = \sum_{i=1}^N \sum_{j=1}^K (c_i + c_j) m(F_i \cap G_j)$$

$$\begin{aligned}
&= \sum_{i=1}^N c_i \sum_{j=1}^K m(F_i \cap G_j) + \sum_{j=1}^K c_j \sum_{i=1}^N m(G_j \cap F_i) \\
&= \sum_{i=1}^N c_i m(F_i) + \sum_{j=1}^K c_j m(G_j) \\
&= \int_E f + \int_E g,
\end{aligned} \tag{5.2}$$

Q.E.D.

PROPOSITION 3. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $f, g : E \rightarrow \mathbb{R}$  be non-negative, bounded, measurable functions. Then

$$\int_E (f + g) = \int_E f + \int_E g. \tag{5.3}$$

PROOF. Let  $f$  have values in  $[0, M]$ ,  $M > 0$ , and let  $0 = y_0 < y_1 < \dots < y_k = M$  be a partition of  $[0, M]$ . Then define  $F_i = \{x \in E : f(x) > y_i\}$  and a function  $s_f \leq f$  by  $s_f(x) = y_{i-1}$  if  $x \in F_{i-1} - F_i$ ,  $i = 1, \dots, k$ . Noticing that  $y_0 = m(F_k) = 0$  and using Theorem 6 we have

$$\begin{aligned}
\int_E s_f &= \sum_{i=1}^k y_{i-1} m(F_{i-1} - F_i) \\
&= \sum_{i=1}^k y_{i-1} (m(F_{i-1}) - m(F_i)) \\
&= \sum_{i=1}^k (y_i - y_{i-1}) m(F_i) \\
&= \sum_{i=1}^k \mu_f(y_i)(y_i - y_{i-1}) \\
&\leq \int_E f,
\end{aligned} \tag{5.4}$$

and, as the norm of the partition approaches zero,

$$\int_E s_f \rightarrow R \int_0^M \mu_f = \int_E f. \tag{5.5}$$

If  $s_g$  is defined for  $g$  as  $s_f$  was defined for  $f$ , applying Proposition 2 to  $s_f$  and  $s_g$ , observing that  $s_f + s_g \leq f + g$ , using Theorem 5(1), and letting the norm of the partition approach zero, we obtain

$$\int_E (f + g) \geq \int_E (s_f + s_g) = \int_E s_f + \int_E s_g \rightarrow \int_E f + \int_E g. \tag{5.6}$$

The first integral above is well defined because  $f + g$  is measurable, nonnegative and bounded. Similarly, if we define a function  $S_f \geq f$  by  $S_f(x) = y_i$  if  $x \in F_{i-1} - F_i$  and if  $S_g$  is defined for  $g$  as  $S_f$  is for  $f$ , a similar argument would yield

$$\int_E (f + g) \leq \int_E f + \int_E g, \tag{5.7}$$

and the conclusion follows from the last two inequalities, Q.E.D.

THEOREM 8. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $f, g \in L(E)$ . Then  $f + g \in L(E)$  and

$$\int_E (f + g) = \int_E f + \int_E g. \tag{5.8}$$

PROOF. Assume first that  $f$  and  $g$  are nonnegative and for  $M > 0$  define  $f^M = \min\{f, M\}$  and, similarly,  $g^M$  and  $(f + g)^M$ . Since  $f^M + g^M \leq f + g$ , Proposition 3 gives

$$\int_E f^M + \int_E g^M = \int_E (f^M + g^M) \leq \int_E (f + g) \quad (5.9)$$

and then Theorem 5(4) implies that

$$\int_E f^M + \int_E g^M \rightarrow \int_E f + \int_E g \leq \int_E (f + g) . \quad (5.10)$$

as  $M \rightarrow \infty$ . It is not difficult to see that  $(f + g)^M \leq f^M + g^M$ , and then

$$\int_E (f + g)^M \leq \int_E (f^M + g^M) = \int_E f^M + \int_E g^M \leq \int_E f + \int_E g . \quad (5.11)$$

Hence

$$\int_E (f + g)^M \rightarrow \int_E (f + g) \leq \int_E f + \int_E g . \quad (5.12)$$

as  $M \rightarrow \infty$ . It follows that

$$\int_E (f + g) = \int_E f + \int_E g , \quad (5.13)$$

and, since the right-hand side is finite,  $f + g \in L(E)$ .

If, instead,  $f \geq 0$  and  $g \leq 0$ , define  $E^+ = \{x \in E : f + g \geq 0\}$  and  $E^- = \{x \in E : f + g < 0\}$ . Since  $f + g$ ,  $-g$  and  $f = f + g + (-g)$  are nonnegative on  $E^+$ , the preceding proof shows that

$$\int_{E^+} f = \int_{E^+} (f + g) + \int_{E^+} (-g) = \int_{E^+} (f + g) - \int_{E^+} g . \quad (5.14)$$

Similarly,  $-(f + g)$ ,  $f$  and  $-g = -(f + g) + f$  are nonnegative on  $E^-$  and

$$\int_{E^-} (-g) = - \int_{E^-} (f + g) + \int_{E^-} f . \quad (5.15)$$

Subtracting the last equation from the previous one and using Theorem 6 proves the desired result in this case.

The two cases proved above imply the truth of the assertion in the general case after decomposing  $E$  into the union of four disjoint subsets, on each of which  $f$  and  $g$  do not change sign, Q.E.D.

## 6. THE CONVERGENCE THEOREMS.

In this section we study the validity of the limit  $\int_E f_N \rightarrow \int_E f$  when  $f_N \rightarrow f$ . In the case of Riemann integration the uniform convergence of this sequence is a sufficient condition. We start by proving a similar result for the Lebesgue integral.

**PROPOSITION 4.** Let  $S \subset \mathbb{R}^n$  be a measurable set with  $m(S) < \infty$  and let  $\{f_N\}$  be a sequence of measurable functions such that  $f_N \in L(S)$  and  $f_N \rightarrow f \in L(S)$  uniformly on  $S$ . Then

$$\int_S f_N \rightarrow \int_S f . \quad (6.1)$$

PROOF. Given  $\epsilon > 0$  there is a  $K \in \mathbb{Z}^+$  such that  $f - \epsilon \leq f_N \leq f + \epsilon$ , and

then  $\mu_{f-\epsilon} \leq \mu_{f_N} \leq \mu_{f+\epsilon}$ , for  $N > K$ .

For  $y > 0$  and  $N > K$ ,

$$\begin{aligned}
 \mu_f(y + \epsilon) &= m\{x \in S : f(x) > y + \epsilon\} \\
 &= m\{x \in S : f(x) - \epsilon > y\} \\
 &= \mu_{f-\epsilon}(y) \\
 &\leq \mu_{f_N}(y) ,
 \end{aligned} \tag{6.2}$$

and then

$$R \int_0^\infty \mu_f - \epsilon m(S) \leq R \int_\epsilon^\infty \mu_f = R \int_0^\infty \mu_f(y + \epsilon) dy \leq R \int_0^\infty \mu_{f_N}(y) . \tag{6.3}$$

For  $y > \epsilon$  and  $N > K$ ,

$$\begin{aligned}
 \mu_{f_N}(y) &\leq \mu_{f+\epsilon}(y) \\
 &= m\{x \in S : f(x) + \epsilon > y\} \\
 &= m\{x \in S : f(x) > y - \epsilon\} \\
 &= \mu_f(y - \epsilon)
 \end{aligned} \tag{6.4}$$

and then

$$\begin{aligned}
 R \int_0^\infty \mu_{f_N} &\leq \epsilon m(S) + R \int_\epsilon^\infty \mu_{f_N} \\
 &\leq \epsilon m(S) + R \int_\epsilon^\infty \mu_f(y - \epsilon) dy \\
 &= \epsilon m(S) + R \int_0^\infty \mu_f .
 \end{aligned} \tag{6.5}$$

Thus, given  $\epsilon > 0$  there is a  $K \in \mathbb{Z}^+$  such that

$$-\epsilon m(S) \leq R \int_0^\infty \mu_{f_N} - R \int_0^\infty \mu_f \leq \epsilon m(S) \tag{6.6}$$

for  $N > K$  and, since  $\epsilon$  is arbitrary,

$$R \int_0^\infty \mu_{f_N} \rightarrow R \int_0^\infty \mu_f \tag{6.7}$$

as  $N \rightarrow \infty$ . Similarly,

$$R \int_{-\infty}^0 \mu_{f_N} \rightarrow R \int_{-\infty}^0 \mu_f \tag{6.8}$$

as  $N \rightarrow \infty$ . The last two limits and Definition 5 establish the desired convergence, Q.E.D.

The following theorem is usually cited as one of the triumphs of the Lebesgue theory because it replaces the uniform convergence of  $\{f_N\}$  by the less restrictive condition that the  $f_N$  be uniformly bounded. It also allows the domain of these functions to have arbitrary Lebesgue measure.

**THEOREM 9.** (The Lebesgue dominated convergence theorem) Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $\{f_N\}$  be a sequence of integrable functions on  $E$  that converges to an integrable function  $f$  on  $E$ . If there is a function  $g \in L(E)$  such

that  $|f_N| \leq g$  for all  $N \in \mathbb{Z}^+$ , then

$$\int_E f_N \rightarrow \int_E f \quad (6.9)$$

as  $N \rightarrow \infty$ .

This theorem is sometimes stated with the  $f_N$  and  $f$  measurable rather than integrable, but, in fact, these functions must be integrable. For the  $f_N$  this is so because  $g \in L(E) \Rightarrow -g \in L(E)$  and  $-g \leq f_N \leq g \Rightarrow \mu_{-g} \leq \mu_{f_N} \leq \mu_g$ . For  $f$  because  $f_N \rightarrow f$  and  $-g \leq f_N \leq g \Rightarrow -g \leq f \leq g$ .

The mechanism through which the uniform boundedness by the integrable function  $g$  operates in this theorem can be better understood by means of the next two lemmas, whose proof is temporarily postponed.

LEMMA 3. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $g \in L(E)$ . For any  $\epsilon > 0$  there is a subset  $F$  of  $E$  with  $m(F) < \infty$  such that

$$|\int_{E-F} g| < \epsilon. \quad (6.10)$$

LEMMA 4. Let  $F \subset \mathbb{R}^n$  be a measurable set, let  $g \in L(F)$  and let  $G$  be a subset of  $F$ . For any  $\epsilon > 0$  there is a  $\delta > 0$  such that if  $m(G) < \delta$  then

$$|\int_G g| < \epsilon. \quad (6.11)$$

Because of Lemma 3, the integrals of all functions involved in Theorem 9 are negligible outside a set  $F$  of finite measure and, because of Lemma 4, the same is true in a sufficiently small subset  $G$  of  $F$ . This is significant because, according to Egorov's theorem (Theorem 4), the set  $G$  can be chosen so that  $f_N \rightarrow f$  uniformly on  $F - G$ . And now, as in the case of Riemann integration, uniform convergence is seen to be the main driving force in the proof of Theorem 9 through Proposition 4.

PROOF OF THEOREM 9. Given  $\epsilon > 0$  let  $F$  be as in Lemma 3, let  $\delta$  be as in Lemma 4 and, for this choice of  $\delta$ , let  $G$  be as in Egorov's theorem. Using the triangle inequality and Theorems 6 and 7 we obtain

$$\begin{aligned} |\int_E f_N - \int_E f| &\leq |\int_F f_N - \int_F f| + \int_{E-F} |f_N| + \int_{E-F} |f| \\ &\leq |\int_F f_N - \int_F f| + 2 \int_{E-F} g \\ &\leq |\int_{F-G} f_N - \int_{F-G} f| + \int_G |f_N| + \int_G |f| + 2 \int_{E-F} g \\ &\leq |\int_{F-G} f_N - \int_{F-G} f| + 2 \int_G g + 2 \int_{E-F} g \\ &\leq |\int_{F-G} f_N - \int_{F-G} f| + 4\epsilon. \end{aligned} \quad (6.12)$$

By Proposition 4,  $N$  can be chosen so large that the first term on the right is smaller than  $\epsilon$ , and, since  $\epsilon$  is arbitrary, the result follows, Q.E.D.

PROOF OF LEMMA 3. In view of Theorem 7 we need only consider the case  $g \geq 0$ . By Definition 5, given  $\epsilon > 0$  there is an  $a > 0$  such that

$$R \int_0^a \mu_g^E < \epsilon. \quad (6.13)$$

If we define  $F = \{x \in E : g(x) > a\}$  we have  $m(F) = \mu_g^E(a) < \infty$  since  $g \in L(E)$ . But  $E - F = \{x \in E : 0 \leq g(x) \leq a\}$ , so that  $\mu_g^{E-F}(y) \leq \mu_g^E(y)$  if  $y < a$  and  $\mu_g^{E-F}(y) = 0$  if  $y \geq a$ . Then

$$\int_{E-F} g = R \int_0^\infty \mu_g^{E-F} \leq R \int_0^a \mu_g^E < \epsilon, \quad (6.14)$$

Q.E.D.

PROOF OF LEMMA 4. In view of Theorem 7 we need only consider the case  $g \geq 0$ . Given  $\epsilon > 0$  there is an  $a > 0$  such that

$$R \int_a^\infty \mu_g^F < \frac{\epsilon}{2}. \quad (6.15)$$

If  $G \subset F$  we have  $\mu_g^G \leq m(G)$  and  $\mu_g^G \leq \mu_g^F$ . Then, if we choose  $\delta = \epsilon/2a$  and if  $m(G) < \delta$ , we obtain

$$\begin{aligned} \int_G g &= R \int_0^a \mu_g^G + R \int_a^\infty \mu_g^G \\ &\leq a m(G) + R \int_a^\infty \mu_g^F \\ &< a m(G) + \frac{\epsilon}{2} < \epsilon, \end{aligned} \quad (6.16)$$

Q.E.D.

In some applications the sequence  $\{f_N\}$  may not be uniformly bounded by an integrable function. Such a case frequently occurs when the  $f_N$  are the partial sums of a series of positive functions. In this type of situation Theorem 11 below is useful, but in order to prove it we need some preparation. First we observe that the Lebesgue integral of a nonnegative function can be characterized in terms of those of bounded integrable functions.

LEMMA 5. Let  $E \subset \mathbb{R}^n$  be a measurable set and let  $f: E \rightarrow \mathbb{R}$  be a nonnegative measurable function. Then

$$\int_E f = \sup \int_E g, \quad (6.17)$$

where the supremum is taken over all bounded functions  $g \in L(E)$  such that  $0 \leq g \leq f$ .

PROOF. If  $f \in L(E)$ , given  $\epsilon > 0$  there is a  $b > 0$  such that

$$0 \leq \int_E f - R \int_0^b \mu_f < \epsilon. \quad (6.18)$$

If we define  $g = \min\{f, b\}$  then  $g$  is bounded,  $0 \leq g \leq f$ ,  $\mu_g(y) = \mu_f(y)$  if  $0 < y < b$  and  $\mu_g(y) = 0$  if  $y \geq b$ . We have

$$\int_E g = R \int_0^b \mu_g = R \int_0^b \mu_f, \quad (6.19)$$

so that  $g \in L(E)$  and

$$0 \leq \int_E f - \int_E g < \epsilon. \quad (6.20)$$

If  $f \notin L(E)$  define  $E_a = \{x \in E : f(x) > a\}$  for each  $a > 0$  and consider two cases.

(1)  $m(E_a) < \infty$  for all  $a > 0$ . For any  $M > 0$  there are  $a, b \in \mathbb{R}$  such that  $0 < a < b$  and

$$R \int_a^b \mu_f > M. \quad (6.21)$$

If we define  $g: E \rightarrow \mathbb{R}$  by  $g = \min\{f, b\}$  on  $E_a$  and  $g = 0$  on  $E - E_a$ , then  $g$  is bounded,  $0 \leq g \leq f$ ,  $\mu_g(y) = m(E_a)$  if  $0 < y < a$ ,  $\mu_g(y) = \mu_f(y)$  if  $a \leq y < b$  and  $\mu_g(y) = 0$  if  $y \geq b$ . It follows that  $g \in L(E)$  and that

$$\int_E g = R \int_0^a m(E_a) + R \int_a^b \mu_f > a m(E_a) + M. \quad (6.22)$$

(2)  $m(E_a) = \infty$  for some  $a > 0$ . Given  $M > 0$  let  $R$  be a rectangle so large that  $m(R \cap E_a) > M/a$ . If we define  $g(x) = a$  if  $x \in R \cap E_a$  and  $g(x) = 0$  otherwise, then  $g$  is bounded,  $0 \leq g \leq f$  and

$$\int_E g = \int_{R \cap E_a} a = a m(R \cap E_a) > M, \quad (6.23)$$

Q.E.D.

**THEOREM 10. (Fatou's lemma)** Let  $E \subset \mathbb{R}^n$  be a measurable set, let  $\{f_N\}$  be a sequence of nonnegative measurable functions on  $E$  and let  $f_N \rightarrow f$ . Then for any  $\epsilon > 0$  there is an  $M \in \mathbb{Z}^+$  such that

$$\int_E f < \int_E f_N + \epsilon \quad (6.24)$$

for  $N > M$ .

**PROOF.** For any  $\epsilon > 0$ , Lemma 5 implies that there is a bounded function  $g \in L(E)$ , such that  $0 \leq g \leq f$  and

$$\int_E f - \int_E g < \frac{\epsilon}{2}. \quad (6.25)$$

If we define  $g_N = \min\{g, f_N\}$  for each  $N \in \mathbb{Z}^+$  then  $g$  is measurable because  $\{x \in E : g_N(x) > y\} = \{x \in E : g(x) > y\} \cap \{x \in E : f_N(x) > y\}$  for each  $y > 0$ , and  $g_N \leq f_N$  implies that

$$\int_E g_N \leq \int_E f_N. \quad (6.26)$$

Also, since  $g_N \rightarrow g$  on  $E$  and  $|g_N| \leq g$ , the Lebesgue dominated convergence theorem implies that

$$\int_E g_N \rightarrow \int_E g \quad (6.27)$$

as  $N \rightarrow \infty$ . Therefore, there is an  $M > 0$  such that

$$\int_E g \leq \int_E f_N + \frac{\epsilon}{2} \quad (6.28)$$

for  $N > M$ , or else we would have

$$\int_E g_N \leq \int_E f_N < \int_E g - \frac{\epsilon}{2} \quad (6.29)$$

for arbitrarily large  $N$ , contradicting the limit above. We conclude that

$$\int_E f < \int_E g + \frac{\epsilon}{2} < \int_E f_N + \epsilon \quad (6.30)$$

for  $N > M$ , Q.E.D.

THEOREM 11. (The Levi monotone convergence theorem) Let  $E \subset \mathbb{R}^n$  be a measurable set, let  $\{f_N\}$  be a sequence of nonnegative functions on  $E$  such that  $f_M \leq f_N$  for  $M < N$ , and let  $f_N \rightarrow f$ . Then

$$\int_E f_N \rightarrow \int_E f \quad (6.31)$$

as  $N \rightarrow \infty$ .

PROOF.  $f_M \leq f_N$  for  $M < N \Rightarrow f_N \leq f$  for all  $N \in \mathbb{Z}^+ \Rightarrow$

$$\int_E f_N \leq \int_E f \quad (6.32)$$

for all  $N \in \mathbb{Z}^+$ . By Fatou's lemma, given  $\epsilon > 0$

$$\int_E f < \int_E f_N + \epsilon \quad (6.33)$$

for  $N$  large enough. These two inequalities imply that

$$\int_E f_N \rightarrow \int_E f \quad (6.34)$$

as  $N \rightarrow \infty$ , Q.E.D.

#### REFERENCES

1. ROYDEN, H.L. Real Analysis, 2nd. ed., Macmillan, New York, 1968.
2. APOSTOL, T.M. Mathematical Analysis, 2nd. ed., Addison-Wesley, Reading, Mass., 1974.
3. MCSHANE, E.J. Unified Integration, Academic Press, Orlando, Fla., 1983.
4. MIKUSINSKI, J. The Bochner Integral, Birkhäuser Verlag, Boston, 1978.
5. REY PASTOR, J. Elementos de la Teoría de Funciones, 3rd. ed., Ibero-American, Madrid, 1953.
6. KESTELMAN, H. Modern Theories of Integration, Oxford University Press, Oxford, 1937.

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

### Guest Editors

**Edson Denis Leonel**, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)