

**A CONSTRUCTION OF A BASE FOR THE M FOLD TENSOR PRODUCT
 OF A BRANCH SPACE**

ROHAN HEMASINHA

The University of West Florida
 Pensacola, Florida 32514

(Received August 3, 1987)

ABSTRACT. Let E be a Banach space with Schauder base $(x_n)_n$. Let $\overset{m}{\bullet}_{\alpha} E$ denote the completion of the m fold tensor product with respect to a reasonable cross norm α . We show that the set $\{x_{i_1} \bullet \dots \bullet x_{i_m} : x_{i_j} \in (x_n)_n\}$ can be enumerated so that for each positive integer k , the first k^m terms are precisely all the elements of the form $x_{i_1} \bullet \dots \bullet x_{i_m}$ with $i_1, \dots, i_m \in \{1, \dots, k\}$ and the set so arranged is a Schauder base for $\overset{m}{\bullet}_{\alpha} E$.

KEYWORDS AND PHRASES. m fold tensor product, projective, injective and reasonable cross norms, Schauder bases.

1980 AMS SUBJECT CLASSIFICATION CODE. 46B15, 46M05.

1. INTRODUCTION.

Given a Banach space E , we will denote by $\overset{m}{\bullet} E$ the m fold algebraic tensor product and by $\overset{m}{\bullet}_{\alpha} E$ its completion with respect to reasonable norm α . If $(x_n)_n$ is a Schauder base for E then $\overset{m}{\bullet}(x_n)$ will denote the set

$$\{x_{i_1} \bullet \dots \bullet x_{i_m} : x_{i_j} \in (x_n)_n, 1 \leq j \leq m\}.$$

For $m > 2$ the projective (injective) tensor norm on $\overset{m}{\bullet} E$ will be denoted by $\gamma_m(\lambda)$. If $m = 2$ this norm will be denoted by $\gamma(\lambda)$.

In [1], [4] it is shown that if E_1, E_2 are Banach spaces with Schauder bases $(x_n)_n, (y_n)_n$ then for any reasonable norm α , the space $E_1 \bullet_{\alpha} E_2$ has a Schauder base (z_n) with the following properties.

- (1) (z_n) is obtained by enumerating the set $(x_n)_n \bullet (y_n)_n$;
- (2) the enumeration is such that for any positive integer k , the first k^2 terms are precisely all elements of the form

$$x_{i_1} \bullet y_{i_2}, \quad i_1, i_2 \in \{1, \dots, k\}.$$

In this paper we show that for any positive integer m , the space ${}^m_{\alpha}E$ has a Schauder base obtained by enumerating ${}^m(x_n)_n$ in such a way that for any positive integer k , the first k^m terms are precisely all elements of the form

$$x_{i_1} \bullet \dots \bullet x_{i_m} \text{ with } i_1, \dots, i_m \in \{1, \dots, k\}.$$

Our proof is via an induction argument and in a forthcoming paper we utilize this construction to derive some properties of the symmetric tensor algebra of a Banach space. Note that even for the case $m = 3$, iteration to $(E \bullet_{\gamma} E) \bullet_{\gamma} E$ of the enumerating scheme described in [1] will not yield a base with the above mentioned properties.

For definitions and terminology regarding Schauder bases, tensor products and tensor norms we refer the reader to [4], [3], and [2].

We denote the following property (*) of the projective tensor norm. For $m > 2$ let u be an element of ${}^m_{\bullet}E$. It can easily be shown that

$$(*) \quad \gamma(u) = \gamma_m(u)$$

where $\gamma(u)$ is the projective norm of u when u is considered as an element of $F \bullet E$ with $F = {}^{m-1}_{\bullet}E$ being endowed with the norm γ_{m-1} . We shall make use of this fact subsequently.

We now state our theorem.

THEOREM. Let (x_n, f_n) , $x_n \in E$, $f_n \in E'$ be biorthogonal system such that $(x_n)_n$ is a Schauder base for E . Let α be a reasonable norm on ${}^m_{\bullet}E$. Then there exists for ${}^m_{\alpha}E$ a biorthogonal system (z_n, g_n) with the following properties.

- (1) As sets $\{z_n : n \in N\} = {}^m_{\bullet}(x_n)_n$
 $\{g_n : n \in N\} = {}^m_{\bullet}(f_n)_n$.
- (2) The enumeration of (z_n, g_n) is such that for each positive integer k the first k^m terms are precisely all the tensors of the form
 $x_{i_1} \bullet \dots \bullet x_{i_m}$,
 $f_{i_1} \bullet \dots \bullet f_{i_m}$ with $i_1, \dots, i_m \in \{1, \dots, k\}$.
- (3) The sequence $(z_n)_n$ is a Schauder base for ${}^m_{\alpha}E$.

PROOF. It is known that [2] $\lambda_m < \alpha < \gamma_m$ for any reasonable norm α . Also ${}^m_{\bullet}E' \subseteq ({}^m_{\lambda_m}E)'$. Thus any enumeration of ${}^m_{\bullet}(x_n)_n$ together with the corresponding enumeration of ${}^m_{\bullet}(f_n)_n$ yields a biorthogonal system for ${}^m_{\alpha}E$. It is easily shown that the linear span of ${}^m_{\bullet}(x_n)_n$ is dense in ${}^m_{\gamma_m}E$ and we shall first establish the assertions of the theorem for $\alpha = \gamma_m$.

Now for $m = 2$ the base constructed in [1] has the stated properties.

Suppose that a biorthogonal system (z_n, g_n) with the stated properties can be constructed for ${}^m_{\gamma_m}E$.

Consider the sequence $(w_n)_{n \in \mathbb{N}}$ in $E^{\otimes m+1}$ defined by the scheme in the accompanying figure. For each positive integer k , the terms

$$w_{k+1}^{\otimes m+1}, w_{k+2}^{\otimes m+1}, \dots, w_{(k+1)^{m+1}}^{\otimes m+1}$$

are described by the tensor beneath each term.

Figure 1

k , a positive integer; ℓ an integer $0 < \ell < k$

$w_{k+1}^{\otimes m+1}$,	$w_{k+2}^{\otimes m+1}$,	$w_{(k+1)^m+k^m(k-1)}$
$x_1 \otimes z_{k+1}^{\otimes m}$	$x_1 \otimes z_{k+2}^{\otimes m}$	$x_1 \otimes z_{(k+1)^m}$
$w_{(k+1)^m+k^m(k-1)+1}$,	$w_{(k+1)^m+k^m(k-1)+2}$,	$w_{2(k+1)^m+k^m(k-2)}$
$x_2 \otimes z_{k+1}^{\otimes m}$	$x_2 \otimes z_{k+2}^{\otimes m}$	$x_2 \otimes z_{(k+1)^m}$
⋮	⋮	⋮
$w_{\ell(k+1)^m+k^m(k-\ell)+1}$,	⋮	$w_{(\ell+1)(k+1)^m+k^m(k-(\ell+1))}$
$x_{\ell+1} \otimes z_{k+1}^{\otimes m}$	⋮	$x_{\ell+1} \otimes z_{(k+1)^m}$
$w_{(k-1)(k+1)^m+k^m+1}$,	⋮	$w_{k(k+1)^m}$
$x_k \otimes z_{k+1}^{\otimes m}$	⋮	$x_k \otimes z_{(k+1)^m}$
$w_{k(k+1)^m+1}$,	$w_{k(k+1)^m+2}$,	$w_{(k+1)^{m+1}}$
$x_{k+1} \otimes z_1$	$x_{k+1} \otimes z_2$	$x_{k+1} \otimes z_{(k+1)^m}$

In the preceding tableau each of the first k rows contains $(k+1)^m - k^m$ entries. The last row has $(k+1)^m$ entries, and so altogether we have exhibited $k((k+1)^m - k^m) + (k+1)^m = (k+1)^{m+1} - k^{m+1}$ entries.

The sequence $(h_n)_{n \in \mathbb{N}}$ is obtained by enumerating the set $\{g_p \otimes f_q \mid g_p \in (g_n), f_q \in (f_n)\}$ according to the same scheme.

In view of the inductive hypothesis the system (w_n, h_n) is a biorthogonal system with the property that the first k^{m+1} terms are all of the form

$$x_{i_1} \bullet \dots \bullet x_{i_{m+1}}, f_{i_1} \bullet \dots \bullet f_{i_{m+1}}$$

with $i_1, \dots, i_{m+1} \in \{1, \dots, k\}$.

Let S_n, T_n, W_n respectively denote the n^{th} partial sum operators of the systems $(x_n, f_n), (z_n, g_n)$ and (w_n, h_n) .

Since $(x_n), (z_n)$ are bases the sequences $(f_n), (T_n)$ are bounded in operator norm by some constant M . To show that (w_n) is a base it thus suffices to show that for some positive M' , $\|W_n\| \leq M'$ for all n ([4], p. 25).

Now, given a positive integer n the defining scheme for (w_j) (see fig. 1) shows that W_n can be expressed as sums of tensor products of the operators S_m, T_m and f_m .

Indeed, let us consider the three possible cases.

Case 1. If $n = k^{m+1}$ for some positive integer k then

$$W_n = \bullet^{m+1} S_k, \text{ the } m+1 \text{ fold tensor product of } S_k.$$

Case 2. If $n = \ell (k+1)^m + k^m(k-\ell) + r$, with $0 < \ell < k$ and

$$1 \leq r < (k+1)^m - k^m \text{ then}$$

$$W_n = \bullet^{m+1} S_k + (T_{(k+1)^m} - T_{k^m}) \bullet S_\ell + (T_{k^{m+r}} - T_{k^m}) \bullet f_{\ell+1}.$$

Case 3. If $n = k(k+1)^m + r$ with $1 \leq r < (k+1)^m$ then

$$W_n = \bullet^{m+1} S_k + (T_{(k+1)^m} - T_{k^m}) \bullet S_k + T_r \bullet f_{k+1}$$

In u is in $\bullet^{m+1} E$ then property (*) together with the fact that γ_{m+1} is reasonable yields the following inequalities

$$(1) \gamma_{m+1}(\bullet^{m+1} S_k(u)) \leq \|S_k\|^{m+1} \gamma_{m+1}(u)$$

$$(2) \gamma_{m+1}((T_{(k+1)^m} - T_{k^m}) \bullet S_\ell(u)) \leq \|T_{(k+1)^m} - T_{k^m}\| \|S_\ell\| \gamma_{m+1}(u)$$

$$(3) \gamma_{m+1}((T_{k^{m+r}} - T_{k^m}) \bullet f_{\ell+1}(u)) \leq \|T_{k^{m+r}} - T_{k^m}\| \|f_{\ell+1}\| \gamma_{m+1}(u).$$

We utilized (*) in deriving (2) and (3). Hence,

$$Y_{m+1}(W_n(u)) \leq (M^{m+1} + 2M^2 + 2M^2)Y_{m+1}(u)$$

in all three cases. It now follows from the uniform boundedness principle that $(\|W_n\|)$ is bounded.

To complete the proof let us recall that if α is a reasonable norm then $\lambda_m \leq \alpha \leq \gamma_m$. Furthermore, $\overset{m}{\circlearrowleft}_{\alpha} E$ is the completion of $\overset{m}{\circlearrowleft} E$ with respect to α . Consequently for all such α , the system (z_n, g_n) is a complete biorthogonal system whose sequence of partial sums T_n is pointwise bounded and hence bounded in operator norm. This means that (z_n) is a Schauder base for $\overset{m}{\circlearrowleft}_{\alpha} E$.

CONCLUDING REMARKS.

It would be interesting to find out whether the base for $\overset{m}{\circlearrowleft}_{\gamma_m} E$ obtained by iteration of the process in [1] is equivalent to the base described in this paper. We hope to investigate this problem in a future paper.

REFERENCES

1. GELBAUM, B.R. and GIL DE LA MADRID, J. Bases for Tensor Products of Banach Spaces, Pacific J. Math. 11 (1961), pp. 1281-1286.
2. CIGLER, J., LOSERT, V. and MICHOR, P. Banach Modules and Functors on Categories of Banach Spaces. Lecture Notes in Pure and Applied Mathematics, Marcel Dekker, Inc., (1979).
3. SCHATTEN, R. A Theory of Cross Spaces. Annals of Mathematics Studies, 26, Princeton University Press, (1950).
4. SINGER, I. Bases for Banach Spaces, Vol. 1. A Series of Comprehensive Studies in Mathematics, 154. Springer-Verlag, (1970).

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru