

FRACTIONAL POWERS OF NON-NEGATIVE OPERATORS IN FRÉCHET SPACES

C. MARTINEZ, M. SANZ and V. CALVO

Universidad de Valencia
Facultad de Matemáticas
Departamento de Matemática Aplicada y Astronomía
Burjassot, Valencia, Spain

(Received April 21, 1988 and in revised form August 16, 1988)

ABSTRACT. In the present paper the theory of fractional powers, which has been restricted to date to certain operators on Banach spaces, is generalized to certain particular operators in Fréchet spaces. The main difficulty consists in the fact that neither the holomorphic functional calculus nor the results on Banach algebras are available for bounded operators on Fréchet spaces.

All the basic properties which a good theory of fractional powers must fulfill are proved, except for the spectral relation,

$$\sigma(A^\alpha) = \{z^\alpha : z \in \sigma(A)\}$$

which remains an unsolved problem.

KEY WORDS AND PHRASES. Fractional powers operators. Fractional derivatives and integrals.
1980 AMS SUBJECT CLASSIFICATION CODE. 47A60, 26A33.

1. INTRODUCTION AND NOTATION.

In [1] we have developed a new theory of fractional powers for non-negative operators, that is, for closed linear operators A in a Banach space X such that $]-\infty, 0[$ is contained in the resolvent set $\rho(A)$ and

$$\sup_{\lambda > 0} ||\lambda(\lambda + A)^{-1}|| < +\infty.$$

To this end, we have used a new definition of fractional powers which coincides with the closure of the operator J^α (defined by A.V. Balakrishnan in [2]) when $D(A)$ is dense in X , and is a strict extension of that operator when A is not densely defined.

The point of view introduced in [1] has proved advantageous in several aspects. First of all, it implies a substantial simplification of the theory and it shows that the denseness hypothesis on the domain $D(A)$, considered as fundamental in previous theories, is irrelevant; furthermore, it provides for the first time in the extensive literature on fractional powers, a satisfactory construction in the case of non-dense domains.

The main theorem of [1] establishes the following relation, for $\epsilon > 0$ and $\operatorname{Re} \alpha > 0$:

$$A^\alpha [(A+\epsilon)^{-1}]^\alpha = [A(A+\epsilon)^{-1}]^\alpha \quad (1.1)$$

In the present paper we generalize the results of [1] to operators A in Fréchet spaces that verify a condition similar to non-negativity. The essential difficulty introduced by substituting the Fréchet structure for the Banach structure consists in the loss of properties undergone by the resolvent function $z \mapsto (z-A)^{-1}$, which is no longer analytical nor even continuous.

In 1984, Wilson Lamb [3] generalizes the concept of non-negative operators to Fréchet spaces in order to obtain certain results in the field of fractional calculus. He gives a definition of fractional powers on operators, A , such that $(\lambda+A)^{-1}$ exists and is bounded for each $\lambda > 0$, and given any seminorm, $\|\cdot\|_p$, among those that describe the space topology, there is another seminorm $\|\cdot\|_{q(p)}$ and a constant $C > 0$ (independent of $\|\cdot\|_p$) such that for each vector ϕ :

$$\|(C\lambda(\lambda+A)^{-1})^n \phi\|_p < \|\phi\|_{q(p)}, \text{ for } n = 1, 2, \dots \quad (1.2)$$

This condition assures that the resolvent function is analytical on a sector centered around the negative real semiaxis, thus allowing application of the same methods and arguments as for Banach spaces.

We propose a different kind of operator which seems more natural and strictly includes those of W. Lamb. Indirectly, this will be shown by means of a counterexample in which the resolvent function associated with a certain operator of our class is not analytical. The lack of this property prevents us from using or generalizing the techniques common to theories on fractional powers previous to [1]). However, we can take advantage of some of the ideas contained in [1].

Being unable to prove identity (1.1) for Fréchet spaces, we have chosen to argue as follows: We show directly that

$$D(A^\alpha) = D[(A+\epsilon)^\alpha], \text{ for all } \epsilon > 0,$$

and, as a consequence, we find an integral expression for the operator

$$A^\alpha = \sum_{p=0}^n (-1)^p \epsilon^p \binom{\alpha}{p} (A + \epsilon)^{\alpha-p} \quad (1.3)$$

on $D(A^\alpha)$, which will enable us to prove the essential property that a good fractional power must have, namely

$$A^\alpha A^\beta = A^{\alpha + \beta} \quad (\text{with } \operatorname{Re} \alpha > 0 \text{ and } \operatorname{Re} \beta > 0).$$

On the other hand, it can be seen that the integral expression for $(A^{\alpha+\beta})^{-1}$ and the relation

$$(A^\alpha)^3 = A^{\alpha\beta} \quad (\text{with } 1 > \alpha > 0 \text{ and } \operatorname{Re} \beta > 0)$$

established in [1] for Banach spaces, are still valid in Fréchet spaces. However, in the latter case the spectral mapping theorem

$$\sigma(A^\alpha) = \{z^\alpha : z \in \sigma(A)\}$$

remains an open problem, at least as far as the present article is concerned.

F will henceforth be a Fréchet space, and $\{\|\cdot\|_p\}_{p \geq 1}$ an increasing sequence of semi-norms defining the topology of F . We shall denote by A a linear operator on F with domain $D(A)$, and by $\rho(A)$ and $\sigma(A)$ the resolvent set and the spectrum of A , respectively.

The proofs of several of the results in the present paper are essentially the same as the proofs of the corresponding results in [1]. These proofs will therefore be omitted.

2. PROPERTIES OF FRACTIONAL OPERATORS.

DEFINITION 2.1. We say that A is non-negative if $\rho(A) \supset \{\lambda e^{i\pi} : \lambda > 0\}$ and for all $n \in \mathbb{N}$, there exists $M_n > 0$ and there exists $m(n) \in \mathbb{N}$ such that

$$\|\lambda(\lambda + A)^{-1}\phi\|_n \leq M_n \|\phi\|_{m(n)} \text{ for all } \phi \in F \text{ and for all } \lambda > 0. \quad (2.1)$$

EXAMPLE 1. Given $p: 1 < p < +\infty$, let

$F = \{f: [0, +\infty[\rightarrow \mathbb{C} \text{ such that } f|_{[0,a]} \in L^p([0,a]), \text{ for all } a > 0\}$ with
 $\|f\|_n = \|f|_{[0,n]}\|_p$.

Taking $A = -d/dx$ on the set

$D(A) = \{f \in F; f|_{[0,n]} \in W^{1,p}([0,n]), \text{ for all } n \in \mathbb{N} \text{ and } f(0) = 0\}$

it is easy to show that A is a non-negative unbounded operator, which is densely defined if and only if $p < +\infty$.

EXAMPLE 2. Let $F = \{f \in C^\infty([0, 1]), \text{ with } f^{(n)}(0) = 0, \text{ for } n=0,1,2,\dots\}$ with the seminorms

$$\|f\|_p = \max_{0 \leq q \leq p} \|f^{(q)}\|_\infty$$

and the (bounded) operator

$$A = -i d/dx$$

It can easily be shown that A has an empty spectrum and that for all $\lambda > 0$ and $g \in F$

$$\|\lambda(A + \lambda)^{-1}g\|_p \leq 2 \|g\|_{p+1}$$

and thus it is a non-negative operator.

We shall now see that for all $w: 0 < w < \pi/2$ the operator $e^{-iw}A$ is not non-negative. This will show that definition 2.1 is not equivalent to W. Lamb's definition; for if an operator satisfies (1.2), this is still true if we multiply the operator by e^{-iw} , if $w > 0$ is small enough.

For $z = \lambda + i\sigma\lambda$ ($\lambda > 0$ and $\sigma = \tan w$) and $g \in F$, we have

$$(A-z)^{-1}g(x) = e^{-i\lambda x + \sigma\lambda x} \int_0^x e^{i\lambda t - \sigma\lambda t} i g(t) dt$$

and consequently

$$|z(A-z)^{-1}g(x)| = \sqrt{1+\sigma^2} e^{\sigma\lambda x} \left| \int_0^x e^{i\lambda t - \sigma\lambda t} g(t) d(\lambda t) \right|.$$

Using $g_\lambda(x) = e^{-i\lambda x} - 1/\lambda x$, we obtain

$$\|z(A-z)^{-1}g_\lambda\|_\infty = \sqrt{1+\sigma^2} e^{\sigma\lambda} \int_0^\lambda e^{-\sigma\xi-1/\xi} d\xi.$$

On the other hand, $g_\lambda^{(n)}(x) = \lambda^n g_1^{(n)}(\lambda x)$ implies $\|g_\lambda^{(n)}\|_\infty = \lambda^n \sup_{\tau \in [0, \lambda]} |g_1^{(n)}(\tau)|$

Thus

$$\lim_{\lambda \rightarrow +\infty} \frac{\|z(A-z)^{-1}g_\lambda\|_\infty}{\|g_\lambda^{(n)}\|_\infty} = \sqrt{1+\sigma^2} \frac{\int_0^{+\infty} e^{-\sigma\xi-1/\xi} d\xi}{\sup_{\tau > 0} |g_1^{(n)}(\tau)|} \lim_{\lambda \rightarrow +\infty} \frac{e^{\sigma\lambda}}{\lambda^n} = +\infty$$

And we can conclude that $e^{-iw}A$ is not non-negative.

REMARK 2.1. It can also be shown that the operator A from example 2 does not satisfy (1.2) even in the case that in this condition the constant C should be dependent on the seminorm $\|\cdot\|_p$, which is contained in the left-hand-side.

In the next definition, as well as all the other statements in this paper, we assume that A is a non-negative operator.

DEFINITION 2.2. The condition that A be non-negative allows us to define the integral operator J^α in the same way as it was defined in [1] for Banach spaces, and the following property holds:

$$J^\alpha J^\beta = J^{\alpha+\beta} \quad (\text{with } \operatorname{Re} \alpha > 0 \text{ and } \operatorname{Re} \beta > 0).$$

If A is continuous, we define

$$A^\alpha = J^\alpha. \quad (2.2)$$

If A is not continuous and $0 \notin \rho(A)$, we have that the operator A^{-1} is a continuous non-negative operator, since

$$\| \lambda(A^{-1} + \lambda)^{-1}\phi \|_n = \| A(A + \lambda^{-1})^{-1}\phi \|_n = \| \phi - \lambda^{-1}(A + \lambda^{-1})^{-1}\phi \|_n <$$

$$< \| \phi \|_n + M_n \| \phi \|_{m(n)} < (1 + M_n) \| \phi \|_{k(n)}$$

(with $k(n) = \max\{n, m(n)\}$) for all $\lambda > 0$ and $\phi \in F$.

$(A^{-1})^\alpha$ is a one-to-one operator, since if $n > \operatorname{Re} \alpha$, we have
 $(A^{-1})^{n-\alpha}(A^{-1})^\alpha = (A^{-1})^n$,

and consequently:

$$(A^{-1})^\alpha \phi = 0 \implies (A^{-1})^n \phi = 0 \implies \phi = 0.$$

We define

$$A^\alpha = ((A^{-1})^\alpha)^{-1} \quad (2.3)$$

It is easy to prove that, for $0 < \operatorname{Re} \alpha < 1$ and $\phi \in D(A)$

$$((A^{-1})^\alpha)^{-1} \phi = \frac{\sin \alpha \pi}{\pi} \int_0^{+\infty} \lambda^{\alpha-1} (\lambda + A)^{-1} A \phi d\lambda.$$

Finally, if A is not continuous and $0 \notin \rho(A)$, we define

$$A^\alpha \phi = \lim_{\epsilon \rightarrow 0} (A + \epsilon)^\alpha \phi \quad (2.4)$$

on the vectors ϕ for which the limit exists.

REMARKS. i) From (2.2) and (2.3) it follows, by direct composition of the powers (or their inverses) in their integral form, as shown in [2], that

$$A^\alpha A^\beta = A^{\alpha+\beta} \quad (\text{with } \operatorname{Re} \alpha > 0 \text{ and } \operatorname{Re} \beta > 0)$$

when A is continuous, and A not being continuous, when $0 \notin \rho(A)$.

ii) Arguing as in [1] it can be proved that when A is not continuous, J^α is a closable operator and A^α is an extension of the closure of J^α .

The subsequent results will give a more precise meaning to the definition of A^α in the more general case in which $0 \notin \rho(A)$ and A is not continuous.

PROPOSITION 2.1. When A is not continuous, the domain of $(A + \epsilon)^\alpha$ is independent of $\epsilon > 0$ (and equal to $D(A^\alpha)$ if $0 \in \rho(A)$).

PROOF. We first assume $0 < \operatorname{Re} \alpha < 1$. Starting with the identity

$$D[(A + \epsilon)^\alpha] = R([(A + \epsilon)^{-1}]^\alpha)$$

and using the integral expression of the operator J^α associated with the operator $(A + \epsilon)^{-1}$, we obtain

$$D[(A + \epsilon)^\alpha] = \left\{ \int_0^{+\infty} \lambda^{-\alpha} (\lambda + \epsilon + A)^{-1} \phi d\lambda : \phi \in F \right\}, \text{ for all } \epsilon > 0.$$

On the other hand, given $\epsilon, n > 0$ the resolvent formula leads to

$$\begin{aligned} & \int_0^{+\infty} \lambda^{-\alpha} (\lambda + \epsilon + A)^{-1} \phi d\lambda - \int_0^{+\infty} \lambda^{-\alpha} (\lambda + n + A)^{-1} \phi d\lambda = \\ & = (n - \epsilon) \int_0^{+\infty} \lambda^{-\alpha} (\lambda + \epsilon + A)^{-1} (\lambda + n + A)^{-1} \phi d\lambda, \text{ for all } \phi \in F, \end{aligned}$$

and the operator A can be introduced into the integrand on the right-hand side with no influence on the convergence of the integral; for this reason

$$[(A + \varepsilon)^{-1}]^\alpha \phi - [(A + \eta)^{-1}]^\alpha \phi \in D(A),$$

and thus

$$[(A + \varepsilon)^{-1}]^\alpha \phi - [(A + \eta)^{-1}]^\alpha \phi \in D[(A + \varepsilon)^\alpha] \cap D[(A + \eta)^\alpha]$$

for all $\phi \in F$, so we get

$$D[(A + \varepsilon)^\alpha] = D[(A + \eta)^\alpha].$$

Next we shall prove by induction on n that, for $0 < \operatorname{Re} \alpha < 2^n$, there exists $\beta(\alpha)$ with $\operatorname{Re} \beta(\alpha) > \operatorname{Re} \alpha$, such that for all $\phi \in F$

$$[(A + \varepsilon)^{-1}]^\alpha \phi - [(A + \eta)^{-1}]^\alpha \phi \in D[(A + \varepsilon)^{\beta(\alpha)}] \cap D[(A + \eta)^{\beta(\alpha)}] \quad (2.5)$$

which will imply the equality of the domains $D[(A + \varepsilon)^\alpha]$ and $D[(A + \eta)^\alpha]$.

This result has already been proved for $n = 0$. Assuming that it holds up to $n-1$, take α such that $0 < \operatorname{Re} \alpha < 2^n$. We have

$$[(A + \varepsilon)^{-1}]^\alpha \phi - [(A + \eta)^{-1}]^\alpha \phi = [(A + \varepsilon)^{-1}]^{\alpha/2} \psi + [(A + \eta)^{-1}]^{\alpha/2} \psi \quad (2.6)$$

$$\text{with } \psi = [(A + \varepsilon)^{-1}]^{\alpha/2} \phi - [(A + \eta)^{-1}]^{\alpha/2} \phi$$

and this element belongs to $D[(A + \varepsilon)^{\beta(\alpha/2)}] \cap D[(A + \eta)^{\beta(\alpha/2)}]$ by the induction hypothesis.

From (2.6) it is easily concluded that $[(A + \varepsilon)^{-1}]^{\alpha/2} \psi$ and $[(A + \eta)^{-1}]^{\alpha/2} \psi$ belong to the set

$$D[(A + \varepsilon)^{\alpha/2 + \beta(\alpha/2)}] \cap D[(A + \eta)^{\alpha/2 + \beta(\alpha/2)}]$$

and relation (2.5) is established (with $\beta(\alpha) = \alpha/2 + \beta(\alpha/2)$).

If $0 \in \rho(A)$, then A^α is defined in the same way as $(A + \varepsilon)^\alpha$, and therefore the argument is still valid if the condition $\eta > 0$ is substituted by $\eta = 0$. This yields $D[(A + \varepsilon)^\alpha] = D(A^\alpha)$.

PROPOSITION 2.2. When A is not continuous we denote by D_α the set $D[(A + \varepsilon)^\alpha]$. Then the function

$$h(t) = (A + t)^\alpha \phi,$$

with $\phi \in D_\alpha$ and $t > 0$, is Hölder continuous in each bounded interval $[0, a]$, with exponent $\operatorname{Re} \alpha / 2^n$, as long as $0 < \operatorname{Re} \alpha < 2^n$. Consequently

$$\lim_{\varepsilon \rightarrow 0} (A + \varepsilon)^\alpha \phi = A^\alpha \phi$$

and $D(A^\alpha) = D_\alpha$.

PROOF. We have for $0 < \operatorname{Re} \alpha < 1$ and $\phi \in D(A)$

$$(A+\varepsilon)^\alpha \phi - (A+\eta)^\alpha \phi = \frac{\sin \alpha\pi}{\pi} \left[\int_0^\delta \lambda^{\alpha-1} (-\lambda(A+\varepsilon+\lambda)^{-1}\phi + \lambda(A+\eta+\lambda)^{-1}\phi) d\lambda + \int_\delta^{+\infty} \lambda^\alpha (\varepsilon-\eta)(A+\varepsilon+\lambda)^{-1}(A+\eta+\lambda)^{-1} \phi d\lambda \right] \quad (2.7)$$

using the identity $(A+\varepsilon)(A+\varepsilon+\lambda)^{-1}\phi = \phi - \lambda(A+\varepsilon+\lambda)^{-1}\phi$ and the resolvent formula. The two integrals are absolutely convergent, for all $\phi \in F$.

Applying the formula (2.7) to the element $(A+1)^{-1}\phi$, for all $\phi \in D_\alpha$ and taking into account that the one-to-one operator $(A+1)^{-1}$ can be placed on the left, since it commutes with the integral sign we conclude that (2.7) holds for all $\phi \in D_\alpha$. Taking $\delta = |\varepsilon - \eta|$, we have

$$\begin{aligned} \left\| (A+\varepsilon)^\alpha \phi - (A+\eta)^\alpha \phi \right\| &< \left| \frac{\sin \alpha\pi}{\pi} \right| \left[\left| \int_0^{|\eta-\varepsilon|} \lambda^{\operatorname{Re} \alpha-1} 2 M_n \|\phi\|_{m(n)} d\lambda \right| + \right. \\ &\quad \left. + \int_{|\eta-\varepsilon|}^{+\infty} \lambda^{\operatorname{Re} \alpha-2} |\eta-\varepsilon| M_n M_{m(n)} \|\phi\|_{m(m(n))} d\lambda \right] \end{aligned}$$

Hence

$$\left\| (A+\varepsilon)^\alpha \phi - (A+\eta)^\alpha \phi \right\| < \left| \frac{\sin \alpha\pi}{\pi} \right| |\eta-\varepsilon|^{\operatorname{Re} \alpha} \left[\frac{2 M_n}{\operatorname{Re} \alpha} \|\phi\|_{m(n)} + \frac{M_n M_{m(n)}}{\operatorname{Re} \alpha - 1} \|\phi\|_{m(m(n))} \right]$$

and consequently $h(t) = (A+t)^\alpha \phi$, with $\phi \in D_\alpha$ and $t > 0$ is Hölder continuous with exponent $\operatorname{Re} \alpha$, so h is continuous and

$$\lim_{\varepsilon \rightarrow 0} (A+\varepsilon)^\alpha \phi$$

exists.

Arguing by induction, it is easy to show that in each bounded interval $[0, a]$ h is a Hölder continuous with exponent $\operatorname{Re} \alpha/2^n$ with $n \in \mathbb{N}$ such that $0 < \operatorname{Re} \alpha < 2^n$. For $n=0$ this statement has actually been proved, and assuming it holds up to $n-1$ and α such that $0 < \operatorname{Re} \alpha < 2^n$, then the relation

$$\left\| (A+\varepsilon)^\alpha \phi - (A+\eta)^\alpha \phi \right\| = \left\| (A+\varepsilon)^{\alpha/2} \psi - (A+\eta)^{\alpha/2} \psi \right\|$$

with

$$\psi = (A+\varepsilon)^{\alpha/2} \phi + (A+\eta)^{\alpha/2} \phi$$

implies, by the induction hypothesis, the property that was claimed for h .

Therefore h is continuous and converges as t tends to zero. Thus, if A is not continuous and $0 \notin \rho(A)$, by the definition of $D(A^\alpha)$, this set is equal to D_α . On the other hand, if $0 \in \rho(A)$, both domains being equal by Proposition 2.1., we also conclude that the following equation holds:

$$\lim_{\varepsilon \rightarrow 0} (A+\varepsilon)^\alpha \phi = A^\alpha \phi, \text{ for all } \phi \in D(A)^\alpha. \quad (2.8)$$

REMARK. Relation (2.8) is obtained as in the classical theory if A is continuous.

LEMMA 2.1. Given $p \in \mathbb{N}$, there exists $q \in \mathbb{N}$ and there exists $K > 0$ such that

$$\|[(A+\epsilon)^{-1}]^\alpha \phi\|_p \leq K \epsilon^{-\operatorname{Re} \alpha} \|\phi\|_q, \text{ for all } \epsilon > 0 \text{ and for all } \phi \in F.$$

PROOF. The case $0 < \operatorname{Re} \alpha < 1$ is immediate from the integral expression for $[(A+\epsilon)^{-1}]^\alpha$, which yields

$$\|[(A+\epsilon)^{-1}]^\alpha \phi\|_p \leq \frac{|\sin \alpha \pi|}{\pi} M_p \|\phi\|_{m(p)} \epsilon^{-\operatorname{Re} \alpha} \int_0^{+\infty} \frac{t^{-\operatorname{Re} \alpha}}{1+t} dt. \quad (2.9)$$

The general case is reduced to the former by repeatedly applying (2.9) after writing

$$[(A+\epsilon)^{-1}]^\alpha = ([(A+\epsilon)^{-1}]^{\alpha/n})^n$$

with a natural number n such that $0 < \operatorname{Re} \alpha/n < 1$.

PROPOSITION 2.3. Given $\phi \in D(A^\alpha)$, $n \in \mathbb{N}$ and $\epsilon > 0$, the integral

$$\int_0^1 t^n (A+\epsilon t)^{\alpha-(n+1)} \phi dt$$

(which may be improper at $t=0$) converges and, furthermore,

$$\begin{aligned} A^\alpha \phi - \sum_{p=0}^n (-1)^p \epsilon^p \binom{\alpha}{p} (A+\epsilon)^{\alpha-p} \phi &= \\ &= (-1)^{n+1} \epsilon^{n+1} \binom{\alpha}{n} (\alpha-n) \int_0^1 t^n (A+\epsilon t)^{\alpha-(n+1)} \phi dt \end{aligned} \quad (2.10)$$

where by $(A+\epsilon)^{\alpha-p}$ we mean $[(A+\epsilon)^{-1}]^p (A+\epsilon)^\alpha$.

PROOF. We argue as in the corresponding Proposition of [1]: first we show that the function h of Proposition 2.2, is indefinitely differentiable, then we write the Taylor expansion of $h(n)$ with respect to $h(\epsilon)$. In the resulting expression we find that we can take the limit for n tending to zero, which leads to (2.10). Only slight changes must be made in order to make up for the differences between Banach and Fréchet spaces, so we omit the details.

3. MAIN RESULTS.

LEMMA 3.1. Given $\alpha, \beta \in \mathbb{C}$, $n \in \mathbb{N}$ and $\epsilon > 0$, with $n+1 > \operatorname{Re} \alpha$, $\operatorname{Re} \beta > 0$ and $\phi \in D[(A+\epsilon)^\alpha]$, we have

$$\int_0^1 t^n (A+\epsilon t)^{\alpha-(n+1)} \phi dt \in D[(A+\epsilon)^\beta]$$

and

$$\lim_{\epsilon \rightarrow 0} \epsilon^{n+1} (A+\epsilon)^\beta \int_0^1 t^n (A+\epsilon t)^{\alpha-(n+1)} \phi dt = 0$$

PROOF. As the operator $(A+\epsilon)^\beta$ is closed, in order to prove that the integral belongs to its domain, it suffices to show that this fact is true for the integrand, and that the following integral (which is improper at $t = 0$ if $0 \notin p(A)$)

$$\int_0^1 t^n (A + \varepsilon)^\beta (A + \varepsilon t)^{\alpha-(n+1)} \phi dt \quad (3.1)$$

is convergent.

The first assertion is obvious, since $n + 1 > \operatorname{Re} \beta$.

We will show that (3.1) converges with respect to any seminorm $\|\cdot\|_p$. To this end, we write

$$(A + \varepsilon)^\beta (A + \varepsilon t)^{\alpha-(n+1)} \phi = [(A + \varepsilon)^{-1}]^{n+1-\beta} \psi$$

with

$$\psi = (A + \varepsilon)^{n+1} (A + \varepsilon t)^{\alpha-(n+1)} \phi.$$

Then, by Lemma 2.1, there exists $q \in \mathbb{N}$ and $K > 0$ such that

$$\|[(A + \varepsilon)^{-1}]^{n+1-\beta} \psi\|_p \leq K \|\psi\|_q \varepsilon^{\operatorname{Re} \beta - (n+1)}$$

In order to estimate $\|\psi\|_q$ by means of a suitable function of t we use the following decomposition:

$$\begin{aligned} (A + \varepsilon)^{n+1} [(A + \varepsilon t)^{-1}]^{n+1} (A + \varepsilon t)^\alpha \phi &= \\ &= \sum_{i=0}^{n+1} \binom{n+1}{i} \varepsilon^i (1-t)^i [(A + \varepsilon t)^{-1}]^i (A + \varepsilon t)^\alpha \phi. \end{aligned}$$

Applying Lemma 2.1 to the last term of the sum and considering the continuity of $(A + \varepsilon t)^\alpha \phi$ as t ranges over the interval $[0, 1]$, we conclude that there exists H_1 and H_2 , positive and independent of t , such that

$$\|\psi\|_q \leq H_1 \sum_{i=0}^n (1-t)^i t^{-i} + H_2 \varepsilon^{\operatorname{Re} \alpha} (1-t)^{n+1} t^{\operatorname{Re} \alpha - (n+1)}.$$

This implies convergence of the integral (3.1). Besides, the estimation of the integrand also implies that

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{n+1} (A + \varepsilon)^\beta \int_0^1 t^n (A + \varepsilon t)^{\alpha-(n+1)} \phi dt = 0$$

as claimed.

LEMMA 3.2. If P is a polynomial with complex coefficients and such that all its roots are non-zero, then if $\phi \in F$ satisfies

$$P[(A + \varepsilon)^{-1}] \phi \in D[(A + \varepsilon)^\alpha]$$

we have $\phi \in D[(A + \varepsilon)^\alpha]$.

PROOF. Let r_1, r_2, \dots, r_h be the roots of P , and k_1, k_2, \dots, k_h their respective multiplicities. Then the hypothesis can be rewritten as

$$[(A + \varepsilon)^{-1} - r_1] \psi \in D[(A + \varepsilon)^\alpha] \quad (3.2)$$

with

$$\psi = [(A + \varepsilon)^{-1} - r_1]^{k_1-1} \prod_{2 \leq i \leq h} [(A + \varepsilon)^{-1} - r_i]^{k_i} \phi$$

and, if we show that $\psi \in D[(A+\varepsilon)^\alpha]$, the argument can be repeated until we arrive at the element ϕ .

Taking a natural number m such that $2^m > \operatorname{Re} \alpha$, (3.2) implies that

$$\sum_{p=0}^{m-1} \left[(A+\varepsilon)^{-1} \right]^{2^p} \left[r_1^{2^p} \right] \left[(A+\varepsilon)^{-1} - r_1 \right] \psi \in D[(A+\varepsilon)^\alpha].$$

Now, carrying out the operation on the left-hand side, we get

$$\left[(A+\varepsilon)^{-1} \right]^{2^m} \psi - r_1^{2^m} \psi \in D[(A+\varepsilon)^\alpha].$$

Since $2^m > \operatorname{Re} \alpha$, we have

$$\left[(A+\varepsilon)^{-1} \right]^{2^m} \psi \in D[(A+\varepsilon)^\alpha],$$

and we obtain $\psi \in D[(A+\varepsilon)^\alpha]$ as claimed.

PROPOSITION 3.1. If $\operatorname{Re} \alpha > 0$ and $\operatorname{Re} \beta > 0$, the following equality holds.

$$A^\beta A^\alpha = A^{\alpha + \beta}.$$

PROOF. We shall see that the operator $A^{\alpha + \beta}$ is an extension of $A^\beta A^\alpha$. Let $\phi \in D(A^\alpha)$ such that $A^\alpha \phi \in D(A^\beta)$. According to (2.10) we have for all $n \in \mathbb{N}$ and for all $\varepsilon > 0$:

$$\begin{aligned} A^\alpha \phi &= \sum_{p=0}^n (-1)^p \varepsilon^p \binom{\alpha}{p} (A+\varepsilon)^{\alpha-p} \phi + \\ &+ (-1)^{n+1} \varepsilon^{n+1} \binom{\alpha}{n} (\alpha-n) \int_0^1 t^n (A+\varepsilon t)^{\alpha-(n+1)} \phi dt. \end{aligned} \quad (3.3)$$

From here, taking $n+1 > \operatorname{Re} \alpha$, $\operatorname{Re} \beta$ and applying Lemmas 3.1 and 3.2 it follows that $(A+\varepsilon)^\alpha \phi \in D[(A+\varepsilon)^\beta]$, that is $\phi \in D(A^{\alpha + \beta})$.

In order to prove that $A^\beta A^\alpha \phi = A^{\alpha + \beta} \phi$ we let $(A+\varepsilon)^\beta$ operate on both sides of (3.3) and take limits for ε tending to zero. By Lemma 3.1 we know that

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{n+1} (A+\varepsilon)^\beta \int_0^1 t^n (A+\varepsilon t)^{\alpha-(n+1)} \phi dt = 0$$

so, in order to establish the equality of $A^\beta A^\alpha \phi$ and $A^{\alpha + \beta} \phi$, it will suffice to prove that

$$\lim_{\varepsilon \rightarrow 0} (A+\varepsilon)^\beta (\varepsilon^i (A+\varepsilon)^{\alpha-i} \phi) = 0 \quad (1 < i < n) \quad (3.4)$$

To this end, let δ be such that $0 < \delta < \min \{1, \operatorname{Re}(\alpha+\beta)\}$. Using Lemma 2.1, we have for all $p \in \mathbb{N}$,

$$\begin{aligned} \left\| (A+\varepsilon)^\beta (A+\varepsilon)^{\alpha-i} \phi \right\|_p &= \left\| \left[(A+\varepsilon)^{-1} \right]^{i-\delta} (A+\varepsilon)^{\alpha+\beta-\delta} \phi \right\|_p \\ &\leq K \varepsilon^{\delta-i} \left\| (A+\varepsilon)^{\alpha+\beta-\delta} \phi \right\|_q, \end{aligned}$$

where K and q do not depend on ϵ and the term within the seminorm is continuous in a neighborhood of zero owing to the existence of the limit

$$\lim_{\epsilon \rightarrow 0} (A + \epsilon)^{\alpha + \beta - \delta} \phi = A^{\alpha + \beta - \delta} \phi$$

(This limit exists because $\phi \in D[(A + \epsilon)^{\alpha + \beta}]$, a set which is included in $D[(A + \epsilon)^{\alpha + \beta - \delta}] = D(A^{\alpha + \beta - \delta})$). Thus we have proved (3.4), as intended.

Next we shall see that $A^\beta A^\alpha$ is an extension of $A^{\alpha + \beta}$.

Let $\phi \in D(A^{\alpha + \beta})$. Then $\phi \in D[(A + \epsilon)^{\alpha + \beta}]$ and consequently $\phi \in D(A^\alpha)$ and $A^\alpha \phi$ can be expressed as in (3.3). Now we argue as in the proof of the inclusion $A^{\alpha + \beta} \supset A^\beta A^\alpha$, except that it is now the second member of (3.3), term by term, of which we know beforehand that it belongs to $D[(A + \epsilon)^\beta]$. Thus we have $A^\alpha \phi \in D[(A + \epsilon)^\beta]$ and taking limits we obtain

$$A^\beta A^\alpha \phi = A^{\alpha + \beta} \phi.$$

PROPOSITION 3.2. A^α is a closed operator.

PROOF. Let $\{\phi_p\}_{p \geq 1}$ be a sequence in $D(A^\alpha)$ such that the following limits exist:

$$\begin{aligned} \lim_{p \rightarrow \infty} \phi_p &= \phi \\ \lim_{p \rightarrow \infty} A^\alpha \phi_p &= \psi \end{aligned} \tag{3.5}$$

Let $n \in \mathbb{N}$ such that $n > \operatorname{Re} \alpha$ and $\mu > 0$. Since A^α is an extension of the closable operator J^α and $D(J^\alpha) \supset D(A^n)$, we have

$$A^\alpha [(A + \mu)^{-1}]^n \phi = [(A + \mu)^{-1}]^n \psi \tag{3.6}$$

Consequently, as by additivity (Proposition 3.1) we know that $D(A^{n-\alpha}) \supset D(A^n)$, we arrive at

$$A^\alpha [(A + \mu)^{-1}]^n \phi \in D(A^{n-\alpha})$$

and

$$A^n [(A + \mu)^{-1}]^n \phi = A^{n-\alpha} [(A + \mu)^{-1}]^n \psi$$

from which it follows, repeating the argument,

$$A^n [(A + \mu)^{-1}]^n \phi \in D(A^\alpha)$$

Now this implies, by Lemma 3.2 that $\psi \in D(A^\alpha)$ and taking this into (3.6), it follows that

$$A^\alpha \phi = \psi,$$

thus A^α is closed.

The following Proposition can be proved by means of the techniques developed in [1] and [2].

PROPOSITION 3.3. We have. (i) When A is not continuous, A^α is an extension of the closure operator J^α , and the equality holds if and only if A is densely defined.

(ii) If μ belongs to the non-empty set

$$\{\mu \in \mathbb{C} \sim \{0\}: \mu + \lambda^\alpha e^{i\theta\alpha} \neq 0 \text{ with } \lambda > 0, -\pi \leq \theta \leq \pi\}$$

then $A^\alpha + \mu$ has a continuous inverse given by the convergent integral

$$\frac{1}{2\pi i} \int_0^+ \left[\frac{1}{\mu + \lambda^\alpha e^{-i\pi\alpha}} - \frac{1}{\mu + \lambda^\alpha e^{i\pi\alpha}} \right] (A + \lambda)^{-1} d\lambda.$$

(iii) If $0 < \alpha < 1$, then A^α is non-negative and we have

$$(A^\alpha)^\beta = A^{\alpha\beta}, \text{ for all } \beta \text{ with } \operatorname{Re} \beta > 0.$$

REMARK 3.1. The concept of a non-negative operator, as well as the results in this paper, can easily be generalized to complete locally convex spaces.

REFERENCES

1. MARTINEZ, C., SANZ, M., MARCO, M., Fractional powers of operators, J. Math. Soc. of Japan (1988), 40(2).
2. BALAKRISHNAN, A.V., Fractional powers of closed operators and the semigroups generated by them, Pacific J. Math. 10 (1960), 419-437 (MR 22, 5899).
3. LAMB, W., Fractional powers of operators defined on a Fréchet space, Proc. Edinburgh Math. Soc. 27, (1984), 165-180.
4. FATTORINI, H.O., The Cauchy problem. Encyclopedia of Mathematics and its applications. Vol. 18, Addison-Wesley, Reading, Mass. (1983).
5. FATTORINI, H.O., Second-Order linear Differential Equations in Banach spaces, North-Holland (1985).
6. HOVEL, H.W. and WESTPHAL, U., Fractional powers of closed operators, Studia Math. 42, (1972), 177-194 (MR 46, 2481).
7. KOMATSU, H., Fractional powers of operators, Pacific J. Math. 19, (1966), 285-346 (MR 34, 1862).
8. KRASNOSEL'SKII, M.A. and SOBOLEVSKII, P.E., Fractional powers of operators acting in Banach spaces, Doklady Acad. Nauk SSSR 129, (1959), 499-502.
9. MCBRIDE, A.C. and ROACH, G.F., Fractional calculus, Research notes in Mathematics 138, Pitman, Advanced Publishing Program 38 (1985).
10. WATANABE, J., On some properties of fractional powers of linear operators, Proc. Japan Acad. 37, (1961), 273-275.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br