

REGULARITY OF "F" METHOD OF SUMMABILITY OF SEQUENCES

YU CHUEN WEI

Department of Mathematics
Castleton State College
Castleton, VT 05735

(Received November 11, 1987 and in revised form January 12, 1988)

ABSTRACT. This paper is to develop theorems concerning the REGULARITY of the method "F" which is more general than Cesaro's, Able's and Riemann's methods in the theory of summability.

KEY WORDS AND PHRASES. Regularity of a Method of Summability of Sequences.

1980 AMS SUBJECT CLASSIFICATION CODE. 40C15.

1. INTRODUCTION.

There are many well-known methods in the theory of summability which has many uses throughout analysis and applied mathematics, for example, Cesaro's, Able's, Riemann's, etc.. Mathematicians have contributed much to the study of these methods which all can be found in books that provide an introduction to Summability Theory. The "F" method is one of methods of Summability, and more general than those mentioned above. But there is less information about Regularity available covering the research in this method. This note concerns the regularity of the "F" method. Five theorems will be given.

2. MAIN RESULTS.

DEFINITION 2.1. Suppose that $\{F_n(x)\}_{n=1}^{\infty}$ is a sequence of functions defined in an interval $0 < x \leq b$ and that for each n

$$\lim_{x \rightarrow 0} F_n(x) = 1,$$

and suppose that

$$F(x) = \sum a_n F_n(x)$$

is convergent in some interval $0 < x \leq c < b$ and

$$\lim_{x \rightarrow 0} F(x) = S.$$

Then we say that $\sum a_n$ is summable (F) to S.

It is not hard to see that if $\sum a_n$ is Cesaro or Abel or Riemann summable, then $\sum a_n$ is summable (F) for suitable functions $F_n(x)$, respectively.

There is a well-known theorem about the regularity of the "F" method. [1,2]:

THEOREM. (REGULARITY) In order that the "F" method should be regular, it is necessary and sufficient that

$$\sum |F_n(x) - F_{n+1}(x)| < H, \quad (2.1)$$

where H is independent of x, in some interval $0 < x \leq c < b$.

It is clear that method "F" is regular if $\{F_n(x)\}$ is monotone and uniformly bounded in some interval $0 < x \leq c < b$. Next first theorem will prove that the "F" method should be regular for some sequence of functions $\{F_n(x)\}$ without monotonicity.

THEOREM 2.1. The condition (2.1) is satisfied if there are two positive sequences $\{m_n\}_{n=0}^{\infty}$ and $\{M_n\}_{n=0}^{\infty}$ such that

$$0 < m_n < m_{n+1}, \quad \text{for all } n$$

and

$$0 < M_n, \quad \sum_{n=0}^{\infty} M_n < \infty$$

and for each n

$$|F_n(x) - 1| < M_n x^{m_n}, \quad 0 < x \leq c < b.$$

PROOF. Since

$$\begin{aligned} & |F_n(x) - F_{n+1}(x)| \\ & \leq |F_n(x) - 1| + |F_{n+1}(x) - 1| \\ & \leq M_n x^{m_n} + M_{n+1} x^{m_{n+1}}, \end{aligned}$$

we have

$$\begin{aligned} & \sum |F_n(x) - F_{n+1}(x)| \\ & \leq \sum (M_n x^{m_n} + M_{n+1} x^{m_{n+1}}) \\ & = M_0 x^{m_0} + 2 \sum_{n=1}^{\infty} M_n x^{m_n}. \end{aligned}$$

If $0 < x < r < 1$, then $x^{m_n} \geq x^{m_{n+1}} \geq \dots > 0$, and for any N

$$\left| \sum_{n=0}^N M_n \right| \leq A. \quad (A \text{ is a constant})$$

For each such x, by Abel's inequality

$$\left| \sum_{n=1}^N M_n x^{m_n} \right| \leq Ax^{m_1} \leq Ar^{m_1}$$

for any N . Let $N \rightarrow \infty$, we have

$$\left| \sum_{n=1}^{\infty} M_n x^{m_n} \right| \leq Ar^{m_1} .$$

Thus

$$\sum |F_n(x) - F_{n+1}(x)| \leq M_0 r^{m_0} + 2 Ar^{m_1} .$$

Let $H = M_0 r^{m_0} + 2Ar^{m_1} + 1$. H is independent of x and

$$\sum |F_n(x) - F_{n+1}(x)| < H$$

in some interval $0 < x \leq c < b$.

THEOREM 2.2. Suppose "F" is regular. Then $\sum a_n F_n(x)$ convergent implies that $\sum a_n F_n^2(x)$ convergent.

PROOF. It follows from the regularity of "F" and $\lim_{x \rightarrow 0} F_n(x) = 1$ for each n , that

$$\sum |F_n(x) - F_{n+1}(x)| < H ,$$

and

$$|F_0(x)| < H_1, \quad |F_n(x)| < H_1 + H . \quad n = 1, \dots \quad (2.2)$$

where H, H_1 are independent of x , in some interval $0 < x \leq c < b$. $\sum a_n F_n(x) = F(x)$, for any $\epsilon > 0$ and each x , we can choose $N_0(\epsilon, x)$, such that

$$\left| \sum_{n=0}^{N-1} a_n F_n(x) - F(x) \right| < \epsilon ,$$

$$N > N_0(\epsilon, x) + 1 .$$

Let

$$S_n = \sum_{i=0}^n a_i F_i(x) ,$$

also

$$\begin{aligned} \left| \sum_{N}^P a_n F_n^2(x) \right| &= \left| \sum_{N}^P a_n F_n(x) F_n(x) \right| \\ &= \left| \sum_{N}^P (S_n(x) - S_{n-1}(x)) F_n(x) \right| \\ &= \left| \sum_{N}^P [(S_n(x) - F(x)) + (F(x) - S_{n-1}(x))] F_n(x) \right| \\ &< |F(x) - S_{N-1}(x)| |F_N(x)| + \sum_{N}^{P-1} [|S_n(x) - F(x)| |F_n(x) - F_{n+1}(x)|] \\ &\quad + |(S_P(x) - F(x)) | |F_P(x)| . \end{aligned}$$

For $P > N > N_0(\epsilon, x) + 1$, it follows

$$\left| \sum_{n=1}^p a_n F_n^2(x) \right| < \varepsilon (H_1 + H) + \varepsilon H + \varepsilon (H_1 + H) = \varepsilon (2H_1 + 3H),$$

Therefore, $\sum a_n F_n^2(x)$ is a convergent in the interval $0 < x \leq c < b$.

COROLLARY 2.1. Suppose that "F" is regular, then $\sum a_n F_n(x)$ convergent implies that $\sum a_n F_n^m(x)$ convergent, where m is a positive integer.

PROOF. It follows from Theorem 2.2 that $m = 2$ the assertion is true. Suppose that $\sum a_n F_n^k(x)$ is convergent and

$$\sum a_n F_n^k(x) = G(x),$$

and let

$$S_n(x) = \sum_{i=0}^n a_i F_i^k(x),$$

then

$$\begin{aligned} \left| \sum_{n=1}^p a_n F_n^{k+1}(x) \right| &= \left| \sum_{n=1}^p a_n F_n^k(x) F_n(x) \right| \\ &= \left| \sum_{n=1}^p (S_n(x) - S_{n-1}(x)) F_n(x) \right| \\ &= \left| \sum_{n=1}^p [(S_n(x) - G(x)) + (G(x) - S_{n-1}(x))] F_n(x) \right|. \end{aligned}$$

Repeating the procedure of the proof of Theorem 2.2, the convergence of $\sum a_n F_n^{k+1}(x)$ can be proved. By the Axiom of Mathematical Induction, for all positive integer m the assertion of the corollary is true.

THEOREM 2.3. Suppose that "F" is regular, then "F^m" is regular, where m is a positive integer.

PROOF. Since

$$\begin{aligned} &\left| \frac{m}{F_n(x)} - \frac{m}{F_{n+1}(x)} \right| \\ &= \left| \frac{m}{F_n(x)} - \frac{m}{F_{n+1}(x)} \right| \left| F_n^{m-1}(x) + F_n^{m-2}(x) F_{n+1}(x) + \dots + F_{n+1}^{m-1}(x) \right|, \end{aligned}$$

it follows from (2.2) that

$$\left| \frac{m}{F_n(x)} - \frac{m}{F_{n+1}(x)} \right| < m(H_1 + H)^{m-1} \left| F_n(x) - F_{n+1}(x) \right|,$$

and

$$\begin{aligned} &\sum \left| \frac{m}{F_n(x)} - \frac{m}{F_{n+1}(x)} \right| \\ &< m(H_1 + H)^{m-1} \sum \left| F_n(x) - F_{n+1}(x) \right|. \end{aligned}$$

Hence for any positive integer m , if "F" is regular, then "F^m" is regular also.

THEOREM 2.4. Suppose that methods "F" and "G" are regular, then

- 1) " $F \pm G$ " are regular;
- 2) " FG " is regular;
- 3) " F^{-1} " is regular, if $\inf_{\substack{0 < x \leq c \\ \text{all } n}} (F_n(x)) \geq e \neq 0$.

PROOF. It follows from (2.1) and (2.2) that

$$\begin{aligned}
 1) \quad & \sum |(F_n \pm G_n) - (F_{n+1} \pm G_{n+1})| \\
 & \leq \sum |(F_n - F_{n+1}) \pm (G_n - G_{n+1})| \\
 & < \sum [|F_n - F_{n+1}| + |G_n - G_{n+1}|] \\
 & < H_F + H_G, \\
 2) \quad & |F_n G_n - F_{n+1} G_{n+1}| \\
 & = |F_n G_n - F_{n+1} G_n + F_{n+1} G_n - F_{n+1} G_{n+1}| \\
 & \leq |F_n G_n - F_{n+1} G_n| + |F_{n+1} G_n - F_{n+1} G_{n+1}| \\
 & \leq |F_n - F_{n+1}| |G_n| + |G_n - G_{n+1}| |F_{n+1}| \\
 & < H_G^1 |F_n - F_{n+1}| + H_F^1 |G_n - G_{n+1}|
 \end{aligned}$$

and

$$\sum |F_n G_n - F_{n+1} G_{n+1}| < H_F H_G^1 + H_F^1 H_G,$$

$$3) \quad |F_n^{-1} - F_{n+1}^{-1}| = \frac{|F_{n+1} - F_n|}{|F_{n+1}| |F_n|} \leq \frac{|F_{n+1} - F_n|}{e^2},$$

and

$$\begin{aligned}
 \sum |F_n^{-1} - F_{n+1}^{-1}| & \leq \sum \frac{|F_{n+1} - F_n|}{e^2} \\
 & \leq \frac{1}{e^2} \sum |F_{n+1}(x) - F_n(x)| \\
 & < \frac{1}{e^2} H_F,
 \end{aligned}$$

where H_F , H_G , H_F^1 , and H_G^1 are independent of x , in some interval $0 < x \leq c < b$. Therefore, the assertions 1), 2) and 3) are true. The proof is completed.

REFERENCES

1. GOLDBERG, R. Methods of Real Analysis, John Wiley and Sons, Inc., New York, 1976.
2. HARDY, G.H. Divergent Series, Oxford University Press, 1949.
3. POWELL, R.E. and SHAH, S.M. Summability Theory and its Applications, Van Nostrand Reinhold Co., London, 1972.
4. KNOPP, K. Theory and Application of Infinite Series, English translation by R.C. Young, Blackie, London, 1928.

Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Lead Guest Editor

Juan J. Nieto, Departamento de Análisis Matemático,
Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;
juanjose.nieto.roig@usc.es

Guest Editor

Donal O'Regan, Department of Mathematics, National
University of Ireland, Galway, Ireland;
donal.oregan@nuigalway.ie