

NOTES ON SUFFICIENT CONDITIONS FOR A GRAPH TO BE HAMILTONIAN

MICHAEL JOSEPH PAUL

School of Computer Science
Florida International University
Miami, Florida 33199

CARMEN BAYTAN SHERSHIN

Mathematics Department
Ransom-Everglades School
Coconut Grove, Florida 33133

and

ANTHONY CONNORS SHERSHIN

Mathematics Department
Florida International University
Miami, Florida 33199

(Received October 1, 1990 and in revised form December 8, 1990)

ABSTRACT. The first part of this paper deals with an extension of Dirac's Theorem to directed graphs. It is related to a result often referred to as the Ghouila-Houri Theorem. Here we show that the requirement of being strongly connected in the hypothesis of the Ghouila-Houri Theorem is redundant.

The Second part of the paper shows that a condition on the number of edges for a graph to be hamiltonian implies Ore's condition on the degrees of the vertices.

KEY WORDS AND PHRASES. hamilton cycle, directed graphs, Ghouila-Houri Theorem, strongly connected digraphs, Dirac's Theorem, Ore's Theorem, edge condition.

1980 AMS SUBJECT CLASSIFICATION CODES. 05C45, 05C20.

1. EXTENDING DIRAC'S THEOREM TO DIRECTED GRAPHS

Establishing whether or not a graph is hamiltonian is nontrivial. Unfortunately, no elegant (convenient) characterization of hamiltonian graphs exists, although several necessary or sufficient conditions are known [1]. Sufficient conditions for a graph, or digraph, to have a hamilton cycle usually take the form of implicitly requiring many edges. One such digraph result was conjectured by C. Berge and was proved by M. Alain Ghouila-Houri [2]. We shall refer to this as the G-H theorem. The hypothesis of this theorem, in addition to requiring many arcs, also stipulates that the digraph be strongly connected. One common variation of the G-H theorem extends Dirac's well known sufficient condition, for hamilton graphs, to directed graphs with the significant additional condition that such digraphs be strongly connected. In the literature this variation has sometimes been called Ghouila-Houri's Theorem [3] even though it is only an immediate corollary of the result stated in [2].

The terms used in this paper are consistent with those used by Chartrand [1]. Only simple graphs and digraphs are considered (i.e. those with no loops, parallel edges or parallel arcs).

In digraph D let $id(v)$ be the indegree of vertex v , $od(v)$ be the outdegree of v , and p be the order (i.e., the number of vertices in D).

1.1 G-H Theorem. If D is a strongly connected simple digraph and if $\text{id}(v) \geq p/2$ and $\text{od}(v) \geq p/2$ for all vertices v in D , then D has a hamilton cycle. [2]

To eliminate the redundancy in the hypothesis of the G-H Theorem, we prove that only weak connectivity need be established for the G-H theorem to hold (as opposed to strong connectivity).

1.2. Lemma. If D is a simple weakly connected digraph, and if for each vertex v , $\text{id}(v) \geq p/2$ and $\text{od}(v) \geq p/2$, then D is strongly connected.

Proof. We shall establish the contrapositive. Suppose D is not strongly connected. Then there are vertices u and v such that there is no dipath from u to v . Partition the vertices of D into two sets A and B where A contains v and all vertices which have a dipath to v , and B contains all the vertices not in A . Since u is in B , both sets are nonempty.

Denote the cardinality of A and B by $|A|$ and $|B|$, respectively. Suppose $|A| \leq |B|$. Thus $|A| \leq p/2$ and $\text{id}(v)$ is at most $(p/2)-1$. Now suppose $|A| > |B|$. This implies that $|B| \leq (p/2)-1$. The vertex u cannot be adjacent to any vertex in A for, if so, it would be in A . (Moreover, u cannot be adjacent to itself since D is simple.) Therefore, u can only be adjacent to the other vertices in B ; that is, $\text{od}(u) < (p/2)-1$.

Thus we have proved that if D is not strongly connected, there is some vertex w such that either $\text{id}(w) < p/2$ or $\text{od}(w) < p/2$. \square

Consequently, Dirac's theorem does generalize in an easy, immediate fashion to digraphs. This result also follows from a stronger result due to Zhang Cun-Quan [4]. Moreover, since verifying that a digraph is strongly connected is nontrivial, this relaxation of the G-H Theorem becomes a useful tool to determine easily whether a digraph is hamiltonian or not.

The 1.2 Lemma is also significant in its own right because verifying that a digraph is strongly connected is an important matter. Indeed, this result, when the condition is satisfied, is easier to use when determining if a large graph is strongly connected than two other commonly used techniques: (a) computing the reachability matrix from the adjacency matrix and seeing whether or not it consists entirely of one's [5]; (b) using the depth-first search algorithm due to R.E. Tarjan [6].

We now state the refinement of (1.1) which follows immediately from it and (1.2).

1.3 Theorem. If D is a simple connected digraph and if for each vertex v , $\text{id}(v) \geq p/2$ and $\text{od}(v) \geq p/2$, then D has a hamilton cycle.

Lastly, the original Ghouila-Houri Theorem in [5] states that if D is strongly connected and if $\text{id}(v) + \text{od}(v) \geq p$ for every vertex v , then D has a hamilton cycle. However, this condition that the minimum total degree be at least p does not guarantee that D is strongly connected: Consider a digraph consisting of two complete symmetric digraphs D and K , with $|D|$ and $|K|$ at least 2, such that D and K are joined completely by all possible arcs from the vertices of D to the vertices of K . This example is mentioned in [7].

2. A FAMILIAR RESULT REVISITED.

Several sufficient conditions for a graph to be hamiltonian have been established. The well known Dirac's condition and Ore's condition deal with the degree of the vertices of a graph. Another condition, less quoted, guarantees a hamilton cycle when there are "enough" edges [8]. Let G be a simple graph having p vertices and m edges. Each of the following is sufficient for a graph to be hamiltonian:

Dirac's condition: For each vertex v of G , $\deg v \geq p/2$.

Ore's Condition: For each pair of nonadjacent vertices u and v of G , $\deg u + \deg v \geq p$.

Edge condition: The number of edges $m \geq (1/2)(p-1)(p-2) + 2$.

The purpose of this section is to show that the edge condition implies Ore's condition, to give simple examples to show that no implication exists between Dirac's condition and the edge condition, and to show that Ore's condition does not necessarily imply either Dirac's or the edge conditions:

The set diagram below (Figure 1) summarizes the relationships among hamilton graphs satisfying Ore's (O), Dirac's (D), and/or the edge condition (E).




Figure 1 Venn diagram summarizing the relationships among hamilton graphs satisfying Ore's, Dirac's, and Edge conditions.

2.1 Theorem. The edge condition implies Ore's condition.

Proof. Let G be a simple graph satisfying the edge condition and let u and v be nonadjacent vertices. The maximum number of possible edges is $(1/2)(p-2)(p-3) + \deg u + \deg v$. Therefore we get $(1/2)(p-2)(p-3) + \deg u + \deg v \geq m \geq (1/2)(p-1)(p-2) + 2$. This yields $\deg u + \deg v \geq (1/2)[(p-1)(p-2) - (p-2)(p-3)] + 2 = p$. \square

Every graph theory student learns that Dirac's condition implies Ore's condition but that the converse is not true. Now we will give simple examples to show that no implication exists between Dirac's condition and the edge condition: A cycle of length four satisfies Dirac's condition, but not the edge condition; the graph of order 5, consisting of K_4 with the addition of a vertex of degree 2 joining it to any two vertices of K_4 , satisfies the edge condition but not Dirac's; a complete graph of order ≥ 3 satisfies both Dirac's and the edge condition. Finally, Figure 2 gives an example of a graph that satisfies Ore's condition (O) but does not satisfy Dirac's (D) nor the edge conditions (E).




Figure 2 A graph in the set $O - (D \cup E)$

ACKNOWLEDGEMENT. The authors thank the referee for suggesting the graph pictured in Figure 2.

REFERENCES

- [1] CHARTRAND, G., Introductory Graph Theory, Dover, 1985.
- [2] GHOUILA-HOURI, M.A., Une condition suffisante d'existence d'un circuit hamiltonien, C.R. Acad. Sci. Paris 251, 1960, 495-497.
- [3] WILSON, R., Introduction to Graph Theory, Longman, 1972, 107.
- [4] CUN-QUAN, Z., Discrete Math. 41, 1982, 79-96.
- [5] HARARY, F., NORMAN, R., and CARTWRIGHT, D., Structural Models: An introduction to the Theory of Directed Graphs, Wiley, 1965, 137.
- [6] AHO, A., HOPCROFT, J., and ULLMAN, J., The design and Analysis of Computer Algorithms, Addison-Wesley, 1974, 189-195.
- [7] BERMOND, J.C., and THOMASSEN, C., Cycles in Digraphs - A Survey, Journal of Graph Theory 5, 1981, 1-43.
- [8] BERGE, C., Graphs and Hypergraphs, North-Holland, 1973, 212-213.

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.oter@usc.es