

FIXED POINT THEOREMS FOR d-COMPLETE TOPOLOGICAL SPACES I

TROY L. HICKS

Department of Mathematics
University of Missouri at Rolla
Rolla, Missouri 65401

(Received May 7, 1991)

ABSTRACT. Generalizations of Banach's fixed point theorem are proved for a large class of non-metric spaces. These include d-complete symmetric (semi-metric) spaces and complete quasi-metric spaces. The distance function used need not be symmetric and need not satisfy the triangular inequality.

KEY WORDS AND PHRASES. Fixed point, d-complete topological spaces.

1980 AMS SUBJECT CLASSIFICATION CODE. 47H10; 54H25

Let (X, t) be a topological space and $d : X \times X \rightarrow [0, \infty)$ such that $d(x, y) = 0$ if and only if $x = y$. X is said to be d-complete if $\sum_{n=1}^{\infty} d(x_n, x_{n+1}) < \infty$ implies that the sequence (x_n) is convergent in (X, t) . In a metric space, such a sequence is a Cauchy sequence. If $T : X \rightarrow X$, $0(x, \infty) = (x, Tx, T^2x, \dots)$ is called the orbit of x . $G : X \rightarrow [0, \infty)$ is T -orbitally lower semi-continuous at x^* if (x_n) is a sequence in $0(x, \infty)$ and $\lim x_n = x^*$ implies $G(x^*) \leq \liminf G(x_n)$. $T : X \rightarrow X$ is w -continuous at x if $x_n \rightarrow x$ implies $Tx_n \rightarrow Tx$.

The basic idea of a d-complete topological space goes back to Kasahara [1] and [2], Iseki [3], and their L-spaces.

LEMMA 1. X is a set, $T : X \rightarrow X$ and $d : X \times X \rightarrow [0, \infty)$. Then there exists $\phi : X \rightarrow [0, \infty)$ such that

(a) $d(x, Tx) \leq \phi(x) - \phi(Tx)$ for all $x \in X$, if and only if

(b) $\sum_{n=0}^{\infty} d(T^n x, T^{n+1} x)$ converges for all x .

(c) If (a) holds for all $y \in 0(x, \infty)$, then $\sum_{n=0}^{\infty} d(T^n x, T^{n+1} x)$ converges.

PROOF. Suppose (a) holds. For $x \in X$, $S_n = \sum_{i=0}^n d(T^i x, T^{i+1} x) \leq \sum_{i=0}^n [\phi(T^i x) - \phi(T^{i+1} x)] = \phi(x) - \phi(T^{n+1} x) \leq \phi(x)$. (S_n) is non-decreasing and bounded above and therefore convergent.

Suppose (b) holds and let $\phi(x) = \sum_{i=0}^{\infty} d(T^i x, T^{i+1} x)$. $\phi(x) - \phi(Tx) = d(x, Tx)$

since $S_n - T_n = \sum_{i=0}^n d(T^i x, T^{i+1} x) - \sum_{i=0}^n d(T^{i+1} x, T^{i+2} x) = d(x, Tx) - d(T^{n+1} x, T^{n+2} x) \rightarrow d(x, Tx)$ as $n \rightarrow \infty$.

The proof of (b) gives (c).

LEMMA 2. X is a topological space, $T : X \rightarrow X$ and $d : X \times X \rightarrow [0, \infty)$ such that $d(x, y) = 0$ if and only if $x = y$. Suppose there exists an $x \in X$ such

that $\lim T^n x = x^*$ exists. Then:

(a) $Tx^* = x^*$ implies $G(x) = d(x, Tx)$ is T -orbitally lower semi-continuous at x^* .

(b) G is T -orbitally lower semi-continuous at x^* and $\liminf d(T^n x, T^{n+1} x) = 0$ imply $Tx^* = x^*$.

PROOF. Assume that $Tx^* = x^*$ and (x_n) is a sequence in $O(x, \infty)$ with $\lim x_n = x^*$. Then $G(x^*) = d(x^*, Tx^*) = 0 \leq \liminf d(x_n, Tx_n) = \liminf G(x_n)$.

If $x_n = T^n x \rightarrow x^*$ and G is T -orbitally lower semi-continuous at x^* , then

$0 \leq d(x^*, Tx^*) = G(x^*) \leq \liminf G(x_n) = \liminf d(T^n x, T^{n+1} x) = 0$. Thus, $Tx^* = x^*$.

The next theorem is a version of Caristi's theorem [4 or 5] in this more general setting. Caristi's theorem for metric spaces is a generalization of Banach's fixed point theorem.

THEOREM 1. Let X be a d -complete topological space. Suppose $T : X \rightarrow X$ and $\phi : X \rightarrow [0, \infty)$. Suppose there exists an x such that

$$d(y, Ty) \leq \phi(y) - \phi(Ty) \text{ for all } y \in O(x, \infty). \quad (1)$$

Then we have:

(a) $\lim T^n x = x^*$ exists.

(b) $Tx^* = x^*$ if and only if $G(x) = d(x, Tx)$ is T -orbitally lower semi-continuous at x .

PROOF. From Lemma 1, $\sum_{n=0}^{\infty} d(T^n x, T^{n+1} x)$ is convergent. X is d -complete so $\lim T^n x = x^*$ exists. Note that $\lim d(T^n x, T^{n+1} x) = 0$. Now apply Lemma 2.

Even in this general setting one obtains a version of Banach's theorem as a corollary of the above theorem.

COROLLARY 1. Let X be a d -complete topological space and $0 < k < 1$. Suppose $T : X \rightarrow X$ and there exists an x such that

$$d(Ty, T^2 y) \leq k d(y, Ty) \text{ for all } y \in O(x, \infty). \quad (2)$$

Then:

(a) $\lim T^n x = x^*$ exists.

(b) $Tx^* = x^*$ if and only if $G(x) = d(x, Tx)$ is T -orbitally lower semi-continuous at x .

PROOF. Set $\phi(y) = \frac{1}{1-k} d(y, Ty)$ for $y \in 0(x, \infty)$. Let $y = T^n x$ in (2). Then $d(T^{n+1}x, T^{n+2}x) \leq k d(T^n x, T^{n+1}x)$ and $d(T^n x, T^{n+1}x) - k d(T^n x, T^{n+1}x) \leq d(T^n x, T^{n+1}x) - d(T^{n+1}x, T^{n+2}x)$. Thus, $d(T^n x, T^{n+1}x) \leq \frac{1}{1-k} [d(T^n x, T^{n+1}x) - d(T^{n+1}x, T^{n+2}x)]$ or $d(y, Ty) \leq \phi(y) - \phi(Ty)$. Apply Theorem 1.

In [6], it was shown that many generalizations of Banach's theorem hold for quasi-metric ($d(x, y) \neq d(y, x)$) spaces. However, we do not have the triangular inequality in the present setting, so standard proofs and theorems do not necessarily hold. Before proving more theorems, we give the definitions of some of the special d -complete topological spaces we had in mind when we formulated the results of this paper.

DEFINITION. A symmetric on a set X is a real-valued function d on $X \times X$ such that:

- (a) $d(x, y) \geq 0$ and $d(x, y) = 0$ if and only if $x = y$; and
- (b) $d(x, y) = d(y, x)$.

Let d be a symmetric on a set X and for any $\epsilon > 0$ and any $x \in X$, let $S(x, \epsilon) = \{y \in X: d(x, y) < \epsilon\}$. We define a topology $t(d)$ on X by $U \in t(d)$ if and only if for each $x \in U$, some $S(x, \epsilon) \subset U$. A symmetric d is a semi-metric if for each $x \in X$ and each $\epsilon > 0$, $S(x, \epsilon)$ is a neighborhood of x in the topology $t(d)$. A topological space X is said to be symmetrizable (semi-metrizable) if its topology is induced by a symmetric (semi-metric) on X .

THEOREM 2. Suppose X is a d -complete Hausdorff topological space, $T : X \rightarrow X$ is w -continuous and satisfies $d(Tx, T^2x) \leq k(d(x, Tx))$ for all $x \in X$, where $k : [0, \infty) \rightarrow [0, \infty)$, $k(0) = 0$, and k is non-decreasing. Then T has a fixed point if and only if

there exists an x in X with $\sum_{n=1}^{\infty} k^n(d(x, Tx)) < \infty$. In this case, $x_n = T^n x \rightarrow p = Tp$.

[k is not assumed to be continuous and $k^2(a) = k(k(a))$].

PROOF. If $Tp = p$, $d(p, Tp) = 0$, and $k^n(0) = 0$ for every n .

Assume the condition holds and let $x_n = T^n x$. $d(Tx, T^2x) \leq k(d(x, Tx))$ and $d(T^2x, T^3x) \leq k(d(Tx, T^2x)) \leq k^2(d(x, Tx))$. By induction,

$$d(x_n, x_{n+1}) = d(T^n x, T^{n+1} x) \leq k^n(d(x, Tx)).$$

Now $\sum_{n=1}^{\infty} d(x_n, x_{n+1}) < \infty$ and X is d -complete. Thus, $\lim x_n = p$ exists. T

w -continuous implies $x_{n+1} = Tx_n \rightarrow Tp$. Since $x_{n+1} \rightarrow p$, $Tp = p$.

In many special cases of Theorem 2, one has $d(Tx, Ty) \leq k(d(x, y))$ for all $x, y \in X$ and the special form of k forces w -continuity of T . It may also force the uniqueness of p and enable us to obtain error bounds. In other cases, T need not be w -continuous so the following theorem is needed.

THEOREM 3. Suppose X is a d -complete Hausdorff topological space, $T : X \rightarrow X$ and there exists a y such that $d(Tx, T^2x) \leq k(d(x, Tx))$ for every $x \in 0(y, \infty)$, where $k : [0, \infty) \rightarrow [0, \infty)$, $k(0) = 0$, and k is non-decreasing. Suppose $\sum_{n=1}^{\infty} k^n(d(y, Ty)) < \infty$

and (x_n) in $O(y, \infty)$ with $\sum_{n=1}^{\infty} d(x_n, x_{n+1}) < \infty$ imply $\lim x_n$ exists. Let $x_n = T^n y$.

Then:

(a) $\lim x_n = p$ exists.

(b) $Tp = p$ if and only if $G(x) = d(x, Tx)$ is T -orbitally lower semi-continuous at p .

PROOF. The proof of Theorem 2 gives (a). Lemma 2 gives (b) since

$$0 = \lim_n d(x_n, x_{n+1}) = \lim_n d(T^n y, T^{n+1} y) = 0.$$

COROLLARY 2. Let $0 < \lambda < 1$. Suppose X is a d -complete Hausdorff topological space, $T : X \rightarrow X$ is w -continuous and satisfies $d(Tx, Ty) \leq \lambda d(x, y)$ for all $x, y \in X$. Then T has a unique fixed point p . For any $x \in X$, $p = \lim T^n x$.

PROOF. Let $k(t) = \lambda t$. Then $k^n(d(x, Tx)) = \lambda^n d(x, Tx)$, so $\sum_{n=1}^{\infty} k^n(d(x, Tx)) < \infty$

for any x in X . T is w -continuous so Theorem 2 gives $x_n = T^n x \rightarrow p = Tp$ for any $x \in X$. Clearly, p is unique since $0 < \lambda < 1$.

In Corollary 2, if one replaces topological by symmetrizable, then $d(Tx, Ty) \leq \lambda d(x, y)$ forces T to be w -continuous. We now give several examples of a specific function k where Theorem 2 or Theorem 3 applies to yield another corollary similar to Corollary 2. To apply the theorems, one needs a non-decreasing function k and an x in X with $\sum_{n=1}^{\infty} k^n(d(x, Tx)) < \infty$. The following examples satisfy these

conditions. The reader can consult [7] for the details that are not obvious and not provided here.

EXAMPLE 1. Suppose $0 < \lambda < 1$. Let $k(t) = \lambda t$ for $t \geq 0$. Then $d(Tx, T^2 x) \leq k(d(x, Tx)) = \lambda d(x, Tx)$ gives $k^n(d(x, Tx)) = \lambda^n d(x, Tx)$.

EXAMPLE 2. Suppose T satisfies $d(Tx, Ty) \leq \phi(d(x, y)) d(x, y)$ for all x, y in X , where $\phi : [0, \infty) \rightarrow [0, 1)$ and ϕ is non-decreasing. Then $k(t) = t \phi(t)$, k is non-decreasing, and $k : [0, \infty) \rightarrow [0, \infty)$. It follows by induction that $k^n(t) \leq t[\phi(t)]^n$.

Since $\phi(t) < 1$, $\sum_{n=1}^{\infty} k^n(t) < \infty$.

EXAMPLE 3. Consider $k(t) = t \phi(t)$ where $\phi : [0, \infty) \rightarrow [0, \infty)$ and $\phi(t) \leq t$ for $t < 1$. If $t < 1$, it follows that $k^n(t) \leq t[\phi(t)]^n$. If k is non-decreasing, the theorems apply.

EXAMPLE 4. $k(t) = t \phi(t)$ where $\phi : [0, \infty) \rightarrow [0, \infty)$ and $\phi(at) \leq \alpha \phi(t)$ for $a \in (0, 1)$. If $\phi(t) < 1$, $k^n(t) \leq (kt)(\phi(t))^n$ for all $n \geq 2$. If k is non-decreasing, the theorems apply.

EXAMPLE 5. Assume k is non-decreasing, k is convex on $[0, 1]$, and $k(t) < t$ for all $0 < t < 1$. Fix $t < 1$. Now $k(t) < t$ gives $k(t) = at$ for some

$0 < a = a(t) < 1$. By induction, $k^n(t) \leq a^n t$ for all n and thus $\sum_{n=1}^{\infty} k^n(t) < \infty$.

THEOREM 4. Suppose (X, d) is a Hausdorff d -complete symmetrizable space,

$T : X \rightarrow X$, and $d(Tx, Ty) \leq [d(x, y)]^p$ where $p > 1$. If there exists x such that $\mu = d(x, Tx) < 1$, then $x_n = T^n x \rightarrow p = Tp$.

PROOF. Let $k(t) = t^p$ for $t \geq 0$. $k(0) = 0$, k is increasing, $k(t) < t$ if $t < 1$, and k is convex. Since $0 \leq \mu < 1$, $\sum_{n=1}^{\infty} k^n(\mu) < \infty$. T is w -continuous so Theorem 2 applies.

REMARKS. Given a topological space (X, t) , when does there exist a distance function d such that X is d -complete? Let X be an infinite set and t any T_1 non-discrete first countable topology for X . Then there exists a complete metric d for X such that $t \leq t_d$ and the metric topology t_d is non-discrete. Now (X, t) is d -complete since $\sum d(x_n, x_{n+1}) < \infty$ implies that (x_n) is a d -Cauchy sequence. Thus, $x_n \rightarrow x$ in t_d and therefore in the topology t . In [8], the construction gives $t \leq t_E$ where t_E is a complete uniform space and the uniformity has a countable base.

Hence, the uniformity is metrizable and the compatible metric d must be complete.

It should also be noted that any complete quasi-metric space (X, d) ($d(x, y) \leq d(y, x)$) is a d -complete topological space. There are several competing definitions for a Cauchy sequence, but $\sum d(x_n, x_{n+1}) < \infty$ will imply that (x_n) is a Cauchy sequence for any reasonable definition. One reasonable definition is obtained by requiring that the filter generated by (x_n) be a Cauchy filter in the quasi-uniformity generated by d . This gives (x_n) is a Cauchy sequence if for each $\epsilon > 0$ there exists a positive integer $n_0 = n(\epsilon)$ and $x = x(\epsilon)$ in X such that $(x_n : n \geq n_0) \subseteq \{y \in X : d(x, y) < \epsilon\}$. The metric space definition of a Cauchy sequence also holds if $\sum d(x_n, x_{n+1}) < \infty$.

REFERENCES

1. S. Kasahara, On some generalizations of the Banach contraction theorem, Math. Seminar Notes, 3(1975), 161-169.
2. S. Kasahara, Some fixed point and coincidence theorems in L -spaces, Math. Seminar Notes, 3(1975), 181-187.
3. K. Iseki, An approach to fixed point theorems, Math. Seminar Notes, 3(1975), 193-202.
4. Alberta Bollenbacher and T. L. Hicks, A fixed point theorem revisited, Proc. Amer. Math. Soc. 102, No. 4, (1988), 898-900.
5. J. Caristi, Fixed point theorems for mappings satisfying inwardness conditions, Trans. Amer. Math. Soc. 215(1976), 241-251.
6. T. L. Hicks, Fixed point theorems for quasi-metric spaces, Math. Japonica 33, No. 2(1988), 231-236.
7. T. L. Hicks, Another view of fixed point theory, Math. Japonica 35, No. 2 (1990), 231-234.
8. Wilson R. Crisler and Troy L. Hicks, A Note on T_1 topologies, Proc. Amer. Math. Soc. 46, No. 1, (1974), 94-96.

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be