

A GENERAL NOTION OF INDEPENDENCE OF SEQUENCES OF INTEGERS

JOHN R. BURKE

Gonzaga University
Department of Mathematics
Gonzaga University
Spokane, WA 99258-0001
USA

(Received February 4, 1992 and in revised form December 10, 1992)

ABSTRACT. In this paper a notion of statistical independence of sequences of integers is developed. The results are generalizations of known results on independent sequences mod m in the integers and more generally, independent sequences on compact sets. All that is assumed is that one has a countable partition of the integers indexed by an ordered set.

KEY WORDS AND PHRASES. Independence mod m , uniform distribution mod m .

1991 AMS SUBJECT CLASSIFICATION CODE. 10K40.

1. INTRODUCTION.

In 1940, Steinhaus and Kac [6] established the concept of independent functions on the unit square, $[0, 1] \times [0, 1]$. Since that time the notion of independence has been developed in several different settings. The independence of sequences of integers mod m was examined by Kuipers, Niederreiter and Shiue [2, 3]. Similar work was done by Kuipers and the author in the Gaussian integers [1]. In both of these situations, one of the key properties used to characterize independent sequences was the existence of a nonprincipal character on the ring structure involved. The question of what can be said about independent sequences when there is no nonprincipal character or indeed no ring structure has also been considered. Niederreiter [5] considered independence in compact spaces. In this article we examine similar questions in a more general setting. While the set \mathbb{Z}_0 , the nonnegative integers, is considered, any ordered countable set may be used.

2. DEFINITIONS AND NOTATION

Let $A = \{a_n\}_{n=0}^{\infty}$ be a sequence in \mathbb{Z}_0 , the set of nonnegative integers. Let Λ be a countable ordered indexing set. If $\mathcal{C} = \{C_{\lambda}\}_{\lambda \in \Lambda}$ is a partition of \mathbb{Z}_0 ($\mathbb{Z}_0 = \bigcup_{\lambda \in \Lambda} C_{\lambda}$, $C_{\lambda} \cap C_{\mu} = \emptyset$ if $\lambda \neq \mu$), define

$$A(C_{\lambda}, n) = \sum_{\substack{a_k \in C_{\lambda} \\ 1 \leq k \leq n}} 1 = |\{a_1, \dots, a_n\} \cap C_{\lambda}|. \text{ If for each } \lambda \in \Lambda \text{ we have that the limit}$$

$$\lim_{n \rightarrow \infty} \frac{A(C_{\lambda}, n)}{n} \text{ exists, then define}$$

$$\phi_{\lambda}(A) = \lim_{n \rightarrow \infty} \frac{A(C_{\lambda}, n)}{n}, \lambda \in \Lambda.$$

$\{\phi_{\lambda}(A)\}_{\lambda \in \Lambda}$ is the asymptotic distribution function (a.d.f) of A with respect to the partition \mathcal{C} .

Now consider two sequences A and B, both with an a.d.f. with respect to the partition \mathcal{C} . Define

$$(A, B)(C_\lambda, C_\mu; n) = \sum_{\substack{a_k \in C_\lambda, b_k \in C_\mu \\ 1 \leq k \leq n}} 1$$

A and B will be said to be acceptable if the limit $\lim_{n \rightarrow \infty} \frac{(A, B)(C_\lambda, C_\mu; n)}{n} = \phi_{\lambda, \mu}(A, B)$ exists for all $\lambda, \mu \in \Lambda$.

Definition 1.1 Let A and B be acceptable sequences with respect to a partition $\mathcal{C} = \{C_\lambda\}_{\lambda \in \Lambda}$. A and B are \mathcal{C} -independent provided that for all $\lambda, \mu \in \Lambda$ we have

$$\phi_{\lambda, \mu}(A, B) = \phi_\lambda(A) \phi_\mu(B)$$

If one chooses the set Λ to be finite then \mathcal{C} -independence is equivalent to that described by Niederreiter [5] in compact spaces. If the equivalence classes of the partition \mathcal{C} are chosen to be residue classes mod m, \mathcal{C} -independence becomes independence mod m as considered by Kuipers, Niederreiter, and Shiue [2,3].

3. MAIN RESULTS

THEOREM 2.1 Let A be a sequence which is acceptable with respect to a partition $\mathcal{C} = \{C_\lambda\}_{\lambda \in \Lambda}$. A and A are \mathcal{C} -independent if and only if $\phi_\lambda(A) = 1$ for some $\lambda \in \Lambda$.

PROOF: Suppose there is a C_λ such that $0 < \phi_\lambda(A) < 1$. If we assume A and A are \mathcal{C} -independent then, since $\phi_\lambda(A) = \phi_{\lambda, \lambda}(A, A)$,

$$0 < [\phi_\lambda(A)]^2 < \phi_\lambda(A) = \phi_{\lambda, \lambda}(A, A).$$

Thus A and A cannot be \mathcal{C} -independent.

Now suppose $\phi_\lambda(A) = 1$ for some $\lambda \in \Lambda$. Then $\phi_\mu(A) = 0$ for $\mu \in \Lambda, \mu \neq \lambda$. It follows that for $\mu, \eta \in \Lambda, \mu \neq \lambda$ or $\eta \neq \lambda$

$$\phi_{\mu, \eta}(A, A) = \phi_\mu(A) \phi_\eta(A) = 0$$

and

$$\phi_{\lambda, \lambda}(A, A) = \phi_\lambda(A) \phi_\lambda(A) = 1.$$

THEOREM 2.2 Given a sequence A, A will be \mathcal{C} -independent of B for every B such that A and B are acceptable if and only if $\phi_\lambda(A) = 1$ for some $\lambda \in \Lambda$.

PROOF: From Theorem 2.1, it follows that $\phi_\lambda(A) = 1$ for some $\lambda \in \Lambda$ in order that A and A be \mathcal{C} -independent.

If $\phi_\lambda(A) = 1$ for some $\lambda \in \Lambda$ and B is any sequence such that A and B are acceptable, we have

$$\frac{B(C_\mu, n)}{n} - \sum_{\substack{\eta \in \Lambda \\ \eta \neq \lambda}} \frac{A(C_\eta, n)}{n} \leq \frac{(A, B)(C_\lambda, C_\mu; n)}{n} \leq \frac{B(C_\mu, n)}{n}$$

Noting that

$$\lim_{n \rightarrow \infty} \sum_{\substack{\eta \in \Lambda \\ \eta \neq \lambda}} \frac{A(C_{\eta}, n)}{n} \leq \sum_{\substack{\eta \in \Lambda \\ \eta \neq \lambda}} \phi_{\eta}(A) = 1 - \phi_{\lambda}(A) = 0$$

it follows that

$$\phi_{\lambda, \mu}(A, B) = \phi_{\mu}(B) = \phi_{\lambda}(A) \phi_{\mu}(B)$$

and

$$\phi_{\eta, \mu}(A, B) = \phi_{\eta}(A) \phi_{\mu}(B) = 0 \text{ if } \eta \neq \lambda.$$

As an example of the relationship between a.d.f. and the notion of \mathcal{C} -independence we have the following result. We will consider the special case where $\Lambda = \mathbb{Z}_0$. If any other countable indexing set is used, the choice of k can be done by choosing any order preserving map $f: \Lambda \rightarrow \mathbb{Z}_0$ and letting $k = f(\lambda)$ where $\phi_{\lambda}(A) \neq 0$, and the subsequence $\{a_{k_n}\}_{n=1}^{\infty}$ of A is determined by $a_{k_n} \in C_{\lambda}$.

THEOREM 2.3 Let A and B be acceptable with respect to a partition $\mathcal{C} = \{C_i\}_{i \in \mathbb{Z}_0}$ with a.d.f. $\{\phi_i(A)\}_{i \in \mathbb{Z}_0}$ and $\{\phi_i(B)\}_{i \in \mathbb{Z}_0}$ respectively. Let $C_k \in \mathcal{C}$ be such that $\phi_k(A) \neq 0$ and let $\{a_{k_n}\}_{n=1}^{\infty}$ be

the subsequence of A such that $a_{k_n} \in C_k$. If A and B are \mathcal{C} -independent, then the sequence

$$B' = \{b_{k_n}\}_{n=1}^{\infty} \text{ has } \{\phi_i(B)\}_{i \in \mathbb{Z}_0} \text{ as its a.d.f.}$$

PROOF: For any $j \in \mathbb{Z}_0$, since $A(C_k, k_n) = n$, we have $(A, B)(C_k, C_j; k_n) = B(C_j, n)$. By the \mathcal{C} -independence of A and B we have

$$\lim_{n \rightarrow \infty} \frac{(A, B)(C_k, C_j; k_n)}{k_n} = \phi_k(A)$$

and

$$\frac{B(C_j, n)}{n} = \frac{(A, B)(C_k, C_j; k_n)}{k_n} \cdot \frac{k_n}{n}$$

we have, by letting $n \rightarrow \infty$

$$\phi_j(B') = \phi_k(A) \phi_j(B) [\phi_k(A)]^{-1} = \phi_j(B)$$

Thus we have some basic properties of \mathcal{C} -independent sequences as well as a method of obtaining sequences with a given a.d.f.. It would, of course, be gratifying to be able to generate acceptable sequences which are \mathcal{C} -independent. In this direction we have a construction following an idea of Nathanson [4].

THEOREM 2.4 Let $A = \{a_n\}_{n=1}^{\infty}$ and $B' = \{b_n\}_{n=1}^{\infty}$ be acceptable with respect to a partition

$\mathcal{C} = \{C_{\lambda}\}_{\lambda \in \Lambda}$ with a.d.f. $\{\phi_{\lambda}(A)\}_{\lambda \in \Lambda}$ and $\{\phi_{\lambda}(B')\}_{\lambda \in \Lambda}$ respectively, where Λ is $\{1, 2, \dots, m\}$ or

$\Lambda = \mathbb{Z}_0$. Then there exists a sequence $B = \{b_n\}_{n=1}^{\infty}$ such that

- A and B are \mathcal{C} -independent

ii) $\{\phi_\lambda(B')\}_{\lambda \in \Lambda}$ is the a.d.f. for B.

PROOF: Let $i_{j,k}$ denote the index of the j^{th} term of A contained in C_k . The sets $\{i_{j,k}\}_{j=1}^\infty$ partition \mathbb{Z}_0 as k runs through Λ . Define $b_n = b_{i_{j,k}} = b'_j$.

First we establish that

$$B(C_\lambda, n) = \sum_{\mu \in \Lambda} B(C_\lambda, A(C_\mu, n)) \text{ tab 80} \quad (1)$$

The relationship (1) can be seen as follows. Consider the elements b_k , $1 \leq k \leq n$ such that $b_k \in C_\lambda$. For fixed μ we count the elements $b_{i_{1,\mu}}, b_{i_{2,\mu}}, \dots, b_{i_{A(C_\mu, n), \mu}}$ (which are the terms $b'_1, \dots, b'_{A(C_\mu, n)}$) that are in C_λ . There are $B(C_\lambda, A(C_\mu, n))$ such terms. Now let μ run through Λ .

We now establish ii). From (*) we have

$$\frac{B(C_\lambda, n)}{n} = \sum_{\mu \in \Lambda} \frac{B(C_\lambda, A(C_\mu, n))}{n} = \sum_{\mu \in \Lambda} \frac{B(C_\lambda, A(C_\mu, n))}{A(C_\mu, n)} \cdot \frac{A(C_\mu, n)}{n}.$$

Letting $n \rightarrow \infty$ we have

$$\phi_\lambda(b) = \sum_{\mu \in \Lambda} \phi_\lambda(B') \phi_\mu(A) = \phi_\lambda(B') \sum_{\mu \in \Lambda} \phi_\mu(A) = \phi_\lambda(B')$$

To establish the C -independence of A and B consider $(A, B)(C_\lambda, C_\mu; n) = B(C_\mu, A(C_\lambda, n))$. Thus

$$\begin{aligned} \phi_{\lambda, \mu}(A, B) &= \lim_{n \rightarrow \infty} \frac{(A, B)(C_\lambda, C_\mu; n)}{n} = \lim_{n \rightarrow \infty} \frac{B(C_\mu, A(C_\lambda, n))}{n} \\ &= \lim_{n \rightarrow \infty} \frac{B(C_\mu, A(C_\lambda, n))}{A(C_\lambda, n)} \cdot \frac{A(C_\lambda, n)}{n} = \phi_\mu(B') \phi_\lambda(A) = \phi_\mu(B) \phi_\lambda(A). \end{aligned}$$

It would be of interest to examine how the structure of the partition C affects the independence of sequences. Perhaps a partition whose classes are determined by the number of prime factors or distinct prime factors would be of interest.

REFERENCES

1. Burke, John R. and Kuipers, L., Asymptotic Distribution and Independence of Sequences of Gaussian Integers, Sim. Stev., 50 e Jaargang (1976-77), 3-21.
2. Kuipers, L. and Niederreiter, H., Asymptotic Distribution mod m and Independence of Sequences of Integers. I, Proc. of the Jap. Ac., Vol. 50 (1974), 256-260.
3. Kuipers, L. and Shiue, J.S., Asymptotic Distribution mod m of Sequences of Integers and the Notion of Independence, Atti della Acc. Naz dei Lincei, Serie V111, Vol. XI (1970), 63-90.
4. Nathanson, M., Asymptotic Distribution and Asymptotic Independence of Sequences of Integers, Acta Math. Acad. Sci. Hungar. 29 (1977) no. 3-4, 207-218.
5. Niederreiter, H., Indépendence de suites Lecture Notes in Math., Vol. 475 Springer Berlin (1975) 54, 120-131.
6. Steinhaus, H. and Kac, M., Fonctions Indépendentes, I-VI, Fund. Math. (1936-40), 46-131.

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.oter@usc.es