

AN OPEN TOSS PROBLEM

PREM N. BAJAJ and G.R. MENDIETA

Department of Mathematics and Statistics
The Wichita State University
Wichita, KS 67208, U.S.A.

(Received January 28, 1992)

ABSTRACT. We discuss some particular cases of the following problem. Each of m persons tosses a coin. Those who get heads stay for the next round. Those who get tails are eliminated; however, if all persons get tails in a round, they move to the next round (as if each of them had got a head). The problem is to find the possibility that exactly one person is left after r rounds. All coins are considered identical but not necessarily fair.

KEY WORDS AND PHRASES. Coin tossing, depletion model.

1991 AMS SUBJECT CLASSIFICATION CODES. Primary, 60C05; Secondary, 62P99.

1. INTRODUCTION.

Numerous games are played with players forming a circle. The opening hand decides the sequence (generally clock-wise as seen from above) of players. In each game, the opening hand is either in the most favorable position or most unfavorable position. One natural way to decide the opening hand is to have each person toss a coin and eliminate those who obtain tails, the case of "all getting tails" being considered equivalent to "all getting heads". With a number of players, it is very unlikely that the issue will be decided in one toss by the m players. Hence, the process is repeated, say r times. It will be interesting to find the probability that one person comes out as a winner after r rounds.

This problem was presented by the first author, at the AMS meeting in 1988 in Providence. An interesting paper [1] by Bruss and O'Cinneide presents some aspects of this problem.

In the following sections, we take up some particular cases of the problem. We invite solution to the general problem.

Throughout p denotes the probability of obtaining a head and $q = 1 - p$ the probability of getting a tail.

2. THEOREM.

Let $P(r, m)$ denote the probability that, starting with m persons, one person wins the toss exactly after r rounds, $m \geq 2, r \geq 1$.

Let $T_s = p^s + q^s, s \geq 2$. Then

$$(a) \quad P(r, 2) = 2pq T_2^{r-1}$$
$$(b) \quad P(r, 3) = 3pq^2 T_3^{r-1} + 6p^3q^2 \sum_{\substack{i+j=r \\ i, j \geq 0, r \geq 2}} T_3^i T_2^j$$

$$(c) \quad P(r, 4) = 4pq^3 T_4^{r-1} + 12p^4 q^3 \sum_{\substack{i+j=r-2 \\ i, j \geq 0, r \geq 2}} T_4^i T_3^j \\ + 12p^3 q^3 \sum_{\substack{i+j=r-2 \\ i, j \geq 0, r \geq 2}} T_4^i T_2^j + 24p^6 q^3 \sum_{\substack{i+j+k=r-3 \\ i, j, k \geq 0, r \geq 3}} T_4^i T_3^j T_2^k$$

$$(d) \quad P(2, m) = mpq^{2m-1} + mq^{m-1} p^2 [(p+1)^m - 1].$$

OUTLINE OF PROOF: Notice that $T_s = p^s + q^s$ is the probability of a tie among s persons (either by all getting tails or all getting heads).

- (a) In this case, there is a tie among two players in each of the first $(r-1)$ rounds.
- (b) Among three players, game is decided in the r th round in one of the two ways:
 - (i) There is a three way tie in each of the first $(r-1)$ rounds.
 - (ii) One player is disqualified sometime in the first $(r-1)$ rounds and another player in the r th round.

In the representation of $P(r, 3)$, first term corresponds to case (i) and the summation to case (ii).

- (c) For four players, the game is decided in one of the following ways, depending upon when three players are out.

- (i) Three players lose the toss in the last round (and there is a 4-way tie in each of the first $(r-1)$ rounds).
- (ii) Two players lose the toss in r th round and one player earlier.
- (iii) One player loses in the last round and two players lose earlier either both in the same round or in two separate rounds.

$$(d) \quad \text{First notice that } P(1, m) = mpq^{m-1}.$$

Now

$$\begin{aligned} P(2, m) &= (p^m + q^m)P(1, m) + \sum_{i=1}^{m-2} \binom{m}{i} p^{m-i} q^i P(1, m-i) \\ &= mpq^{2m-1} + m \sum_{i=0}^{m-2} \binom{m-1}{i} p^{m-i+1} q^{m-1} \\ &= mpq^{2m-1} + mq^{m-1} p^2 [(p+1)^m - 1]. \end{aligned}$$

3. REMARK.

$P(2, m)/P(1, m) = q^m + p[(1+p)^m - 1]$ approaches $+\infty$ with m . Thus for large m , there is much greater chance for the game to be decided in two rounds than in one round.

4. THEOREM.

If the coin is fair ($p = 1/2$), then

$$P(r, m) = m2^{-rm} \sum_{k=1}^{2^r-1} (-1)^{k+1} k^{m-1} \quad (1)$$

PROOF. For $k = 1, \dots, m$ let $P(r, m, k)$ be the probability that starting with m persons at the end of r rounds there are k players in the game. Then $P(r, m)$ above is just $P(r, m, 1)$.

First note that for $m \geq 2$ and $k = 1, \dots, m$

$$P(1, m, k) = \left\{ \binom{m}{k} + \delta(k-m) \right\} 2^{-m},$$

where $\delta(x)$ is the indicator function of $\{0\}$. In particular, we have that

$$P(1, m, 1) = m2^{-m}.$$

We will use induction to show that (1) holds.

We will show below that if (1) holds for $r = n$ then it also holds for $r = n + 1$.

By conditioning on the outcome of the first toss we get that

$$\begin{aligned} P(n+1, m, 1) &= \sum_{j=2}^m P(1, m, j)P(n, j, 1) \\ &= \sum_{j=2}^m \binom{m}{j} 2^{-m} j 2^{-nj} \sum_{k=1}^{2^n-1} (-1)^{k+1} k^{j-1} + \\ &\quad + m 2^{-(n+1)m} \sum_{k=1}^{2^n-1} (-1)^{k+1} k^{m-1} \\ &= 2^{-m} \sum_{k=1}^{2^n-1} (-1)^{k+1} m 2^{-n} \sum_{l=1}^{m-1} \binom{m-1}{l} k^l 2^{-nl} + \\ &\quad + m 2^{-(n+1)m} \sum_{k=1}^{2^n-1} (-1)^{k+1} k^{m-1} \\ &= m 2^{-(n+1)m} \sum_{k=1}^{2^{n+1}} (-1)^{k+1} k^{m-1} \end{aligned}$$

after some simplification. This concludes the proof.

REFERENCES

1. BRUSS, F.T. & O'CINNEIDE, C.A., On the maximum and its uniqueness for geometric random samples, J. Appl. Prob. 27 (1990), 598-610.

Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Lead Guest Editor

Juan J. Nieto, Departamento de Análisis Matemático,
Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;
juanjose.nieto.roig@usc.es

Guest Editor

Donal O'Regan, Department of Mathematics, National
University of Ireland, Galway, Ireland;
donal.oregan@nuigalway.ie