

AN EXTENSION OF A BEST APPROXIMATION THEOREM

A. CARBONE
Dipartimento di Matematica
Università della Calabria
87036 Arcavacata di Rende (Cosenza), ITALY

(Received November 14, 1994 and in revised form February 10, 1995)

ABSTRACT. In this paper an extension of a theorem of Prolla is given and several interesting corollaries are derived. Fixed point results are also given in the end.

KEY WORDS AND PHRASES. Almost affine map, coincidence point, approximatively compact set, fixed points.

1991 AMS SUBJECT CLASSIFICATION CODES. Primary 47H10, Secondary 54X10.

1. INTRODUCTION.

Recently several researchers have given extensions of the following well-known theorem of Ky Fan [4] on best approximation.

THEOREM 1. Let C be a compact convex subset of a normed linear space X and $f : C \rightarrow X$ a continuous function. Then there is an $x_0 \in C$ such that

$$\|x_0 - fx_0\| = d(fx_0, C) = \inf\{\|fx_0 - y\| : y \in C\}.$$

There are several proofs of this theorem using the KKM-map principle, variational inequality and fixed point theory. The above theorem has interesting applications in fixed point theory and approximation theory.

Prolla [8] gave the following.

THEOREM 2. Let C be a nonempty compact convex subset of a normed linear space X , and $g : C \rightarrow C$ continuous, almost-affine, onto map. If $f : C \rightarrow X$ is a continuous function then there is a $y_0 \in C$ such that

$$\|gy_0 - fy_0\| = d(fy_0, C).$$

Note. In case $g = I$, an identity function, then Theorem 1 is obtained.

The purpose of this paper is to extend Theorem 2 and derive a few interesting corollaries.

We need the following.

Let X be a Banach space and C a nonempty subset of X . Let $x \in X$ and denote $d(x, C) = \inf\{\|x - y\| : y \in C\}$. In case $d(x, C) = \|x - y\|$ for $y \in C$, then y is said to be an element of best approximation to x . The set of best approximation to x is given by

$$P(x) = \{y \in C : \|x - y\| = d(x, C)\}.$$

The map $P : X \rightarrow 2^C$ is called the metric projection onto C . If $Px \neq \emptyset$ for all $x \in X$ then C is called a proximinal set. In case $P(x)$ contains at most one element for each $x \in X$ then C is called a Chebyshev set.

A subset C of X is called an approximatively compact if for every $x \in X$ and every sequence $\{y_n\}$ in C with $\lim_{n \rightarrow \infty} \|x - y_n\| = d(x, C)$ there exists a subsequence $\{y_{n_i}\}$ converging to an element of C .

A compact set is always approximatively compact but converse is not true. For example, a closed convex set in a Hilbert space is approximatively compact but not compact.

For an approximately compact set C the following holds.

- i) $Px \neq \emptyset$ for each $x \in X$;
- ii) C is closed;
- iii) Px is compact;
- iv) if C is convex then Px is convex;
- v) the metric projection $P : X \rightarrow 2^C$ is upper semicontinuous (see [9] or [11]);
- vi) $P(A) = \cup \{P(x) : x \in A\}$ is compact for any compact subset A in X .

Let X and Y be normed linear spaces and 2^Y denote the set of all nonempty subsets of Y . A multivalued mapping $F : X \rightarrow 2^Y$ is upper semicontinuous (usc) if $F^{-1}(A) = \{x \in X : Fx \cap A \neq \emptyset\}$ is closed in X for each closed set A in Y .

A multivalued map F is said to be compact if $F(X)$ is contained in a compact subset of Y . F is said to be acyclic if Fx is nonempty, compact and acyclic subset of Y for each $x \in X$.

A multivalued map $F : X \rightarrow X$ (X is a metric space) is said to be admissible if there are maps

$$F_i : X_i \rightarrow X_{i+1} \quad i = 0, 1, 2, \dots, n \quad X_0 = X_{n+1} = X$$

such that:

- i) $F = F_n F_{n-1} \dots F_0$;
- ii) F_i is acyclic and usc for each i ;
- iii) X_i are metric spaces for each $i = 1, 2, \dots, n$ (see [7]).

The following theorem will be used [7] in our work.

THEOREM 3. Let C be a convex subset of a Banach space X and $F : C \rightarrow C$ an admissible compact map. Then F has a fixed point.

Let C be a convex subset of X . A map $g : C \rightarrow X$ is almost-affine if it satisfies

$$\|g(\lambda x_1 + (1 - \lambda)x_2) - y\| \leq \lambda \|gx_1 - y\| + (1 - \lambda) \|gx_2 - y\|$$

for all $x_1, x_2 \in C$, $y \in X$ and $0 < \lambda < 1$.

g is an affine map if

$$g(\lambda x_1 + (1 - \lambda)x_2) = \lambda gx_1 + (1 - \lambda)gx_2, \quad \lambda \in (0, 1).$$

If $g : C \rightarrow C$ is a single-valued function then g is said to be proper if $g^{-1}(A)$ is compact for A compact.

The following is the main result.

THEOREM 4. Let C be a nonempty convex subset of a normed linear space X and $P : X \rightarrow 2^C$ the metric projection satisfying

- i) $P(x) = \{y \in C : \|x - y\| \leq \|x - z\| \text{ for all } z \in C\} \neq \emptyset \text{ for each } x \in X$
and
- ii) P sends compact subsets of X onto compact subsets of C .

Let $g : C \rightarrow C$ be a continuous, onto, proper and $g^{-1}(z)$ an acyclic subset of C for every $z \in C$.

Then for every continuous map $f : C \rightarrow X$ with $f(C)$ relatively compact there exists a $y_0 \in C$ such that

$$\|gy_0 - fy_0\| = d(fy_0, C).$$

Note. 1. In case C is an approximatively compact set then conditions i) and ii) are satisfied by P .

2. In case C is a compact convex set then the condition that $f(C)$ is relatively compact is not required since the continuous image of a compact set is compact. The condition that g is proper is also not needed, since g is continuous so for any compact set D in C , $g^{-1}(D)$ is a closed subset of a compact set C and hence is compact.

The proof is on the same lines as in [3].

PROOF. Let $P : X \rightarrow 2^C$ be the metric projection. Define a multivalued map $F : C \rightarrow C$ by

$$Fx = \{y \in C : gy \in P(f(x))\}.$$

Then $Fx \neq \emptyset$ for each $x \in C$ since g is an onto map. $F(x)$ is closed and acyclic (see [3]). We show that F is upper semicontinuous. This will be done as in [10].

Let B be a closed subset of C and z a limit point of $F^{-1}(B)$. We choose $\{z_n\} \subseteq F^{-1}(B) \subseteq C$ such that $z_n \rightarrow z$. We show that $z \in F^{-1}(B)$. Since for each n , $F(z_n) \cap B \neq \emptyset$ we have $\{y_n\} \subseteq C$ with $y_n \subseteq F(z_n) \cap B$.

Then for each n ,

$$\|gy_n - fz_n\| = d(fz_n, C). \quad (*)$$

Let $A = \overline{f(C)}$ and $D = P(A)$. By ii) D is a compact subset of C . Since g is a proper map $g^{-1}(D)$ is a compact subset of C .

Now, for each n ,

$$y_n \in F(z_n) = g^{-1}Pf(z_n) \subseteq g^{-1}Pf(C) \subseteq g^{-1}(D).$$

Consequently, there exists $y \in C$ and a subsequence $\{y_{n_i}\}$ of $\{y_n\}$ such that $y_{n_i} \rightarrow y$. By $(*)$ we get that

$$\|gy - fz\| = d(fz, C).$$

This gives $y \in F(z) \cap B$, that is, $z \in F^{-1}(B)$ and F is an upper semicontinuous multifunction.

Now, the map $F = g^{-1} \circ P \circ f : C \rightarrow C$ is an admissible map. Since $f(C)$ is relatively compact therefore $F(C) = g^{-1}(P(f(C)))$ is also relatively compact because the image of a compact set under upper semicontinuous map with compact values is compact.

Then F satisfies conditions of Theorem 3 and has a fixed point, say $y_0 \in F(y_0)$. This implies that

$$\|gy_0 - fy_0\| = d(fy_0, C).$$

An almost affine map satisfies the condition that $g^{-1}(z)$ is an acyclic set for each $z \in C$. As a consequence we have the following [10].

COROLLARY 1. Let C be a nonempty convex subset of a normed linear space X and $P : X \rightarrow 2^C$ the metric projection satisfying i) and ii) of Theorem 4. Let $g : C \rightarrow C$ be a continuous, onto, almost affine and proper map. Then for each continuous map $f : C \rightarrow X$ with $f(C)$ relatively compact there exists a $y_0 \in C$ such that

$$\|gy_0 - fy_0\| = d(fy_0, C).$$

In case C is an approximatively compact set then conditions i) and ii) are satisfied and we have the following [3].

COROLLARY 2. Let C be an approximatively compact convex set of a normed linear space X and $g : C \rightarrow C$ continuous, onto, proper and $g^{-1}(z)$ is an acyclic subset of C for every $z \in C$. Then for each continuous function $f : C \rightarrow X$ with $f(C)$ relatively compact there exists an $x_0 \in C$ such that

$$\|gx_0 - fx_0\| = d(fx_0, C).$$

In case C is an approximatively compact and $g = I$, an identity function, then we get a well-known result of Reich [9].

Recently in [2] almost quasi convex g was considered.

The function $g : C \rightarrow X$ is said to be almost quasi convex if

$$\|g(\lambda x_1 + (1 - \lambda)x_2) - y\| \leq \max(\|gx_1 - y\|, \|gx_2 - y\|)$$

for $x_1, x_2 \in C$, $y \in X$ and $0 < \lambda < 1$.

As in [3] it is easy to see that almost quasi convex implies that $g^{-1}(z)$ is an acyclic set. Therefore a recent result due to Park, Singh and Watson [6] given below, follows as a corollary.

COROLLARY 3. Let C be a nonempty convex subset of a normed linear space X and $P : X \rightarrow 2^C$ the metric projection satisfying

- i) $Px \neq \emptyset$ for each $x \in X$, and
- ii) P sends compact subset of X onto compact subsets of C .

Let $g : C \rightarrow C$ be a continuous, almost quasi convex, onto and proper map.

Then for each continuous map $f : C \rightarrow X$ with $\overline{f(C)}$ compact there is a $y_0 \in C$ such that

$$\|gy_0 - fy_0\| = d(fy_0, C).$$

Note. They [6] concluded that either

- i) f and g have a coincidence point $w \in C$, i.e., $fw = gw$, or
- ii) there is a $w \in C$ such that $gw \in \partial C$ and

$$0 < \|gw - fw\| \leq \|z - fw\| \quad \text{for all } z \in I_C(gw).$$

Recall that if $f : C \rightarrow X$ is a map, then the inward set $I_C(x)$, of C at x is defined by

$$I_C(x) = \{y : y \in X, \text{ there exist } u \in C \text{ and } r > 0 \text{ such that } y = x + r(u - x)\}.$$

The closure is denoted by $\overline{I_C(x)}$. The function f is said to be inward map if $f(x) \in I_C(x)$ for every $x \in C$ and weakly inward if $f(x) \in \overline{I_C(x)}$.

In case C is a compact convex subset of a normed linear space X we get the following.

COROLLARY 4. Let C be a compact convex subset of a normed linear space X and $g : C \rightarrow C$ continuous, onto, $g^{-1}(z)$ is an acyclic set for each $z \in C$. Then for each continuous function $f : C \rightarrow X$ there is a $y_0 \in C$ such that

$$\|gy_0 - fy_0\| = d(fy_0, C).$$

If $g = I$, an identity function, in Corollary 4, then we get a well-known result of Ky Fan [4].

If in Corollary 4, g is almost-affine, continuous, onto then we get a theorem of Prolla [8].

If $fy_0 \in C$ for all y_0 then Corollary 4 yields a coincidence theorem, that is, there is a $y_0 \in C$ such that

$$gy_0 = fy_0.$$

In case g is an identity function in Corollary 4 and f has an additional condition then we get a fixed point theorem given below.

COROLLARY 5. If all the hypotheses of Corollary 4 are satisfied with $g = I$, and in addition: for any $x \in \partial C$ with $x \neq fx$ there exists a number λ (real or complex depending on whether the vector space is real or complex) such that $|\lambda| < 1$ and

$$y = \lambda x + (1 - \lambda)fx \in C,$$

then f has a fixed point.

PROOF. By Corollary 4 we have that

$$\|y_0 - fy_0\| = d(fy_0, C).$$

Take $y_0 \in \partial C$ and assume that $y_0 \neq fy_0$. Then

$$\begin{aligned} 0 < \|y_0 - fy_0\| &\leq \|y - fy_0\| = \|\lambda y_0 + (1 - \lambda)fy_0 - fy_0\| \\ &= \|\lambda(y_0 - fy_0)\| \\ &= |\lambda| \|y_0 - fy_0\| < \|y_0 - fy_0\| \end{aligned}$$

a contradiction. Hence $y_0 = fy_0$.

In case $g = I$, an identity function, then we could derive a very well-known result due to Browder [1] as a corollary.

COROLLARY 6. Let X be a Banach space, C a compact, convex subset of X and $f : C \rightarrow X$ a continuous function. Suppose that for each $x \in C$ with $x \neq fx$ there exists a $y \in I_C(x)$ such that

$$\|y - fx\| < \|x - fx\|.$$

Then f has a fixed point.

PROOF. By Corollary 4 (with $g = I$), we have

$$\|x - fx\| = d(fx, C)$$

i.e. $\|x - fx\| \leq \|fx - z\|$ for all $z \in C$.

This inequality remains valid (see [5]) for all $y \in I_C(x)$, i.e.

$$\|x - fx\| \leq \|fx - y\| \quad \text{for all } y \in I_C(x). \quad (**)$$

By hypothesis for each $x \in C$ with $x \neq fx$ we have $y \in I_C(x)$ such that

$$\|y - fx\| < \|x - fx\|$$

a contradiction to (**). Hence f has a fixed point.

Further work in this direction due to Park [5] gives several applications.

REFERENCES

- [1] BROWDER, F.E., On a sharpened form of the Schauder fixed point theorem, *Proc. Natl. Acad. Sci. USA* **74** (1977), 4749-4751.
- [2] CARBONE, A., A note on a theorem of Prolla, *Indian J. Pure Appl. Math.* **23** (1991), 257-260.
- [3] CARBONE, A. and CONTI, G., Multivalued maps and existence of best approximants, *J. Approx. Theory* **64** (1991), 203-208.
- [4] FAN, K., Extensions of two fixed point theorems of F. E. Browder, *Math. Z.* **112** (1969), 234-240.
- [5] PARK, S., On generalization of Ky Fan's theorem on best approximations, *Numer. Funct. Anal. Optimiz.* **9** (1987), 619-628.
- [6] PARK, S., SINGH, S.P. and WATSON, B., Remarks on best approximations and fixed points, *J. Pure Appl. Math.* **25** (5) (1994), 459-462.
- [7] POWERS, M.J., Lefschetz fixed point theorems for a new class of multivalued maps, *Pacif. J. Math.* **42** (1972), 211-220.
- [8] PROLLA, J.B., Fixed point theorems for set-valued mappings and existence of best approximants, *Numer. Funct. Anal. Optimiz.* **5** (1982-83), 449-455.

- [9] REICH, S., Approximate selections, best approximations, fixed points, and invariant sets, *J. Math. Anal. Appl.* **62** (1978), 104-113
- [10] SEHGAL, V.M. and SINGH, S.P., A theorem on best approximations, *Numer. Funct. Anal. Optimiz.* **10** (1989), 181-184
- [11] SINGER, I., *Best Approximations in Normed Linear Spaces by Elements of Linear Subspaces*, Springer-Verlag (1970), 368

Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Lead Guest Editor

Juan J. Nieto, Departamento de Análisis Matemático,
Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;
juanjose.nieto.roig@usc.es

Guest Editor

Donal O'Regan, Department of Mathematics, National
University of Ireland, Galway, Ireland;
donal.oregan@nuigalway.ie