

A SIMPLIFICATION OF D'ALARCAO'S IDEMPOTENT SEPARATING EXTENSIONS OF INVERSE SEMIGROUPS

CONSTANCE C. EDWARDS

Department of Mathematical Sciences
Indiana University-Purdue University at Fort Wayne
Fort Wayne, Indiana 46805 U.S.A.

(Received August 29, 1977)

ABSTRACT. In [2] D'Alarcao states necessary and sufficient conditions for the attainment of an idempotent-separating extension of an inverse semigroup. To do this D'Alarcao needed essentially three mappings satisfying thirteen conditions. In this paper we show that one can achieve the same results with two mappings satisfying eight conditions.

KEY WORDS AND PHRASES. Inverse semigroup and idempotent-separating extension.

AMS(MOS) SUBJECT CLASSIFICATION (1970) CODES. 20M10

I. THE CONSTRUCTION.

Unless otherwise mentioned, the terminology and notation of Clifford and Preston [1] shall be assumed. The reader is referred to [2] for all other undefined terms.

LEMMA 1. If A and B are inverse semigroups and f is a homomorphism of A onto $E(B)$ which is one-to-one on $E(A)$, then

- (i) $af = (a^{-1}a)f$ for all $a \in A$;
- (ii) $a^{-1}ga = ga^{-1}a$ for all $a \in A$ and $g \in E(A)$; and
- (iii) $ga = ag$ for all $a \in A$ and $g \in E(A)$.

PROOF. (i) Since $af \in E(B)$, by Lallement's result [3], there exists $e \in E(A)$ such that $af = ef$. Consequently, $ef = e^{-1}f = (ef)^{-1} = (af)^{-1} = a^{-1}f$, and thus $af = ef = (a^{-1}f)(af) = (a^{-1}a)f$.

(ii) By part (i), for each $g \in E(A)$ and $a \in A$,

$$((ga)^{-1}ga)f = (ga)f = gf \cdot af = gf \cdot (a^{-1}a)f = (ga^{-1}a)f.$$

Since f is one-to-one on $E(A)$, we have $a^{-1}ga = (ga)^{-1}ga = ga^{-1}a$.

(iii) By part (ii), $ga = a(a^{-1}ga) = a(ga^{-1}a) = ag$. This completes the proof of the lemma.

Let A and B be inverse semigroups and let f be a homomorphism of A onto $E(B)$ which is one-to-one on $E(A)$. (The existence of such an f implies that A is a semilattice of groups; see e.g. [4].) Suppose there is a mapping w of $B \times B$ into A (denoted by $(b,c)w = b^c$) and that for each $b \in B$ there is a mapping U_b of A into A (denoted by $aU_b = a^b$) such that:

- (P1) For each $b, c \in B$, $b^c \in (c^{-1}b^{-1}bc)f^{-1}$.
- (P2) For each $b \in B$, U_b is a homomorphism.
- (P3) If $b, c \in B$ and $a \in (b^{-1}b)f^{-1}$, then $a^c \in (c^{-1}b^{-1}bc)f^{-1}$.
- (P4) $(bc)^d(b^c)^d = b^{cd}c^d$ for all $b, c, d \in B$.
- (P5) If $b \in E(B)$ and $a \in A$, then $a^b = ea$ where $\{e\} = E(A) \cap bf^{-1}$.
- (P6) For each $b \in B$, $(bb^{-1})^b = e$ where $\{e\} = E(A) \cap (b^{-1}b)f^{-1}$.
- (P7) $b^c(a^b)^c = a^{bc}b^c$ for all $a \in A$ and $b, c \in B$.
- (P8) For each $b, c \in E(B)$, $b^c = eg$ where $\{e\} = E(A) \cap bf^{-1}$ and $\{g\} = E(A) \cap cf^{-1}$.

Let $S^* = \{(b,a) : b \in B \text{ and } a \in (b^{-1}b)f\}$. Let equality in S^* be defined in the usual manner and let multiplication be defined by $(b,a)(c,p) = (bc, b^ca^p)$.

Moreover, let $\gamma: S^* \rightarrow B$ be defined by $(b,a)\gamma = b$.

THEOREM 1. Under the above conditions, (S^*, γ) is an idempotent-separating extension of A by B . Moreover, $\gamma|_A = f$.

PROOF. Note that since fof^{-1} is idempotent separating, $fof^{-1} \subseteq \mathbb{N}$. Consequently, if $e \in E(A)$ and $e, a \in (b^{-1}b)f^{-1}$, then $ea = a = ae$.

As is shown in [2], closure in S^* follows from (P1) and (P3) and, associativity follows from (P2), (P4) and (P7).

Our proof that S^* is regular is the same as that found in [2] but is more informative. Let $(b,a) \in S^*$ and let $x = (a^{b^{-1}})^{-1}(b^{b^{-1}})^{-1}$. By (P1) and (P3), $((a^{b^{-1}})^{-1})f = bb^{-1} = ((b^{b^{-1}})^{-1})f$. Consequently $(b^{-1}, x) \in S^*$ since $xf = ((a^{b^{-1}})^{-1}(b^{b^{-1}})^{-1})f = bb^{-1}$. Also, by (P1) and (P3), $(b^{b^{-1}}a^{b^{-1}}x)^b f = b^{-1}b$. Since $(b^{b^{-1}}a^{b^{-1}}x)^b \in E(A) \cap (b^{-1}b)f^{-1}$ we have $(b^{b^{-1}}a^{b^{-1}}x)^b a = a$. It now follows from (P6) that $(b,a)(b^{-1},x)(b,a) = (b, (bb^{-1})^b (b^{b^{-1}}a^{b^{-1}}x)^b a) = (b, ea) = (b,a)$ since $\{e\} = E(A) \cap (b^{-1}b)f^{-1}$. Consequently, S^* is regular.

As shown in [2], it follows from (P5) and (P6) that if $(g,e) \in E(S^*)$ then $g \in E(B)$ and $\{e\} = E(A) \cap gf^{-1}$. Consider the subset $A^* = \{(b,a) : b \in E(B)\} \subseteq S^*$ which contains $E(S^*)$. Let $\delta: A^* \rightarrow A$ by $(b,a) \rightarrow a$. To show δ is a homomorphism, let $(b,a), (c,p) \in A^*$. Then $a = ea$ and $p = gp$ where $\{e\} = E(A) \cap bf^{-1}$ and $\{g\} = E(A) \cap cf^{-1}$. By (P8), (P5) and lemma lili, $(b,a)(c,p) = (bc, b^ca^cp) = (bc, eg \cdot ga \cdot p) = (bc, ea \cdot gp) = (bc, ap)$. Thus δ is a homomorphism which obviously has A for its range. It follows from the definitions of S^* and δ that δ is one-to-one. Consequently the idempotents of S^* commute since A^* is isomorphic to the inverse semigroup A . We now have that S^* is an inverse semigroup.

It is obvious that γ is a homomorphism of S^* onto B and $\{E(B)\}\gamma^{-1} = A^*$.

If $(b,a), (c,p) \in E(A^*)$ such that $(b,a)\gamma = (c,p)\gamma$, then $b = c$ and $af = b = c = pf$. Since $a, p \in E(A)$, $a = p$. Consequently γ is one-to-one on $E(A^*)$. It is obvious that $\gamma|_{A^*} = f$. Since $A^* = A$, this completes the proof of the theorem.

2. THE STRUCTURE THEOREM.

Since our two mappings and eight properties are the same as, but fewer than, those used by D'Alarcao, the following theorem has been proven by D'Alarcao ([2], Theorem 2).

THEOREM 2. Let A and B be inverse semigroups and let (S,f) be an idempotent-separating extension of A by B . Then for each $b \in B$ there exists a mapping U_b of A into A (denoted by $aU_b = a^b$) and there exists a mapping w of $B \times B$ into A satisfying (P1) through (P8). Moreover, S is isomorphic to S^* .

REFERENCES

1. Clifford, A. H. and G. B. Preston The Algebraic Theory of Semigroups, Vol. I, Math Surveys of the American Math. Soc. 7 (Providence, RI, 1961).
2. D'Alarcao, H. Idempotent-Separating Extensions of Inverse Semigroups, J. Aust. Math. Soc. 9 (1969) 211-17.
3. Lallement, G. Congruences et Équivalences de Green sur un Demi-groupe Régulier, C. R. Acad. Sc. Paris Ser A 262 (1966) 613-6.
4. Preston, G. B. Inverse Semi-groups, J. of the London Math. Soc. 29 (1954) 396-403.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br