

DUALITY IN THE OPTIMAL CONTROL FOR DAMPED HYPERBOLIC SYSTEMS WITH POSITIVE CONTROL

MI JIN LEE, JONG YEOUL PARK, and YOUNG CHEL KWON

Received 10 September 2002

We study the duality theory for damped hyperbolic equations. These systems have positive controls and convex cost functionals. Our main results lie in the application of duality theorem, that is, $\inf J = \sup K$, on various cost functions.

2000 Mathematics Subject Classification: 49J20, 49N15, 93C20.

1. Introduction. Lions [2] introduced optimal control problems of the variety of distributed parameter systems, for example, elliptic, parabolic, and hyperbolic. Here, we see that adjoint state systems are given cost functional and distributed parameter systems. Duality theory is $\inf J = \sup K$, where J satisfies systems and K satisfies adjoint state systems. The duality theory for the corresponding parabolic systems has been given by Chan [1] and Tanimoto [7]. Park and Lee [5, 6] studied the duality theory for hyperbolic systems. Also, they [4] obtained same results for hyperbolic systems with damping terms. In this paper, we study the duality theory for damped hyperbolic systems with positive controls. These systems have various convex cost functionals. The main objective is to prove the duality theorem for damped hyperbolic systems with positive controls and various cost functions. The main tools are integration by parts and Green's formula.

2. Preliminaries. Let X be a Hilbert space $(\cdot, \cdot)$ and let $\|\cdot\|_X$ denote the inner product and the induced norm on X ; X' denotes the dual space of X and $\langle \cdot, \cdot \rangle_{X', X}$ denotes the dual pairing between X' and X . We introduce underlying Hilbert spaces to describe damped second-order evolution equations. Let H be a real pivot Hilbert space; its norm $\|\cdot\|_H$ is simply denoted by $|\cdot|_H$. For $i = 1, 2$, let V_i be a real separable Hilbert space. Assume that each pair (V_i, H) is a Gelfand triple space with a notation $V_i \hookrightarrow H \equiv H' \hookrightarrow V'_i$. We suppose that V_1 is continuously embedded in V_2 . Then we see that $V_1 \hookrightarrow V_2 \hookrightarrow H \equiv H' \hookrightarrow V'_2 \hookrightarrow V'_1$ and the equalities $\langle \phi, \psi \rangle_{V'_1, V_1} = \langle \phi, \psi \rangle_{V'_2, V_2}$ for $\phi \in V'_2$, $\psi \in V$, and $\langle \phi, \psi \rangle_{V'_1, V_1} = (\phi, \psi)_H$ for $\phi \in H$, $\psi \in V$ hold. Let T be a positive number. We define a function space $W(0, T)$ by

$$W(0, T) = \{y \mid y \in L^2(0, T; V_1), y' \in L^2(0, T; V_2), y'' \in L^2(0, T; V'_1)\} \quad (2.1)$$

with an inner product

$$(\mathcal{Y}_1, \mathcal{Y}_2)_{W(0,T)} = \int_0^T \{ (\mathcal{Y}_1(t), \mathcal{Y}_2(t))_{V_1} + (\mathcal{Y}'_1(t), \mathcal{Y}'_2(t))_{V_2} + (\mathcal{Y}''_1(t), \mathcal{Y}''_2(t))_{V'_1} \} dt. \quad (2.2)$$

This becomes a Hilbert space with norm

$$\|\mathcal{Y}\|_{W(0,T)} = \left(\|\mathcal{Y}\|_{L^2(0,T;V_1)}^2 + \|\mathcal{Y}'\|_{L^2(0,T;V_2)}^2 + \|\mathcal{Y}''\|_{L^2(0,T;V'_1)}^2 \right)^{1/2}, \quad (2.3)$$

where $' = d/dt$ and $'' = d^2t/dt^2$.

From now on, we set $V = V_2 = H_0^1(\Omega)$ and $H = L^2(\Omega)$, where Ω is a bounded open set in $\mathbb{R}^n$ with smooth boundary Γ , and let $Q = \Omega \times (0, T)$. We will give an exact description of damped second-order evolution equations. We consider the bilinear forms defined by

$$\begin{aligned} a_1(t; \phi, \psi) &= \sum_{i,j=1}^n \int_{\Omega} a_{ij}(t, x) \frac{\partial \phi(x)}{\partial x_i} \frac{\partial \psi(x)}{\partial x_j} dx \\ &\quad + \int_{\Omega} a_0(t, x) \phi(x) \psi(x) dx \quad \forall \phi, \psi \in V, \\ a_2(t; \phi, \psi) &= \sum_{i,j=1}^n \int_{\Omega} b_{ij}(t, x) \frac{\partial \phi(x)}{\partial x_i} \frac{\partial \psi(x)}{\partial x_j} dx \\ &\quad + \int_{\Omega} b_0(t, x) \phi(x) \psi(x) dx \quad \forall \phi, \psi \in V, \end{aligned} \quad (2.4)$$

where a_{ij} , b_{ij} , a_0 , and b_0 are the functions satisfying the following properties:

- (i) $a_{ij} = a_{ji}$ and $b_{ij} = b_{ji}$,
- (ii) $a_{ij}, b_{ij}, a_0, b_0 \in C^1([0, T]; L^\infty(\Omega))$,
- (iii) $\sum_{i,j=1}^n a_{ij}(t, x) \xi_i \xi_j \geq c_1 (\xi_1^2 + \cdots + \xi_n^2)$, $c_1 > 0$, $\xi_i \in \mathbb{R}$,
- (iv) $\sum_{i,j=1}^n b_{ij}(t, x) \xi_i \xi_j \geq c_2 (\xi_1^2 + \cdots + \xi_n^2)$, $c_2 > 0$, $\xi_i \in \mathbb{R}$.

Using the above properties, we can show the coercivity condition of a_1 and a_2 . Indeed, by (i) and (ii), there exists $K > 0$ such that $|a_0(t, x)| \leq K$ a.e., $x \in \Omega$, and for all $t \in [0, T]$. The coercivity condition of a_1 follows from

$$\begin{aligned} a_1(t; \phi, \psi) &\geq c_1 \sum_{i=1}^n \int_{\Omega} \left| \frac{\partial \phi(x)}{\partial x_i} \right|^2 dx - K \int_{\Omega} |\phi(x)|^2 dx \\ &\geq c_1 \|\phi\|_{H_0^1(\Omega)}^2 - K \|\phi\|_{L^2(\Omega)}^2. \end{aligned} \quad (2.5)$$

Similarly, we can show the coercivity condition of a_2 . Then we can define the operator $A_i(t) \in \mathcal{L}(V_i, V'_i)$ for $t \in [0, T]$ deduced by the relation

$$a_i(t; \phi, \psi) = \langle A_i(t) \phi, \psi \rangle_{V'_i, V_i} \quad \forall \phi, \psi \in V_i, \quad i = 1, 2. \quad (2.6)$$

Let $y_0 \in H_0^1(\Omega)$ and $y_1 \in L^2(\Omega)$. Then by Nakagiri and Ha [3], there exists a solution $y \in W(0, T)$ of

$$\begin{aligned} \frac{\partial^2 y(u; t)}{\partial t^2} + A_2(t) \frac{\partial y(u; t)}{\partial t} + A_1(t) y(u; t) &= u \quad \text{in } (0, T), \\ y(u; 0) &= y_0, \quad y'(u; 0) = y_1 \quad \text{in } \Omega, \quad y = 0 \quad \text{on } \Sigma. \end{aligned} \quad (2.7)$$

3. Duality. We choose a control variable space $\mathcal{U} = L^2(Q)$. Let $u \in \mathcal{U}_{ad} = \{u \mid u \geq 0 \text{ a.e. in } Q\} \subset \mathcal{U}$ and y satisfying (2.7). we consider a cost functional given by

$$J(y, u) = \frac{1}{2} \int_0^T |y(u) - z_d|^2 dt + \frac{1}{2} \int_0^T (Nu, u) dt. \quad (3.1)$$

For our purpose, we consider the following systems:

$$\begin{aligned} y''(u; t) + A_2(t) y'(u; t) + A_1(t) y(u; t) &= u \quad \text{in } Q, \\ y(u; 0) &= y_0, \quad y'(u; 0) = y_1 \quad \text{in } \Omega, \\ u &\geq 0 \quad \text{a.e. in } Q, \quad y(0) \geq 0, \quad y'(0) \geq 0, \quad y = 0 \quad \text{on } \Sigma, \end{aligned} \quad (3.2)$$

$$\begin{aligned} p''(u; t) - A_2(t) p'(u; t) + (A_1(t) - A_2'(t)) p(u; t) &= y - z_d \quad \text{in } Q, \\ p &= 0 \quad \text{on } \Sigma, \quad p(u; T) = 0, \quad p'(u; T) = 0 \\ p + Nu &\geq 0 \quad \text{in } Q, \quad p(0) \geq 0, \quad p'(0) \geq 0, \end{aligned} \quad (3.3)$$

$$\begin{aligned} u(p + Nu) &= 0, \\ p(0) y(0) &= 0 \quad \text{in } \Omega, \quad y'(0) p(0) = 0, \quad y(0) p'(0) = 0. \end{aligned} \quad (3.4)$$

THEOREM 3.1. Let $J = (1/2) \int_0^T |y - z_d|^2 dt + (1/2) \int_0^T (Nu, u) dt$ and $K = -(1/2) \int_0^T |y|^2 dt + (1/2) \int_0^T |z_d|^2 dt - (1/2) \int_0^T (Nu, u) dt$. Assume that y_0, u_0 , and p_0 satisfy (3.2), (3.3), and (3.4), respectively, y and u in J satisfy (3.2), and y and u in K satisfy (3.3). Then

$$\inf_{(3.2)} J = J(y_0, u_0) = K(y_0, u_0) = \sup_{(3.3)} K. \quad (3.5)$$

PROOF. (i) We begin by showing that $J = K$ at (y_0, u_0, p_0) .

$$\begin{aligned} J(y_0, u_0) &= J(y_0, u_0) - \int_0^T (u_0, p_0) dt - \int_0^T (u_0, Nu_0) dt \\ &= J(y_0, u_0) - \int_0^T (y_0'' + A_2(t) y_0' + A_1(t) y_0, p_0) dt - \int_0^T (u_0, Nu_0) dt \end{aligned}$$

$$\begin{aligned}
&= J(y_0, u_0) - \int_0^T (y_0, p_0'' - A_2'(t)p_0 - A_2(t)p_0' + A_1(t)p_0) dt \\
&\quad - \int_0^T (u_0, Nu_0) dt \\
&= J(y_0, u_0) - \int_0^T (y_0, y_0 - z_d) dt - \int_0^T (u_0, Nu_0) dt.
\end{aligned} \tag{3.6}$$

(ii) To show $\inf_{(3.2)} J = J(y_0, u_0)$, we must check that $J(y, u) \geq J(y_0, u_0)$ under (3.2) for (y, u, p) and under (3.2), (3.3), and (3.4) for (y_0, u_0, p_0) . Now, we have

$$\begin{aligned}
J(y, u) - J(y_0, u_0) &\geq \int_0^T (y_0 - z_d, y - y_0) dt + \int_0^T (Nu_0, u - u_0) dt \\
&= \int_0^T (p_0'' - A_2(t)p_0' + (A_1(t) - A_2'(t))p_0, y - y_0) dt \\
&\quad + \int_0^T (Nu_0, u - u_0) dt \\
&= \int_0^T (p_0, y'' + A_2(t)y' + A_1(t)y) dt \\
&\quad - \int_0^T (p_0, y_0'' + A_2(t)y_0' + A_1(t)y_0) dt \\
&\quad + \int_0^T (Nu_0, u - u_0) dt \\
&= \int_0^T (p_0, u - u_0) dt + \int_0^T (Nu_0, u - u_0) dt \\
&\geq 0.
\end{aligned} \tag{3.7}$$

(iii) We claim that $K(y, u) \leq K(y_0, u_0)$ under (3.3) for (y, u, p) :

$$\begin{aligned}
&J(y_0, u_0) - J(y, u) \\
&\geq \int_0^T (y - z_d, y - y_0) dt + \int_0^T (Nu, u - u_0) dt \\
&= \int_0^T (p'' - A_2(t)p' + (A_1(t) - A_2'(t))p, y_0 - y) dt + \int_0^T (Nu, u_0 - u) dt \\
&\quad + \int_0^T (y_0'' + A_2(t)y_0' + A_1(t)y_0 - u_0, p_0 - p) dt \\
&= - \int_0^T (y - z_d, y) dt - \int_0^T (Nu, u) dt + \int_0^T (y_0, y_0 - z_d) dt \\
&\quad + \int_0^T (Nu_0, u_0) dt + \int_0^T (Nu + p, u_0) dt - \int_0^T (u_0, p_0 + Nu_0) dt
\end{aligned}$$

$$\begin{aligned}
&\geq - \int_0^T (\gamma - z_d, \gamma) dt - \int_0^T (Nu, u) dt \\
&\quad + \int_0^T (\gamma_0, \gamma_0 - z_d) dt + \int_0^T (Nu_0, u_0) dt.
\end{aligned} \tag{3.8}$$

Therefore

$$K(\gamma_0, u_0) \geq K(\gamma, u). \tag{3.9}$$

This completes the proof. $\square$

Now, we observe the terminal value of $\gamma(u; t)$. Since the observation $z(u)$ is given by $\gamma(u; T)$, the cost function is given as

$$J(\gamma, u) = \frac{1}{2} |\gamma(T) - z_d|^2 + \frac{1}{2} \int_0^T (Nu, u) dt. \tag{3.10}$$

We introduce the following systems:

$$\begin{aligned}
&\gamma''(u; t) + A_2(t)\gamma'(u; t) + A_1(t)\gamma(u; t) = u \quad \text{in } Q, \\
&\gamma(u; 0) = \gamma_0, \quad \gamma'(u; 0) = \gamma_1 \quad \text{in } \Omega, \\
&u \geq 0 \quad \text{a.e. in } Q, \quad \gamma(0) \geq 0, \quad \gamma'(0) \geq 0, \quad \gamma = 0 \quad \text{on } \Sigma,
\end{aligned} \tag{3.11}$$

$$\begin{aligned}
&p''(u; t) - A_2(t)p'(u; t) + (A_1(t) - A'_2(t))p(u; t) = 0 \quad \text{in } Q, \\
&p = 0 \quad \text{on } \Sigma, \\
&p(u; T) = 0, \quad p'(u; T) = \gamma(T) - z_d, \\
&-p + Nu \geq 0 \quad \text{in } Q, \quad p(0) \geq 0, \quad p'(0) \geq 0,
\end{aligned} \tag{3.12}$$

$$\begin{aligned}
&u(-p + Nu) = 0, \\
&p(0)\gamma(0) = 0 \quad \text{in } \Omega, \quad \gamma'(0)p(0) = 0, \quad \gamma(0)p'(0) = 0.
\end{aligned} \tag{3.13}$$

THEOREM 3.2. Let $J = (1/2)|\gamma(T) - z_d|^2 + (1/2) \int_0^T (Nu, u) dt$ and $K = -(1/2)|\gamma(T)|^2 + (1/2)|z_d|^2 - (1/2) \int_0^T (Nu, u) dt$. Assume that γ_0 , u_0 , and p_0 satisfy (3.11), (3.12), and (3.13), respectively, γ and u in J satisfy (3.11), and γ and u in K satisfy (3.12). Then

$$\inf_{(3.11)} J = J(\gamma_0, u_0) = K(\gamma_0, u_0) = \sup_{(3.12)} K. \tag{3.14}$$

PROOF. (i) We now prove that $J(\gamma_0, u_0) = K(\gamma_0, u_0)$:

$$\begin{aligned}
 J(\gamma_0, u_0) &= J(\gamma_0, u_0) + \int_0^T (p_0, u_0) dt - \int_0^T (Nu_0, u_0) dt \\
 &= J(\gamma_0, u_0) + \int_0^T (\gamma_0'' + A_2(t)\gamma_0' + A_1(t)\gamma_0, p_0) dt - \int_0^T (Nu_0, u_0) dt \\
 &= \frac{1}{2} |\gamma_0(T) - z_d|^2 - (\gamma_0(T) - z_d, \gamma_0(T)) - \frac{1}{2} \int_0^T (Nu_0, u_0) dt \\
 &= -\frac{1}{2} |\gamma_0(T)|^2 + \frac{1}{2} |z_d|^2 - \frac{1}{2} \int_0^T (Nu_0, u_0) dt \\
 &= K(\gamma_0, u_0).
 \end{aligned} \tag{3.15}$$

(ii) We show that $J(\gamma, u) \geq J(\gamma_0, u_0)$ under (3.11) for (γ, u, p) and under (3.11), (3.12), and (3.13) for (γ_0, u_0, p_0) :

$$\begin{aligned}
 J(\gamma, u) - J(\gamma_0, u_0) &\geq (\gamma_0(T) - z_d, \gamma(T) - \gamma_0(T)) + \int_0^T (Nu_0, u - u_0) dt \\
 &= (\gamma_0(T) - z_d, \gamma(T) - \gamma_0(T)) + \int_0^T (Nu_0, u - u_0) dt \\
 &\quad - \int_0^T (p'' - A_2(t)p' + (A_1(t) - A_2'(t))p, \gamma - \gamma_0) dt \\
 &= (\gamma_0(T) - z_d, \gamma(T) - \gamma_0(T)) \\
 &\quad + \int_0^T (Nu_0, u - u_0) dt - (\gamma_0(T) - z_d, \gamma(T) - \gamma_0(T)) \\
 &\quad - \int_0^T (p_0, \gamma'' - \gamma_0'' + A_2(t)\gamma' - A_2(t)\gamma_0' \\
 &\quad \quad + A_1(t)\gamma - A_1(t)\gamma_0) dt \\
 &= \int_0^T (-p_0 + Nu_0, u - u_0) dt \\
 &\geq 0.
 \end{aligned} \tag{3.16}$$

(iii) We have to check that $K(\gamma, u) \leq K(\gamma_0, u_0)$ under (3.12) for (γ, u, p) :

$$\begin{aligned}
 J(\gamma_0, u_0) - J(\gamma, u) &\geq (\gamma(T) - z_d, \gamma_0(T) - \gamma(T)) + \int_0^T (Nu, u_0 - u) dt \\
 &= (\gamma(T) - z_d, \gamma_0(T) - \gamma(T)) + \int_0^T (Nu, u_0 - u) dt \\
 &\quad - \int_0^T (\gamma_0'' + A_2(t)\gamma_0' + A_1(t)\gamma_0 - u_0, p_0 - p) dt
 \end{aligned}$$

$$\begin{aligned}
&= -(\gamma(T) - z_d, \gamma(T)) - \int_0^T (Nu, u) + (\gamma_0(T) - z_d, \gamma_0(T)) dt \\
&\quad + \int_0^T (Nu_0, u_0) dt + \int_0^T (-p + Nu, u_0) dt \\
&\quad + \int_0^T (p_0 - Nu_0, u_0) dt \\
&\geq -(\gamma(T) - z_d, \gamma(T)) - \int_0^T (Nu, u) dt \\
&\quad + (\gamma_0(T) - z_d, \gamma_0(T)) + \int_0^T (Nu_0, u_0) dt.
\end{aligned} \tag{3.17}$$

This shows that

$$K(\gamma_0, u_0) \geq K(\gamma, u). \tag{3.18}$$

Therefore, [Theorem 3.2](#) is proved. $\square$

When the observation $z(u)$ is given by $z(u) = \gamma'(u)$, the cost function is defined as

$$J(\gamma', u) = \frac{1}{2} \int_0^T |\gamma'(u) - z_d|^2 dt + \frac{1}{2} \int_0^T (Nu, u) dt. \tag{3.19}$$

We will consider the following systems:

$$\begin{aligned}
&\gamma''(u; t) + A_2(t)\gamma'(u; t) + A_1(t)\gamma(u; t) = u \quad \text{in } (0, T), \\
&\gamma(u; 0) = \gamma_0, \quad \gamma'(u; 0) = \gamma_1 \quad \text{in } \Omega, \\
&u \geq 0 \quad \text{a.e. in } Q, \quad \gamma(0) \geq 0, \quad \gamma'(0) \geq 0, \quad \gamma = 0 \quad \text{on } \Sigma,
\end{aligned} \tag{3.20}$$

$$\begin{aligned}
&p''(u; t) - A_2(t)p'(u; t) + A_1(t)p(u; t) + \int_t^T A_1(\sigma)p(u; \sigma) d\sigma \\
&\quad = \gamma'(u; t) - z_d \quad \text{in } Q,
\end{aligned} \tag{3.21}$$

$$\begin{aligned}
&p = 0 \quad \text{on } \Sigma, \quad p(u; T) = 0, \quad p'(u; T) = 0, \\
&-p' + Nu \geq 0 \quad \text{in } Q, \quad p(0) \geq 0, \quad p'(0) \geq 0,
\end{aligned}$$

$$u(-p + Nu) = 0$$

$$\begin{aligned}
&p(0)\gamma(0) = 0, \quad \gamma'(0)p(0) = 0, \quad \gamma(0)p'(0) = 0, \\
&\gamma(0) \int_0^T A'(\sigma)p(\sigma) d\sigma = 0 \quad \text{in } \Omega.
\end{aligned} \tag{3.22}$$

THEOREM 3.3. Let $J = (1/2) \int_0^T |\gamma' - z_d|^2 dt + \int_0^T (1/2)(Nu, u) dt$ and $K = -(1/2) \int_0^T |\gamma'|^2 dt + (1/2) \int_0^T |z_d|^2 dt - (1/2) \int_0^T (Nu, u) dt$. Assume that γ_0, u_0 , and p_0 satisfy (3.20), (3.21), and (3.22), respectively, γ and u in J satisfy (3.20),

and y and u in K satisfy (3.21). Then

$$\inf_{(3.20)} J = J(y'_0, u_0) = K(y'_0, u_0) = \sup_{(3.21)} K. \quad (3.23)$$

PROOF. (i) First, we claim that $J(y'_0, u_0) = K(y'_0, u_0)$:

$$\begin{aligned} J(y'_0, u_0) &= J(y'_0, u_0) + \int_0^T (p'_0, u_0) dt - \int_0^T (Nu_0, u_0) dt \\ &= J(y'_0, u_0) + \int_0^T (p'_0, y''_0 + A_2(t)y'_0 + A_1(t)y_0) dt - \int_0^T (Nu_0, u_0) dt \\ &= J(y'_0, u_0) + \int_0^T \left(-p''_0 + A_2(t)p'_0 - A_1(t)p_0 \right. \\ &\quad \left. - \int_t^T A'(\sigma)p(\sigma)d\sigma, y' \right) dt - \int_0^T (Nu_0, u_0) dt \\ &= J(y'_0, u_0) - \int_0^T (y'_0 - z_d) dt - \int_0^T (Nu_0, u_0) dt \\ &= -\frac{1}{2} \int_0^T |y'_0|^2 dt + \frac{1}{2} \int_0^T |z_d|^2 dt - \frac{1}{2} \int_0^T (Nu_0, u_0) dt. \end{aligned} \quad (3.24)$$

(ii) Second, we must show that $\inf J = J(y'_0, u_0)$:

$$\begin{aligned} J(y', u) - J(y'_0, u_0) &\geq \int_0^T (y'_0 - z_d, y' - y'_0) dt + \int_0^T (Nu_0, u - u_0) dt \\ &= \int_0^T \left(p''_0 - A_2(t)p'_0 + A_1(t)p_0 \right. \\ &\quad \left. + \int_t^T A'_1(\sigma)p_0(\sigma)d\sigma, y' - y'_0 \right) dt \\ &\quad + \int_0^T (Nu_0, u - u_0) dt \\ &= \int_0^T (p'_0, -(y'' - y''_0) - A_2(t)(y' - y'_0) - A_1(t)(y - y_0)) dt \\ &\quad + \int_0^T (Nu_0, u - u_0) dt \\ &= \int_0^T (-p'_0 + Nu_0, u - u_0) dt \\ &\geq 0. \end{aligned} \quad (3.25)$$

(iii) Finally, we check that $\sup K = K(y'_0, u_0)$:

$$\begin{aligned}
 & J(y'_0, u_0) - J(y', u) \\
 & \geq \int_0^T (y' - z_d, y'_0, y') dt + \int_0^T (Nu, u_0 - u) dt \\
 & = \int_0^T \left(p'' - A_2(t)p' + A_1(t)p + \int_t^T A'_1(\sigma)p(\sigma) d\sigma, y'_0 - y' \right) dt \\
 & \quad - \int_0^T (y''_0 + A_2(t)y'_0 + A_1(t)y_0 - u_0, p'_0 - p') dt + \int_0^T (Nu, u_0 - u) dt \\
 & = - \int_0^T (y' - z_d, y') dt + \int_0^T (p', -y''_0 - A_2(t)y'_0 - A_1(t)y_0) dt \\
 & \quad - \int_0^T \left(y'_0 - p'_0 + A_2(t)p'_0 - A_1(t)p_0 - \int_t^T A'(\sigma)p_0(\sigma) d\sigma \right) dt \\
 & \quad + \int_0^T (Nu, u_0) dt - \int_0^T (Nu, u) dt + \int_0^T (u_0, p') dt \\
 & \quad + \int_0^T (u_0, p'_0) dt - \int_0^T (u_0, p') dt \\
 & = - \int_0^T (y' - z_d, y') dt - \int_0^T (Nu, u) dt + \int_0^T (y'_0, y'_0 - z_d) dt \\
 & \quad + \int_0^T (Nu, u_0) dt + \int_0^T (u_0, p'_0) dt \\
 & \quad - \int_0^T (u_0, p') dt + \int_0^T (Nu_0, u_0) dt - \int_0^T (Nu_0, u_0) dt \\
 & \geq - \int_0^T (y' - z_d, y') dt - \int_0^T (Nu, u) dt \\
 & \quad + \int_0^T (y'_0 - z_d, y'_0) dt + \int_0^T (Nu_0, u_0) dt. \tag{3.26}
 \end{aligned}$$

This implies that

$$K(y'_0, u_0) \geq K(y, u). \tag{3.27}$$

So, we claimed [Theorem 3.3](#). $\square$

In this case, we observe the terminal value $y'(u; T)$. Since the observation $z(u)$ is given by $y'(u; T)$, the cost functional is given as

$$J(y', u) = \frac{1}{2} |y'(u; T) - z_d|^2 + \frac{1}{2} \int_0^T (Nu, u) dt. \tag{3.28}$$

We introduce the following systems:

$$\begin{aligned} \gamma''(u;t) + A_2(t)\gamma'(u;t) + A_1(t)\gamma(u;t) &= u \quad \text{in } Q, \\ \gamma(u;0) &= \gamma_0, \quad \gamma'(u;0) = \gamma_1 \quad \text{in } \Omega, \end{aligned} \quad (3.29)$$

$$u \geq 0 \quad \text{a.e. in } Q, \quad \gamma(0) \geq 0, \quad \gamma'(0) \geq 0 \quad \text{in } \Omega, \quad \gamma = 0 \quad \text{on } \Sigma,$$

$$\begin{aligned} p''(u;t) - A_2(t)p'(u;t) + (A_1(t) - A_2'(t))p(u;t) &= 0 \quad \text{in } Q, \\ p &= 0 \quad \text{on } \Sigma, \end{aligned} \quad (3.30)$$

$$p(u;T) = \gamma'(T) - z_d, \quad p'(u;T) = A_2(T)\gamma'(T) - z_d,$$

$$p + Nu \geq 0 \quad \text{in } Q,$$

$$p(0) \geq 0, \quad p'(0) \geq 0 \quad \text{in } \Omega,$$

$$u(p + Nu) = 0, \quad (3.31)$$

$$p(0)\gamma(0) = 0, \quad \gamma'(0)p(0) = 0, \quad \gamma(0)p'(0) = 0 \quad \text{in } \Omega.$$

THEOREM 3.4. Let $J = (1/2)|\gamma'(T) - z_d|^2 + (1/2)\int_0^T (Nu, u)dt$ and $K = -(1/2)|\gamma'(T)|^2 + (1/2)|z_d|^2 - (1/2)\int_0^T (Nu, u)dt$. Assume that γ_0 , u_0 , and p_0 satisfy (3.29), (3.30), and (3.31), respectively, γ and u in J satisfy (3.29), and γ and u in K satisfy (3.30). Then

$$\inf_{(3.29)} J = J(\gamma'_0, u_0) = K(\gamma'_0, u_0) = \sup_{(3.30)} K. \quad (3.32)$$

PROOF. (i) First, we show that $J = K$ at (γ'_0, u_0, p_0) :

$$\begin{aligned} J(\gamma'_0, u_0) &= J(\gamma'_0, u_0) - \int_0^T (u_0, p_0)dt - \int_0^T (u_0, Nu_0)dt \\ &= J(\gamma'_0, u_0) - \int_0^T (\gamma_0'' + A_2(t)\gamma_0' + A_1(t)\gamma_0, p_0)dt - \int_0^T (Nu_0, u_0)dt \\ &= \frac{1}{2}|\gamma_0'(T) - z_d|^2 + \frac{1}{2}\int_0^T (Nu_0, u_0)dt - (\gamma_0'(T) - z_d, \gamma_0'(T)) \\ &\quad - \int_0^T (Nu_0, u_0)dt \\ &= -\frac{1}{2}|\gamma_0'(T)|^2 + \frac{1}{2}|z_d|^2 - \frac{1}{2}\int_0^T (Nu_0, u_0)dt \\ &= K(\gamma'_0, u_0). \end{aligned} \quad (3.33)$$

(ii) Second, we check that $J(\mathbf{y}', u) \geq J(\mathbf{y}'_0, u_0)$:

$$\begin{aligned}
 & J(\mathbf{y}', u) - J(\mathbf{y}'_0, u_0) \\
 & \geq (\mathbf{y}'_0(T) - z_d, \mathbf{y}'(T) - \mathbf{y}'_0(T)) + \int_0^T (Nu_0, u - u_0) dt \\
 & = (\mathbf{y}'_0(T) - z_d, \mathbf{y}'(T) - \mathbf{y}'_0(T)) + \int_0^T (Nu_0, u - u_0) dt \\
 & \quad + \int_0^T (\mathbf{p}'' - A_2(t)\mathbf{p}' + (A_1(t) - A'_2(t))\mathbf{p}, \mathbf{y} - \mathbf{y}_0) dt \\
 & = \int_0^T (\mathbf{p}, \mathbf{y}'' - \mathbf{y}''_0 + A_2(t)\mathbf{y}' - A_2(t)\mathbf{y}'_0 + A_1(t)\mathbf{y} - A_1(t)\mathbf{y}_0) dt \\
 & \quad + \int_0^T (Nu_0, u - u_0) dt \\
 & = \int_0^T (\mathbf{p} + Nu_0, u - u_0) dt \\
 & \geq 0.
 \end{aligned} \tag{3.34}$$

(iii) Finally, we claim that $K(\mathbf{y}', u) \leq K(\mathbf{y}'_0, u_0)$:

$$\begin{aligned}
 & J(\mathbf{y}'_0, u_0) - J(\mathbf{y}', u) \\
 & \geq (\mathbf{y}'_0(T) - z_d, \mathbf{y}'(T) - \mathbf{y}'_0(T)) + \int_0^T (Nu_0, u - u_0) dt \\
 & = (\mathbf{y}'_0(T) - z_d, \mathbf{y}'(T) - \mathbf{y}'_0(T)) + \int_0^T (Nu_0, u - u_0) dt \\
 & \quad + \int_0^T (\mathbf{y}''_0 + A_2(t)\mathbf{y}'_0 + A_1(t)\mathbf{y}_0 - u_0, \mathbf{p}_0 - \mathbf{p}) dt \\
 & = -(\mathbf{y}'(T) - z_d, \mathbf{y}'(T)) - \int_0^T (Nu, u) dt + (\mathbf{y}'_0(T) - z_d, \mathbf{y}'(T)) \\
 & \quad + \int_0^T (Nu_0, u_0) dt + \int_0^T (Nu + \mathbf{p}, u_0) dt - \int_0^T (\mathbf{p}_0 + Nu_0, u_0) dt \\
 & \geq -(\mathbf{y}'(T) - z_d, \mathbf{y}'(T)) - \int_0^T (Nu, u) dt + (\mathbf{y}'_0(T) - z_d, \mathbf{y}'(T)) \\
 & \quad + \int_0^T (Nu_0, u_0) dt.
 \end{aligned} \tag{3.35}$$

This implies that

$$K(\mathbf{y}'_0, u_0) \geq K(\mathbf{y}', u). \tag{3.36}$$

This completes the proof. $\square$

Acknowledgment. This work was supported by the Korea Research Foundation Grant KRF-2001-037-DA0010.

REFERENCES

- [1] W. L. Chan, *Duality in the optimal control of non-well-posed parabolic equations with positive controls*, J. Math. Anal. Appl. **107** (1985), no. 2, 509–519.
- [2] J.-L. Lions, *Optimal Control of Systems Governed by Partial Differential Equations*, Die Grundlehren der mathematischen Wissenschaften, vol. 170, Springer-Verlag, New York, 1971.
- [3] S.-I Nakagiri and J. Ha, *Optimal control problems for hyperbolic distributed parameter systems with damping terms*, Ph.D. thesis, Department of Applied Mathematics, Kobe University, Japan, 1996.
- [4] J. Y. Park and M. J. Lee, *Duality in the optimal control problems for hyperbolic distributed parameter systems with damping terms*, J. Math. Anal. Appl. **227** (1998), no. 2, 449–461.
- [5] ———, *Duality in the optimal control of distributed parameter systems governed by hyperbolic equations*, Indian J. Pure Appl. Math. **31** (2000), no. 4, 451–460.
- [6] ———, *Duality in the optimal control of hyperbolic equations with positive controls*, Int. J. Math. Math. Sci. **23** (2000), no. 3, 181–188.
- [7] S. Tanimoto, *Duality in the optimal control of non-well-posed distributed systems*, J. Math. Anal. Appl. **171** (1992), no. 1, 277–287.

Mi Jin Lee: Department of Mathematics, Pusan National University (PNU), Pusan 609-735, Korea

E-mail address: jin8719@hanmail.net

Jong Yeoul Park: Department of Mathematics, Pusan National University (PNU), Pusan 609-735, Korea

E-mail address: jyepark@pusan.ac.kr

Young Chel Kwon: Department of Mathematics, Dong-A University, Pusan 604-714, Korea

E-mail address: yckwon@daunet.donga.ac.kr

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru