

OPTIMAL ORDER YIELDING DISCREPANCY PRINCIPLE FOR SIMPLIFIED REGULARIZATION IN HILBERT SCALES: FINITE-DIMENSIONAL REALIZATIONS

SANTHOSH GEORGE and M. THAMBAN NAIR

Received 13 June 2003

Simplified regularization using finite-dimensional approximations in the setting of Hilbert scales has been considered for obtaining stable approximate solutions to ill-posed operator equations. The derived error estimates using an a priori and a posteriori choice of parameters in relation to the noise level are shown to be of optimal order with respect to certain natural assumptions on the ill posedness of the equation. The results are shown to be applicable to a wide class of spline approximations in the setting of Sobolev scales.

2000 Mathematics Subject Classification: 65R10, 65J10, 46E35, 47A50.

1. Introduction. Many of the inverse problems that occur in science and engineering are ill posed, in the sense that a unique solution that depends continuously on the data does not exist. A typical example of an ill-posed equation that often occurs in practical problems, such as in geological prospecting, computer tomography, steel industry, and so forth, is the Fredholm integral equation of the first kind (cf. [2, 6, 8]). Many such problems can be put in the form of an operator equation $Ax = y$, where $A : X \rightarrow Y$ is a bounded linear operator between Hilbert spaces X and Y with its range $R(A)$ not closed in Y .

Regularization methods are to be employed for obtaining a stable approximate solution for an ill-posed problem. Tikhonov regularization is a simple and widely used procedure to obtain stable approximate solutions to an ill-posed operator equation (2.1). In order to improve the error estimates available in Tikhonov regularization, Natterer [17] carried out error analysis in the framework of Hilbert scales. Subsequently, many authors extended, modified, and generalized Natterer's work to obtain error bounds under various contexts (cf. Neubauer [18], Hegland [7], Schröter and Tautenhahn [20], Mair [10], Nair et al. [16], and Nair [13, 15]). Finite-dimensional realizations of the Hilbert scales approach has been considered by Engl and Neubauer [3].

If $Y = X$ and A itself is a positive selfadjoint operator, then the simplified regularization introduced by Lavrentiev is better suited than Tikhonov regularization in terms of speed of convergence and condition numbers of the resulting equations in the case of finite-dimensional approximations (cf. Schock [19]).

In [4], the authors introduced the Hilbert scales variant of the simplified regularization and obtained error estimates under a priori and a posteriori parameter choice strategies which are optimal in the sense of the "best possible worst error" with respect to certain source set. Recently (cf. [5]), the authors considered a new discrepancy

principle yielding optimal rates which does not involve certain restrictive assumptions as in [4]. The purpose of this paper is to obtain a finite-dimensional realization of the results in [5].

2. Preliminaries. Let H be a Hilbert space and $A : H \rightarrow H$ a positive, bounded self-adjoint operator on H . The inner product and the corresponding norm are denoted by $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$, respectively. Recall that A is said to be a positive operator if $\langle Ax, x \rangle \geq 0$ for every $x \in H$. For $\gamma \in R(A)$, the range of A , consider the operator equation

$$Ax = \gamma. \tag{2.1}$$

Let \hat{x} be the minimal norm solution of (2.1). It is well known that if $R(A)$ is not closed in H , then the problem of solving (2.1) for \hat{x} is ill posed in the sense that small perturbations in the data γ can cause large deviations in the solution. A prototype of an ill-posed equation (2.1) is an integral equation of the first kind,

$$\int_0^1 k(\xi, t)x(t)dt = \gamma(\xi), \quad 0 \leq \xi \leq 1, \tag{2.2}$$

where $k(\cdot, \cdot)$ is a nondegenerate kernel which is square integrable, that is,

$$\int_0^1 \int_0^1 |k(\xi, t)|^2 dt d\xi < \infty, \tag{2.3}$$

satisfying $k(\xi, t) = k(t, \xi)$ for all ξ, t in $[0, 1]$, and such that the eigenvalues of the corresponding integral operator $A : L^2[0, 1] \rightarrow L^2[0, 1]$,

$$(Ax)(\xi) = \int_0^1 k(\xi, t)x(t)dt, \quad 0 \leq \xi \leq 1, \tag{2.4}$$

are all nonnegative (cf. [14]). For example, one of the important ill-posed problems which arise in applications is the *backward heat equation* problem: the problem is to determine the initial temperature $\varphi_0 := u(\cdot, 0)$ from the measurements of the final temperature $\varphi_T := u(\cdot, T)$, where $u(\xi, t)$ satisfies

$$\begin{aligned} u_t - u_{\xi\xi} &= 0, & (\xi, t) &\in (0, 1) \times (0, T), \\ u(0, t) &= u(1, t) = 0, & t &\in [0, T]. \end{aligned} \tag{2.5}$$

We recall from elementary theory of partial differential equations that the solution $u(\xi, t)$ of the above heat equation is given by (cf. Weinberger [23])

$$u(\xi, t) = \sum_{n=1}^{\infty} \hat{\varphi}_0(n)e^{-n^2\pi^2t} \sin(n\pi\xi), \tag{2.6}$$

where $\hat{\varphi}_0(n)$ for $n \in \mathbb{N}$ are the Fourier coefficients of the initial temperature $\varphi_0(\xi) := u(\xi, 0)$. Hence,

$$u(\xi, T) = \sum_{n=1}^{\infty} \hat{\varphi}_0(n)e^{-n^2\pi^2T} \sin(n\pi\xi). \tag{2.7}$$

The above equation can be written as

$$\varphi_T(s) = \sum_{n=1}^{\infty} e^{-n^2\pi^2T} \langle \varphi_0, u_n \rangle u_n(\xi) \quad \text{with } u_n(\xi) = \sqrt{2} \sin(n\pi\xi). \quad (2.8)$$

Thus the problem is to solve the operator equation

$$A\varphi_0 = \varphi_T, \quad (2.9)$$

where $A : L^2[0, 1] \rightarrow L^2[0, 1]$ is the operator defined by

$$(A\varphi)(\xi) = \sum_{n=1}^{\infty} e^{-n^2\pi^2T} \langle \varphi, u_n \rangle u_n(\xi) = \int_0^1 k(\xi, t) \varphi(t) dt, \quad 0 \leq \xi \leq 1, \quad (2.10)$$

where

$$k(\xi, t) := \sum_{n=1}^{\infty} e^{-n^2\pi^2T} u_n(\xi) u_n(t). \quad (2.11)$$

Note that the above integral operator is compact, positive, and selfadjoint with positive eigenvalues $e^{-n^2\pi^2T}$ and corresponding eigenvectors $u_n(\cdot)$ for $n \in \mathbb{N}$.

For considering the regularization of (2.1) in the setting of Hilbert scales, we consider a Hilbert scale $\{H_t\}_{t \in \mathbb{R}}$ generated by a strictly positive operator $L : D(L) \rightarrow H$ with its domain $D(L)$ dense in H satisfying

$$\|Lx\| \geq \|x\|, \quad x \in D(L). \quad (2.12)$$

By the operator L being strictly positive, we mean that $\langle Lx, x \rangle > 0$ for all nonzero $x \in H$. Recall (cf. [9]) that the space H_t is the completion of $D := \bigcap_{k=0}^{\infty} D(L^k)$ with respect to the norm $\|x\|_t$, induced by the inner product

$$\langle u, v \rangle_t = \langle L^t u, L^t v \rangle, \quad u, v \in D. \quad (2.13)$$

Moreover, if $\beta \leq \gamma$, then the embedding $H_\gamma \hookrightarrow H_\beta$ is continuous, and therefore the norm $\|\cdot\|_\beta$ is also defined in H_γ and there is a constant $c_{\beta,\gamma}$ such that

$$\|x\|_\beta \leq c_{\beta,\gamma} \|x\|_\gamma \quad \forall x \in H_\beta. \quad (2.14)$$

An important inequality that we require in the analysis is the *interpolation inequality*

$$\|x\|_\lambda \leq \|x\|_r^\theta \|x\|_t^{1-\theta}, \quad x \in H_t, \quad (2.15)$$

where

$$r \leq \lambda \leq t, \quad \theta = \frac{t-\lambda}{t-r}, \quad (2.16)$$

and the *moment inequality*

$$\|B^u x\| \leq \|B^v x\|^{u/v} \|x\|^{1-u/v}, \quad 0 \leq u \leq v, \quad (2.17)$$

where B is a positive selfadjoint operator (cf. [2]).

We assume that the ill-posed nature of the operator A is related to the Hilbert scale $\{H_t\}_{t \in \mathbb{R}}$ according to the relation

$$c_1 \|x\|_{-a} \leq \|Ax\| \leq c_2 \|x\|_{-a}, \quad x \in H, \tag{2.18}$$

for some positive reals a, c_1 , and c_2 .

For the example of the integral operator considered in (2.4), one may take L to be defined by

$$Lx := \sum_{j=1}^{\infty} j^2 \langle x, u_j \rangle u_j, \tag{2.19}$$

where $u_j(t) := \sqrt{2} \sin(j\pi t)$, $j \in \mathbb{N}$ with domain of L as

$$D(L) := \left\{ x \in L^2[0, 1] : \sum_{j=1}^{\infty} j^4 |\langle x, u_j \rangle|^2 < \infty \right\}. \tag{2.20}$$

In this case, it can be seen that

$$H_t = \left\{ x \in L^2[0, 1] : \sum_{j=1}^{\infty} j^{4t} |\langle x, u_j \rangle|^2 < \infty \right\} \tag{2.21}$$

and the constants a, c_1 , and c_2 in (2.18) are given by $a = 1, c_1 = c_2 = 1/\pi^2$ (see Schröter and Tautenhahn [20, Section 4]).

The regularized approximation of \hat{x} , considered in [4] is the solution of the well-posed equation

$$(A + \alpha L^s)x_\alpha = y, \quad \alpha > 0, \tag{2.22}$$

where s is a fixed nonnegative real number. Note that if $D(L) = X$ and $L = I$, then the above procedure is the simplified or Lavrentiev regularization.

Suppose the data y is known only approximately, say \tilde{y} in place of y with $\|y - \tilde{y}\| \leq \delta$ for a known error level $\delta > 0$. Then, in place of (2.22), we have

$$(A + \alpha L^s)\tilde{x}_\alpha = \tilde{y}. \tag{2.23}$$

It can be seen that the solution \tilde{x}_α of the above equation is the unique minimizer of the function

$$x \mapsto \langle Ax, x \rangle - 2\langle \tilde{y}, x \rangle + \alpha \langle L^s x, x \rangle, \quad x \in D(L). \tag{2.24}$$

One of the crucial results for proving the results in [4, 5] as well as the results in this paper is the following proposition, where the functions f and g are defined by

$$f(t) = \min \{c_1^t, c_2^t\}, \quad g(t) = \max \{c_1^t, c_2^t\}, \quad t \in \mathbb{R}, |t| \leq 1, \tag{2.25}$$

respectively, with c_1, c_2 as in (2.18).

PROPOSITION 2.1 (cf. [4, Proposition 3.1]). *For $s > 0$ and $|\nu| \leq 1$,*

$$f\left(\frac{\nu}{2}\right) \|x\|_{-\nu(s+a)/2} \leq \|A_s^{\nu/2} x\| \leq g\left(\frac{\nu}{2}\right) \|x\|_{-\nu(s+a)/2}, \quad x \in H, \quad (2.26)$$

where $A_s = L^{-s/2} A L^{-s/2}$.

Using the above proposition, the following result has been proved by George and Nair [4].

THEOREM 2.2 (cf. [4, Theorem 3.2]). *Suppose $\hat{x} \in H_t$, $0 < t \leq s + a$, and $\alpha > 0$, and \tilde{x}_α is as in (2.23). Then*

$$\|\hat{x} - \tilde{x}_\alpha\| \leq \phi(s, t) \alpha^{t/(s+a)} \|\hat{x}\|_t + \psi(s) \alpha^{-a/(s+a)} \delta, \quad (2.27)$$

where

$$\phi(s, t) = \frac{g((s-2t)/(2s+2a))}{f(s/(2s+2a))}, \quad \psi(s) = \frac{g(-s/(2s+2a))}{f(s/(2s+2a))}. \quad (2.28)$$

In particular, if $\alpha = c_0 \delta^{(s+a)/(t+a)}$ for some constant $c_0 > 0$, then

$$\|\hat{x} - \tilde{x}_\alpha\| \leq \eta(s, t) \delta^{t/(t+a)}, \quad (2.29)$$

where

$$\eta(s, t) = \max \left\{ \phi(s, t) \|x\|_t c_0^{t/(t+a)}, \psi(s) c_0^{-a/(s+a)} \right\}. \quad (2.30)$$

For proposing a finite-dimensional realization, we consider a family $\{S_h : h > 0\}$ of finite-dimensional subspaces of H_k for some $k \geq s$, and consider the minimizer $\tilde{x}_{\alpha, h}$ of the map defined in (2.24) when x varies over S_h . Equivalently, $\tilde{x}_{\alpha, h}$ is the unique element in S_h satisfying the equation

$$\langle (A + \alpha L^s) \tilde{x}_{\alpha, h}, \varphi \rangle = \langle \hat{y}, \varphi \rangle \quad \forall \varphi \in S_h. \quad (2.31)$$

As in Engl and Neubauer [3], we assume the following approximation properties for S_h .

There exists a constant $\kappa > 0$ such that for every $u \in H_r$ with $r > k \geq s$,

$$\inf \{ \|u - \varphi\|_k : \varphi \in S_h \} \leq \kappa h^{r-k} \|u\|_r, \quad h > 0. \quad (2.32)$$

As already exemplified in [3], the above assumption is general enough to include a wide variety of approximation spaces, such as spline spaces and finite element spaces.

We will also make use of the following result from Engl and Neubauer [3, Lemma 2.2].

LEMMA 2.3. *Under the assumption (2.32), there exists a constant $c > 0$ such that for every $u \in H_s$ and $h > 0$,*

$$\inf_{\varphi \in S_h} \{ h^{-a/2} \|u - \varphi\|_{-a/2} + h^{s/2} \|u - \varphi\|_{s/2} \} \leq c h^s \|u\|_s. \quad (2.33)$$

3. General error estimates. For a fixed $s > 0$, let \tilde{x}_α and $\tilde{x}_{\alpha,h}$ be as in (2.23) and (2.31), respectively. We will obtain estimate for $\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|$ so that we get an estimate for $\|\hat{x} - \tilde{x}_{\alpha,h}\|$ using Theorem 2.2 and the relation

$$\|\hat{x} - \tilde{x}_{\alpha,h}\| \leq \|\hat{x} - \tilde{x}_\alpha\| + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|. \tag{3.1}$$

In view of the interpolation inequality (2.15), by taking $\rho = -a/2$, $\tau = s/2$, and $\lambda = 0$ in (2.15), we get

$$\|x\| \leq \|x\|_{-a/2}^{s/(s+a)} \|x\|_{s/2}^{a/(s+a)}, \quad x \in H_{s/2}. \tag{3.2}$$

Thus, we can deduce an estimate for $\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|$ once we have estimates for $\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{-a/2}$ and $\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{s/2}$. For this purpose, we first prove the following.

LEMMA 3.1. *Let \tilde{x}_α and $\tilde{x}_{\alpha,h}$ be as in (2.23) and (2.31), respectively. Then*

$$\|A^{1/2}(\tilde{x}_\alpha - \tilde{x}_{\alpha,h})\|^2 + \alpha\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{s/2}^2 = \inf_{\varphi \in S_h} \{ \|A^{1/2}(\tilde{x}_\alpha - \varphi)\|^2 + \alpha\|\tilde{x}_\alpha - \varphi\|_{s/2}^2 \}. \tag{3.3}$$

PROOF. It can be seen (cf. [16]) that

$$\langle u, v \rangle_* := \langle Au, v \rangle + \alpha \langle L^s u, v \rangle, \quad u, v \in D(L), \tag{3.4}$$

defines a complete inner product on $D(L)$. Let $\|\cdot\|_*$ be the norm induced by $\langle \cdot, \cdot \rangle_*$, that is,

$$\|u\|_* = (\langle Au, u \rangle + \alpha \langle L^s u, u \rangle)^{1/2} = (\|A^{1/2}u\|^2 + \alpha\|u\|_{s/2}^2)^{1/2}. \tag{3.5}$$

Let X be the space $D(L)$ with the inner product $\langle \cdot, \cdot \rangle_*$ and let P_h be the orthogonal projection of X onto the space S_h . Then from (2.23) and (2.31) we have

$$\langle (A + \alpha L^s)(\tilde{x}_\alpha - \tilde{x}_{\alpha,h}), \varphi \rangle = 0 \quad \forall \varphi \in S_h, \tag{3.6}$$

that is,

$$\langle \tilde{x}_\alpha - \tilde{x}_{\alpha,h}, \varphi \rangle_* = 0 \quad \forall \varphi \in S_h. \tag{3.7}$$

Hence

$$P_h(\tilde{x}_\alpha - \tilde{x}_{\alpha,h}) = 0 \tag{3.8}$$

so that

$$\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_* = \inf_{\varphi \in S_h} \|(\tilde{x}_\alpha - \tilde{x}_{\alpha,h}) - \varphi\|_* = \inf_{\varphi \in S_h} \|\tilde{x}_\alpha - \varphi\|_*. \tag{3.9}$$

Now the result follows using the definition of $\|\cdot\|_*$. □

Next we obtain estimate for $\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|$ using the estimates for $\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{-a/2}$ and $\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{s/2}$. We will use the notation

$$A_s := L^{-s/2} A L^{-s/2} \quad (3.10)$$

and observe that for $\alpha > 0$,

$$(A + \alpha L^s)x = L^{s/2}(A_s + \alpha I)L^{s/2}x \quad \forall x \in H_s, \quad (3.11)$$

THEOREM 3.2. *Suppose $\hat{x} \in H_t$ and assumption (2.32) holds, and let \tilde{x}_α and $\tilde{x}_{\alpha,h}$ be as in (2.23) and (2.31), respectively. Then*

$$\begin{aligned} & \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\ & \leq f\left(\frac{1}{2}\right)^{-s/(s+a)} \max\{\mathcal{F}(s,a), \mathcal{G}(s,t,a)\} \Phi(s,h,\alpha) \alpha^{-a/(2s+2a)} \left(\frac{\delta}{\alpha} + \alpha^{(t-s)/(s+a)}\right) h^s, \end{aligned} \quad (3.12)$$

where f and g are as in (2.25), and

$$\begin{aligned} \mathcal{F}(s,a) &= \frac{g(-s/(2s+2a))}{f(-s/(2s+2a))}, \quad \mathcal{G}(s,t,a) = \frac{g((s-2t)/(2s+2a))}{f(-s/(2s+2a))} \|\hat{x}\|_t, \\ \Phi(s,h,\alpha) &= c \max\left\{g\left(\frac{1}{2}\right) h^{a/2}, \alpha^{1/2} h^{-s/2}\right\}. \end{aligned} \quad (3.13)$$

PROOF. First we prove

$$\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{-a/2} \leq \frac{1}{f(1/2)} \Phi(s,h,\alpha) h^s \|\tilde{x}_\alpha\|_s, \quad (3.14)$$

$$\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{s/2} \leq \Phi(s,h,\alpha) \alpha^{-1/2} h^s \|\tilde{x}_\alpha\|_s, \quad (3.15)$$

$$\|\tilde{x}_\alpha\|_s \leq \mathcal{F}(s,a) \alpha^{-1} \delta + \mathcal{G}(s,t,a) \alpha^{(t-s)/(s+a)} \quad (3.16)$$

with $\mathcal{F}(s,a)$, $\mathcal{G}(s,t,a)$, and $\Phi(s,h,\alpha)$ as in the statement of the theorem.

By Lemma 3.1 and Proposition 2.1, it follows that

$$\begin{aligned} & f\left(\frac{1}{2}\right)^2 \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{-a/2}^2 + \alpha \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{s/2}^2 \\ & \leq \inf_{\varphi \in S_h} \left\{ g\left(\frac{1}{2}\right)^2 \|\tilde{x}_\alpha - \varphi\|_{-a/2}^2 + \alpha \|\tilde{x}_\alpha - \varphi\|_{s/2}^2 \right\}. \end{aligned} \quad (3.17)$$

Note that

$$\begin{aligned} & g\left(\frac{1}{2}\right)^2 \|\tilde{x}_\alpha - \varphi\|_{-a/2}^2 + \alpha \|\tilde{x}_\alpha - \varphi\|_{s/2}^2 \\ & \leq \left[g\left(\frac{1}{2}\right) \|\tilde{x}_\alpha - \varphi\|_{-a/2} + \alpha^{1/2} \|\tilde{x}_\alpha - \varphi\|_{s/2} \right]^2. \end{aligned} \quad (3.18)$$

But

$$\begin{aligned} g\left(\frac{1}{2}\right) \|\tilde{x}_\alpha - \varphi\|_{-a/2} + \alpha^{1/2} \|\tilde{x}_\alpha - \varphi\|_{s/2} \\ \leq \omega_{h,\alpha,s} \left[h^{-a/2} \|\tilde{x}_\alpha - \varphi\|_{-a/2} + h^{s/2} \|\tilde{x}_\alpha - \varphi\|_{s/2} \right], \end{aligned} \tag{3.19}$$

where

$$\omega(h, \alpha, s) := \max \left\{ g\left(\frac{1}{2}\right) h^{a/2}, \alpha^{1/2} h^{-s/2} \right\}. \tag{3.20}$$

Hence, by Lemma 2.3, we have

$$f\left(\frac{1}{2}\right)^2 \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{-a/2}^2 + \alpha \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{s/2}^2 \leq (\omega(h, \alpha, s) ch^s \|\tilde{x}_\alpha\|_s)^2. \tag{3.21}$$

In particular,

$$\begin{aligned} f\left(\frac{1}{2}\right) \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{-a/2} &\leq \omega(h, \alpha, s) ch^s \|\tilde{x}_\alpha\|_s, \\ \alpha^{1/2} \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|_{s/2} &\leq \omega(h, \alpha, s) ch^s \|\tilde{x}_\alpha\|_s. \end{aligned} \tag{3.22}$$

From these, we obtain (3.14) and (3.15).

Now, to prove (3.16), observe from (2.23) and (3.11) that

$$\tilde{x}_\alpha = L^{-s/2} (A_s + \alpha I)^{-1} L^{-s/2} \tilde{y}. \tag{3.23}$$

By Proposition 2.1, taking $\nu = -s/(s+a)$, we have

$$\begin{aligned} &\|L^{s/2} (A_s + \alpha I)^{-1} L^{-s/2} (\tilde{y} - \gamma)\| \\ &\leq \frac{1}{f(-s/(2s+2a))} \|A_s^{-s/(2s+2a)} (A_s + \alpha I)^{-1} L^{-s/2} (\tilde{y} - \gamma)\| \\ &\leq \frac{\|(A_s + \alpha I)^{-1}\|}{f(-s/(2s+2a))} \|A_s^{-s/(2s+2a)} L^{-s/2} (\tilde{y} - \gamma)\| \\ &\leq \frac{\alpha^{-1} g(-s/(2s+2a))}{f(-s/(2s+2a))} \|L^{-s/2} (\tilde{y} - \gamma)\|_{s/2} \end{aligned} \tag{3.24}$$

so that

$$\|L^{s/2} (A_s + \alpha I)^{-1} L^{-s/2} (\tilde{y} - \gamma)\| \leq \mathfrak{F}(s, a) \alpha^{-1} \delta. \tag{3.25}$$

Since $L^{-s/2} \gamma = A_s L^{s/2} \hat{x}$, we have

$$\begin{aligned} &\|L^{s/2} (A_s + \alpha I)^{-1} L^{-s/2} \gamma\| \\ &\leq \frac{1}{f(-s/(2s+2a))} \|A_s^{-s/(2s+2a)} (A_s + \alpha I)^{-1} A_s L^{s/2} \hat{x}\| \\ &\leq \frac{1}{f(-s/(2s+2a))} \|(A_s + \alpha I)^{-1} A_s^{(a+t)/(a+s)}\| \|A_s^{(s-2t)/(2a+2s)} L^{s/2} \hat{x}\|, \end{aligned} \tag{3.26}$$

where

$$\|A_s^{(s-2t)/(2a+2s)} L^{s/2} \hat{x}\| \leq g\left(\frac{s-2t}{2s+2a}\right) \|\hat{x}\|_t. \quad (3.27)$$

Since

$$\|(A_s + \alpha I)^{-1} A_s^\tau\| \leq \alpha^{\tau-1}, \quad 0 < \tau \leq 1, \quad (3.28)$$

it follows from the above relations that

$$\begin{aligned} \|L^{s/2} (A_s + \alpha I)^{-1} L^{-s/2} \mathcal{Y}\| &\leq \frac{g((s-2t)/(2s+2a)) \|\hat{x}\|_t}{f(-s/(2s+2a))} \alpha^{(t-s)/(s+a)} \\ &= \mathcal{G}(s, t, a) \alpha^{(t-s)/(s+a)}. \end{aligned} \quad (3.29)$$

Thus, (3.25) and (3.29) give

$$\begin{aligned} \|\tilde{x}_\alpha\|_s &\leq \|L^{s/2} (A_s + \alpha I)^{-1} L^{-s/2} \tilde{\mathcal{Y}}\| \\ &\leq \|L^{s/2} (A_s + \alpha I)^{-1} L^{-s/2} (\tilde{\mathcal{Y}} - \mathcal{Y})\| + \|L^{s/2} (A_s + \alpha I)^{-1} L^{-s/2} \mathcal{Y}\| \\ &\leq \mathcal{F}(s, a) \alpha^{-1} \delta + \mathcal{G}(s, t, a) \alpha^{(t-s)/(s+a)}. \end{aligned} \quad (3.30)$$

Now, the estimates (3.14) and (3.15) together with the interpolation inequality (3.2) give

$$\begin{aligned} \|\tilde{x}_\alpha - \tilde{x}_{\alpha, h}\| &\leq \|\tilde{x}_\alpha - \tilde{x}_{\alpha, h}\|_{-a/2}^{s/(s+a)} \|\tilde{x}_\alpha - \tilde{x}_{\alpha, h}\|_s^{a/(s+a)} \\ &\leq f\left(\frac{1}{2}\right)^{-s/(s+a)} \alpha^{-a/2(s+a)} \Phi(s, h, \alpha) h^s \|\tilde{x}_\alpha\|_s. \end{aligned} \quad (3.31)$$

From this, the result follows by making use of the estimate (3.16) for \tilde{x}_α . \square

4. A priori error estimates. Now we choose the regularization parameter α and discretization parameter h a priori depending on the noise level δ such that optimal order $O(\delta^{t/(t+a)})$ yields whenever $\hat{x} \in H_t$.

THEOREM 4.1. *Suppose $\hat{x} \in H_t$ with $0 < t \leq s + a$ and assumption (2.32) holds. Suppose, in addition, that*

$$\alpha = c_0 \delta^{(s+a)/(t+a)}, \quad h = d_0 \delta^{1/(t+a)} \quad (4.1)$$

for some constants $c_0, d_0 > 0$. Then, using the notations in Theorems 2.2 and 3.2,

$$\|\hat{x} - \tilde{x}_{\alpha, h}\| \leq [\eta(s, t) + \xi(s, t)] \delta^{t/(t+a)}, \quad (4.2)$$

where

$$\begin{aligned} \eta(s, t) &= \max \left\{ \phi(s, t) \|x\|_t c_0^{t/(t+a)}, \psi(s) c_0^{-a/(s+a)} \right\}, \\ \xi(s, t) &= c \left[f\left(\frac{1}{2}\right) \right]^{-s/(s+a)} d_0^s \left(c_0^{-1} + c_0^{(t-s)/(t+a)} \right) \\ &\quad \times \max \left\{ \mathcal{F}(s, a), \mathcal{G}(s, t, a) \right\} \max \left\{ g\left(\frac{1}{2}\right) d_0^{a/2}, c_0^{1/2} d_0^{-s/2} \right\}. \end{aligned} \tag{4.3}$$

PROOF. Using the choice (4.1), it is seen that

$$\begin{aligned} \Phi(s, h, \alpha) \alpha^{-a/2(s+a)} &= c c_0^{-a/2(s+a)} \max \left\{ g\left(\frac{1}{2}\right) d_0^{a/2}, c_0^{1/2} d_0^{-s/2} \right\}, \\ \delta \alpha^{-1} h^s &= c_0^{-1} d_0^s \delta^{t/(t+a)}, \\ \alpha^{(t-s)/(s+a)} h^s &= c_0^{(t-s)/(t+a)} d_0^s \delta^{t/(t+a)}. \end{aligned} \tag{4.4}$$

Therefore, by Theorem 3.2, we have

$$\|\tilde{x}_\alpha - \tilde{x}_{\alpha, h}\| \leq \xi(s, t) \delta^{t/(t+a)}. \tag{4.5}$$

Also, from Theorem 2.2, we have

$$\|\hat{x} - \tilde{x}_\alpha\| \leq \eta(s, t) \delta^{t/(t+a)}. \tag{4.6}$$

Thus the result follows from the inequality

$$\|\hat{x} - \tilde{x}_\alpha\| \leq \|\hat{x} - \tilde{x}_\alpha\| + \|\tilde{x}_\alpha - \tilde{x}_{\alpha, h}\|. \tag{4.7}$$

□

REMARK 4.2. We observe that the error bound obtained is of the same order as of Theorem 2.2, and this order is optimal with respect to the source set

$$M_{\rho, t} = \{x \in H_t : \|x\|_t \leq \rho\} \tag{4.8}$$

in the sense of the best possible worst error (cf. [4]).

5. Discrepancy principle. In this section, we consider a discrepancy principle to choose the regularization parameter α depending on the noise level δ and the discretization parameter h . This is a finite-dimensional variant of the discrepancy principle considered in [5].

We assume throughout that $y \neq 0$. Suppose that $\tilde{y} \in H$ is such that

$$\|y - \tilde{y}\| \leq \delta \quad (5.1)$$

for a known error level $\delta > 0$ and $P_h \tilde{y} \neq 0$, where P_h is the orthogonal projection of H onto S_h . We assume, throughout this section, that

$$\|A(P_h - I)\| \leq c_3 h, \quad h > 0, \quad (5.2)$$

for some $c_3 > 0$, independent of h . Let

$$R_\alpha := (A_s + \alpha I)^{-1}. \quad (5.3)$$

We will make use of the relation

$$\|R_\alpha A_s^\tau\| \leq \alpha^{\tau-1}, \quad \alpha > 0, \quad 0 < \tau \leq 1, \quad (5.4)$$

which follows from the spectral properties of the selfadjoint operator A_s , $s > 0$.

Let s, a be fixed positive real numbers. For $\alpha > 0$ and $x \in H$, consider the functions

$$F(\alpha, x) = \frac{\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|}. \quad (5.5)$$

Note that, by assumption (2.18), $\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|$ is nonzero for every $x \in H$ with $P_h x \neq 0$, so that the function $F(\alpha, x)$ is well defined for all such x . We observe that the assumption $P_h x \neq 0$ is satisfied for $x \neq 0$ and h small enough, if $P_h x \rightarrow x$ as $h \rightarrow 0$ for every $x \in H$.

In the following, we assume that h is such that $P_h \tilde{y} \neq 0$.

In order to choose the regularization parameter α , we consider the discrepancy principle

$$F(\alpha, \tilde{y}) = b\delta + dh \quad (5.6)$$

for some $b, d > 0$. In the due course, we will make use of the relation

$$f\left(\frac{-s}{2s+2a}\right) \|x\| \leq \|A_s^{-s/(2s+2a)} L^{-s/2} x\| \leq g\left(\frac{-s}{2s+2a}\right) \|x\| \quad (5.7)$$

which can easily be derived from [Proposition 2.1](#).

First we prove the monotonicity of the function $F(\alpha, x)$ defined in (5.5).

THEOREM 5.1. *Let $x \in H$ be such that the function $\alpha \mapsto F(\alpha, x)$ for $\alpha > 0$ in (5.5) is well defined. Then, $F(\cdot, x)$ is increasing and it is continuously differentiable with $F'(\alpha, x) \geq 0$ for all $\alpha > 0$. In addition,*

$$\lim_{\alpha \rightarrow 0} F(\alpha, x) = 0, \quad \lim_{\alpha \rightarrow \infty} F(\alpha, x) = \|A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|. \tag{5.8}$$

PROOF. Using the definition (5.5) of $F(\alpha, \cdot)$, we have

$$\begin{aligned} & \frac{\partial}{\partial \alpha} F(\alpha, x) \\ &= \frac{(\partial/\partial \alpha)(F^2(\alpha, x))}{2F(\alpha, x)} \\ &= \frac{2\alpha \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2 \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2}{2\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2} \\ & \quad \times \frac{(\partial/\partial \alpha) [\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2]}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^3} \\ & \quad - \frac{\alpha^2 \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^4 (\partial/\partial \alpha) [\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2]}{2\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2 \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^3} \\ &= \frac{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2 (\partial/\partial \alpha) [\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2]}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^3} \\ & \quad - \frac{\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2 (\partial/\partial \alpha) [\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2]}{2 \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^3}. \end{aligned} \tag{5.9}$$

Let $\{E_\lambda : 0 \leq \lambda \leq a\}$ be the spectral family of A_s , where $a \geq \|A_s\|$. Then

$$\begin{aligned} & \frac{\partial}{\partial \alpha} (\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2) \\ &= \frac{\partial}{\partial \alpha} \int_0^a \frac{\alpha}{\lambda^{s/(s+a)} (\lambda + \alpha)^3} d\langle E_\lambda L^{-s/2} P_h x, L^{-s/2} P_h x \rangle \\ &= \int_0^a \left[\frac{1}{\lambda^{s/(s+a)} (\lambda + \alpha)^3} - \frac{3\alpha}{\lambda^{s/(s+a)} (\lambda + \alpha)^4} \right] d\langle E_\lambda L^{-s/2} P_h x, L^{-s/2} P_h x \rangle \\ &= \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2 - 3\alpha \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2. \end{aligned} \tag{5.10}$$

Similarly, we obtain

$$\frac{\partial}{\partial \alpha} (\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|) = -4 \|R_\alpha^{5/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h x\|^2. \tag{5.11}$$

Therefore, from (5.9), by using (5.10) and (5.11), we get

$$\begin{aligned} \frac{\partial}{\partial \alpha} F(\alpha, \mathbf{x}) &= \frac{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^3} \\ &\times \left[\|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 - 3\alpha \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \right] \\ &+ \frac{2\alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \|R_\alpha^{5/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^3}. \end{aligned} \quad (5.12)$$

The above equation can be rewritten as

$$\begin{aligned} \frac{\partial}{\partial \alpha} F(\alpha, \mathbf{x}) &= \frac{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^3} \\ &\times \left[\|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 - \alpha \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \right] \\ &+ \frac{2\alpha}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^3} \\ &\times \left[\|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \|R_\alpha^{5/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \right. \\ &\left. - \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^4 \right]. \end{aligned} \quad (5.13)$$

Since

$$\begin{aligned} &\|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \\ &= \langle (A_s + \alpha I)^{-3} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle, \\ &\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \\ &= \langle (A_s + \alpha I)^{-3} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, (A_s + \alpha I)^{-1} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle, \end{aligned} \quad (5.14)$$

we see that

$$\begin{aligned} &\|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \\ &= \alpha \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \\ &+ \langle (A_s + \alpha I)^{-3} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, A_s (A_s + \alpha I)^{-1} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle \\ &= \alpha \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 + \|A_s^{a/(2s+2a)} R_\alpha^2 L^{-s/2} P_h \mathbf{x}\|^2. \end{aligned} \quad (5.15)$$

Also, we have

$$\begin{aligned}
 & \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^4 \\
 &= [\langle R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle]^2 \\
 &= [\langle R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, R_\alpha^{5/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle]^2 \\
 &\leq \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 \|R_\alpha^{5/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2.
 \end{aligned} \tag{5.16}$$

Hence,

$$\frac{\partial}{\partial \alpha} (F(\alpha, \mathbf{x})) \geq 0. \tag{5.17}$$

To prove the last part of the theorem, we observe that

$$\begin{aligned}
 & \alpha^2 \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\| - F(\alpha, \mathbf{x}) \\
 &= \frac{\alpha^2 \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 - \alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|}.
 \end{aligned} \tag{5.18}$$

We note that

$$\begin{aligned}
 \alpha^2 \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 &= \alpha \langle R_\alpha^3 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, \alpha R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle, \\
 \alpha \|R_\alpha^{3/2} A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|^2 &= \alpha \langle R_\alpha^3 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle.
 \end{aligned} \tag{5.19}$$

Since

$$\alpha R_\alpha - I = A_s R_\alpha = R_\alpha A_s, \tag{5.20}$$

it follows that

$$\begin{aligned}
 & \alpha^2 \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\| - F(\alpha, \mathbf{x}) \\
 &= \frac{-\alpha \langle R_\alpha^3 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, A_s R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|} \\
 &= \frac{-\alpha \|A_s^{a/(2s+2a)} R_\alpha^2 L^{-s/2} P_h \mathbf{x}\|^2}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|} \\
 &\leq 0
 \end{aligned} \tag{5.21}$$

so that

$$F(\alpha, \mathbf{x}) \geq \alpha^2 \|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\| \geq \alpha^2 \frac{\|A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|}{(\|A_s\| + \alpha)^2}. \tag{5.22}$$

Also, we have

$$\begin{aligned} F(\alpha, \mathbf{x}) &= \frac{\alpha \langle R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}, R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x} \rangle}{\|R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|} \\ &\leq \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|. \end{aligned} \quad (5.23)$$

Hence

$$\left(\frac{\alpha}{\|A_s\| + \alpha} \right)^2 \|A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\| \leq F(\alpha, \mathbf{x}) \leq \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|. \quad (5.24)$$

From this we can conclude that

$$\lim_{\alpha \rightarrow 0} F(\alpha, \mathbf{x}) = 0, \quad \lim_{\alpha \rightarrow \infty} F(\alpha, \mathbf{x}) = \|A_s^{-s/(2s+2a)} L^{-s/2} P_h \mathbf{x}\|. \quad (5.25)$$

This completes the proof. \square

For the next theorem, in addition to (5.1), we assume that the inexact data $\tilde{\mathbf{y}}$ satisfies the relation

$$\|A_s^{-s/(2s+2a)} L^{-s/2} P_h \tilde{\mathbf{y}}\| \geq b\delta + dh. \quad (5.26)$$

This assumption is satisfied for small enough h and δ , if, for example,

$$(b + \tilde{f}(s))\delta + (d + c_3 \tilde{f}(s) \|\hat{\mathbf{x}}\|)h \leq \tilde{f}(s) \|\mathbf{y}\|, \quad (5.27)$$

where $\tilde{f}(s) = f(-s/(2s+2a))$, because

$$\|P_h \tilde{\mathbf{y}}\| \geq \|\mathbf{y}\| - \|(I - P_h)A\hat{\mathbf{x}}\| - \delta, \quad (5.28)$$

and by (5.7),

$$\|A_s^{-s/(2s+2a)} L^{-s/2} P_h \tilde{\mathbf{y}}\| \geq \tilde{f}(s) \|P_h \tilde{\mathbf{y}}\|. \quad (5.29)$$

Now the following theorem is a consequence of [Theorem 5.1](#).

THEOREM 5.2. *Assume that (5.1) and (5.26) are satisfied. Then there exists a unique $\alpha := \alpha(\delta, h)$ satisfying*

$$F(\alpha, \tilde{\mathbf{y}}) = b\delta + dh. \quad (5.30)$$

In order to obtain an estimate for the error $\|\hat{\mathbf{x}} - \tilde{\mathbf{x}}_{\alpha, h}\|$ with the parameter choice strategy (5.30), we will make use of (3.31). The next lemma gives an error estimate for $\|\tilde{\mathbf{x}}_\alpha\|_s$ in terms of $\alpha = \alpha(\delta, h)$, δ , and h .

LEMMA 5.3. *Let $\alpha := \alpha(\delta, h)$ be the unique solution of (5.30). Then for any fixed $\tau > 0$,*

$$\|\tilde{\mathbf{x}}_\alpha\|_s \leq c_4 (\delta + h)^{\tau/(\tau+1)} \alpha^{-1}, \quad (5.31)$$

where

$$c_4 \geq \max \{b + \tilde{g}(s), c + c_3 \|\hat{x}\| \tilde{g}(s)\} \tilde{g}(s) \|\tilde{y}\| \quad (5.32)$$

with $\tilde{g}(s) := g(-s/(2s+2a))$.

PROOF. By (3.30), we have

$$\|\tilde{x}_\alpha\|_s \leq \|L^{s/2} R_\alpha L^{-s/2} \tilde{y}\| \leq \tilde{f}^{-1}(s) \alpha^{-1} \|\alpha R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}\|. \quad (5.33)$$

To obtain an estimate for $\|\alpha R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}\|$, we will make use of the moment inequality (2.17). Precisely, we use (2.17) with

$$u = \tau, \quad v = 1 + \tau, \quad B = \alpha R_\alpha, \quad x = \alpha^{1-\tau} R_\alpha^{1-\tau} A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}. \quad (5.34)$$

Then, since

$$\|x\| \leq \|A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}\| \leq g\left(\frac{-s}{2s+2a}\right) \|\tilde{y}\|, \quad (5.35)$$

we have

$$\begin{aligned} & \|\alpha R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}\| \\ &= \|B^\tau x\| \leq \|B^{\tau+1} x\|^{\tau/(\tau+1)} \|x\|^{1/(\tau+1)} \\ &= \|\alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}\|^{\tau/(\tau+1)} \left[g\left(\frac{-s}{2s+2a}\right) \|\tilde{y}\| \right]^{1/(\tau+1)}. \end{aligned} \quad (5.36)$$

Further, by (5.21),

$$\begin{aligned} & \|\alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}\| \\ & \leq \|\alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} (I - P_h) \tilde{y}\| + \|\alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \tilde{y}\| \\ & \leq \tilde{g}(s) \|(I - P_h) \tilde{y}\| + F(\alpha, \tilde{y}) \\ & \leq \tilde{g}(s) [\|(I - P_h)(\tilde{y} - y)\| + \|(I - P_h)A\hat{x}\|] + F(\alpha, \tilde{y}) \\ & \leq \tilde{g}(s) [\delta + c_3 \|\hat{x}\| h] + F(\alpha, \tilde{y}). \end{aligned} \quad (5.37)$$

Therefore, if $\alpha := \alpha(\delta, h)$ is the unique solution of (5.30), then we have

$$\|\alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} \tilde{y}\| \leq (b + \tilde{g}(s))\delta + (d + \tilde{g}(s)c_3 \|\hat{x}\|)h. \quad (5.38)$$

Now the result follows from (5.33), (5.36), (5.37), and (5.38). \square

LEMMA 5.4. *Suppose that \hat{x} belongs to H_t for some $t \leq s$, and $\alpha := \alpha(\delta, h) > 0$ is the unique solution of (5.30), where $b > \tilde{g}(s)$ and $d > c_3 \|\hat{x}\| \tilde{g}(s)$ with $\tilde{g}(s) := g(-s/(2s+2a))$. Then*

$$\alpha \geq c_0 \delta^{(s+a)/(t+a)}, \quad c_0 = \frac{\min \{b - \tilde{g}(s), d - c_3 \|\hat{x}\| \tilde{g}(s)\}}{g((s-2t)/(2s+2a))\rho}. \quad (5.39)$$

PROOF. Note that by (5.23), Proposition 2.1, and (2.18), we have

$$\begin{aligned}
 F(\alpha, \tilde{y}) &\leq \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h \tilde{y}\| \\
 &\leq \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h (\tilde{y} - y)\| \\
 &\quad + \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} (P_h - I) y\| + \alpha \|R_\alpha A_s^{-s/(2s+2a)} A_s L^{s/2} \hat{x}\| \\
 &\leq \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h (\tilde{y} - y)\| \\
 &\quad + \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} (P_h - I) y\| + \alpha \|R_\alpha A_s^{(s+2a)/(2s+2a)} L^{s/2} \hat{x}\| \\
 &\leq \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} P_h (\tilde{y} - y)\| \\
 &\quad + \alpha \|R_\alpha A_s^{-s/(2s+2a)} L^{-s/2} (P_h - I) y\| + \alpha \|R_\alpha A_s^{(t+a)/(s+a)} A_s^{(s-2t)/(2s+2a)} L^{s/2} \hat{x}\| \\
 &\leq \tilde{g}(s) [\delta + c_3 \|\hat{x}\| h] + \|\alpha R_\alpha A_s^{(t+a)/(s+a)}\| \|A_s^{(s-2t)/(2s+2a)} L^{s/2} \hat{x}\| \\
 &\leq \tilde{g}(s) [\delta + c_3 \|\hat{x}\| h] + g\left(\frac{s-2t}{2s+2a}\right) \rho \alpha^{(t+a)/(s+a)}.
 \end{aligned} \tag{5.40}$$

Thus

$$\min\{b - \tilde{g}(s), d - c_3 \|\hat{x}\| \tilde{g}(s)\} (\delta + h) \leq g\left(\frac{s-2t}{2s+2a}\right) \rho \alpha^{(t+a)/(s+a)}, \tag{5.41}$$

which implies

$$\alpha \geq c_0 (\delta + h)^{(s+a)/(t+a)}, \quad c_0 = \frac{\min\{b - \tilde{g}(s), d - c_3 \|\hat{x}\| \tilde{g}(s)\}}{g((s-2t)/(2s+2a)) \rho}. \tag{5.42}$$

This completes the proof. \square

THEOREM 5.5. Under the assumptions in Lemma 5.4, for any fixed $\tau > 0$,

$$\|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \leq c_5 (\delta + h)^\zeta, \tag{5.43}$$

where

$$\zeta := \frac{\tau}{\tau+1} + \frac{s}{2} - \frac{s+2a}{2t+2a} + \gamma, \quad c_5 \geq cc_4 f\left(\frac{1}{2}\right)^{-s/(s+a)} \max\left\{g\left(\frac{1}{2}\right), 1\right\} \tag{5.44}$$

with

$$\gamma := \begin{cases} 0, & \text{if } t \geq 1-a, \\ \frac{s+a}{2} \left(1 - \frac{1}{t+a}\right), & \text{if } t < 1-a. \end{cases} \tag{5.45}$$

PROOF. Note that by (3.31) and Lemmas 5.3 and 5.4,

$$\begin{aligned}
 & \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\
 & \leq f\left(\frac{1}{2}\right)^{-s/(s+a)} \Phi(s, h, \alpha) \alpha^{-a/(2s+2a)} h^s \|\tilde{x}_\alpha\|_s \\
 & \leq c f\left(\frac{1}{2}\right)^{-s/(s+a)} \max\left\{g\left(\frac{1}{2}\right) h^{a/2}, \alpha^{1/2} h^{-s/2}\right\} c_4 \alpha^{-a/(2s+2a)} h^s \alpha^{-1} (\delta + h)^{\tau/(\tau+1)} \\
 & \leq c f\left(\frac{1}{2}\right)^{-s/(s+a)} \max\left\{g\left(\frac{1}{2}\right) h^{(s+a)/2} \alpha^{-1/2}, 1\right\} c_4 \alpha^{-a/(2s+2a)-1/2} h^{s/2} (\delta + h)^{\tau/(\tau+1)} \\
 & \leq c f\left(\frac{1}{2}\right)^{-s/(s+a)} \max\left\{g\left(\frac{1}{2}\right) (\delta + h)^{(s+a)/2-(s+a)/(2t+2a)}, 1\right\} \\
 & \quad \times c_4 (\delta + h)^{\tau/(\tau+1)+s/2-a/(2t+2a)-(s+a)/(2t+2a)} \\
 & \leq c f\left(\frac{1}{2}\right)^{-s/(s+a)} \max\left\{g\left(\frac{1}{2}\right), 1\right\} c_4 (\delta + h)^{\tau/(\tau+1)+s/2-a/(2t+2a)-(s+a)/(2t+2a)+y}.
 \end{aligned}
 \tag{5.46}$$

This completes the proof. □

THEOREM 5.6. Under the assumptions in Lemma 5.4,

$$\|\hat{x} - x_\alpha\| = O((\delta + h)^{t/(t+a)}).
 \tag{5.47}$$

PROOF. Since x_α is the solution of (2.22), we have

$$\begin{aligned}
 \hat{x} - x_\alpha &= \hat{x} - (A + \alpha L^s)^{-1} \gamma \\
 &= \alpha L^{-s/2} (A_s + \alpha I)^{-1} L^{s/2} \hat{x} \\
 &= \alpha L^{-s/2} R_\alpha L^{s/2} \hat{x}.
 \end{aligned}
 \tag{5.48}$$

Therefore, by (5.7), we have

$$f\left(\frac{s}{2s+2a}\right) \|\hat{x} - x_\alpha\| \leq \|\alpha A_s^{s/(2s+2a)} R_\alpha L^{s/2} \hat{x}\|.
 \tag{5.49}$$

To obtain an estimate for $\|\alpha A_s^{s/(2s+2a)} R_\alpha L^{s/2} \hat{x}\|$, first we will make use the moment inequality (2.17) with

$$u = \frac{t}{a}, \quad v = 1 + \frac{t}{a}, \quad B = \alpha R_\alpha A_s^{a/(s+a)}, \quad x = \alpha^{1-t/a} R_\alpha^{1-t/a} A_s^{(s-2t)/(2s+2a)} L^{s/2} \hat{x}.
 \tag{5.50}$$

Then, since

$$\|x\| \leq \|A_s^{(s-2t)/(2s+2a)} L^{s/2} \hat{x}\| \leq g \left(\frac{s-2t}{2s+2a} \right) \|L^{s/2} \hat{x}\|_{t-s/2} \leq g \left(\frac{s-2t}{2s+2a} \right) \rho, \quad (5.51)$$

we have

$$\begin{aligned} & \| \alpha A_s^{s/(2s+2a)} R_\alpha L^{s/2} \hat{x} \| \\ &= \| B^{t/a} x \| \\ &\leq \| B^{1+t/a} x \|^{t/(t+a)} \| x \|^{a/(t+a)} \\ &\leq \| \alpha^2 R_\alpha^2 A_s^{(2a+s)/(2s+2a)} L^{s/2} \hat{x} \|^{t/(t+a)} \| x \|^{a/(t+a)} \\ &\leq \| \alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} y \|^{t/(t+a)} \| x \|^{a/(t+a)} \\ &\leq g \left(\frac{s-2t}{2s+2a} \right)^{a/(t+a)} \rho^{a/(t+a)} \| \alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} y \|^{t/(t+a)}. \end{aligned} \quad (5.52)$$

Further, by (5.2), (5.7), and (5.21),

$$\begin{aligned} & \| \alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} y \| \\ &\leq \| \alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} (y - \tilde{y}) \| + \| \alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} (I - P_h) \tilde{y} \| \\ &\quad + \| \alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} P_h \tilde{y} \| \\ &\leq g \left(-\frac{s}{2s+2a} \right) (\delta + c_3 \|\hat{x}\| h) + F(\alpha, \tilde{y}). \end{aligned} \quad (5.53)$$

Therefore, if $\alpha := \alpha(\delta, h)$ is the unique solution of (5.30), then we have

$$\| \alpha^2 R_\alpha^2 A_s^{-s/(2s+2a)} L^{-s/2} y \| \leq [\tilde{g}(s) + b] \delta + [\tilde{g}(s) c_3 \|\hat{x}\| + d] h. \quad (5.54)$$

Now the result follows from (5.49), (5.52), (5.53), and (5.54). \square

THEOREM 5.7. *Under the assumptions in Lemma 5.4, for any fixed $\tau > 0$,*

$$\|\hat{x} - \tilde{x}_{\alpha, h}\| \leq c_6 (\delta + h)^\mu, \quad \mu := \min \left\{ \frac{t}{t+a}, \zeta \right\} \quad (5.55)$$

for some $c_6 > 0$, and ζ as in Theorem 5.5.

PROOF. Let x_α and \tilde{x}_α be the solutions of (2.22) and (2.23), respectively. Then by triangle inequality, (5.4), and Proposition 2.1,

$$\begin{aligned}
 & \|\hat{x} - \tilde{x}_{\alpha,h}\| \\
 & \leq \|\hat{x} - x_\alpha\| + \|x_\alpha - \tilde{x}_\alpha\| + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\
 & = \|\hat{x} - x_\alpha\| + \|L^{-s/2}R_\alpha L^{-s/2}(\mathcal{Y} - \tilde{\mathcal{Y}})\| + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\
 & \leq \|\hat{x} - x_\alpha\| + \frac{1}{f(s/(2s+2a))} \|A_s^{s/(2s+2a)}R_\alpha L^{-s/2}(\mathcal{Y} - \tilde{\mathcal{Y}})\| + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\
 & \leq \|\hat{x} - x_\alpha\| + \frac{1}{f(s/(2s+2a))} \|A_s^{s/(s+a)}R_\alpha A_s^{-s/(2s+2a)}L^{-s/2}(\mathcal{Y} - \tilde{\mathcal{Y}})\| + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\
 & \leq \|\hat{x} - x_\alpha\| + \frac{1}{f(s/(2s+2a))} \|A_s^{s/(s+a)}R_\alpha\| \|A_s^{-s/(2s+2a)}L^{-s/2}(\mathcal{Y} - \tilde{\mathcal{Y}})\| + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\
 & \leq \|\hat{x} - x_\alpha\| + \frac{g(-s/(2s+2a))}{f(s/(2s+2a))} \delta \alpha^{-a/(s+a)} + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\| \\
 & \leq \|\hat{x} - x_\alpha\| + \frac{g(-s/(2s+2a))}{f(s/(2s+2a))} (\delta + h) \alpha^{-a/(s+a)} + \|\tilde{x}_\alpha - \tilde{x}_{\alpha,h}\|.
 \end{aligned}
 \tag{5.56}$$

The proof now follows from Lemma 5.4 and Theorems 5.5 and 5.6. □

COROLLARY 5.8. *If t, s, a satisfy $\max\{0, 1 - a\} < t \leq s$ and τ is large enough such that*

$$\gamma + \frac{s}{2} \left[1 - \frac{1}{t+a} \right] \geq \frac{1}{\tau+1},
 \tag{5.57}$$

then

$$\|\hat{x} - \tilde{x}_{\alpha,h}\| \leq c_6 (\delta + h)^{t/(t+a)}
 \tag{5.58}$$

with c_6 as in Theorem 5.7.

PROOF. Let ζ, μ be as in Theorems 5.5 and 5.7, respectively. Then we observe that

$$\mu = \frac{t}{t+a} \quad \text{if } \gamma + \frac{s}{2} \left[1 - \frac{1}{t+a} \right] \geq \frac{1}{\tau+1}.
 \tag{5.59}$$

Hence the result follows from Theorem 5.7. □

6. Order optimality of the error estimates. In order to measure the quality of an algorithm to solve an equation of the form (2.1), Micchelli and Rivlin [12] considered the quantity

$$e(M, \delta) := \sup \{ \|x\| : x \in M, \|Ax\| \leq \delta \}
 \tag{6.1}$$

and showed that

$$e(M, \delta) \leq E(M, \delta) \leq 2e(M, \delta),
 \tag{6.2}$$

where

$$E(M, \delta) = \inf_R \sup \{ \|x - Rv\| : x \in M, v \in H, \|Ax - v\| \leq \delta \} \quad (6.3)$$

is the best possible worst error. Here the stabilizing set M is assumed to be convex such that $M = -M$ with $0 \in M$ (see also, Väinikko and Veretennikov [22]), and infimum is taken over all algorithms $R : Y \rightarrow X$. Since H is a Hilbert space and A is assumed to be selfadjoint and positive, we, in fact, have (cf. Melkman and Micchelli [11])

$$e(M, \delta) = E(M, \delta). \quad (6.4)$$

Now using the assumption (2.18), and taking $r = -a$, $\lambda = 0$ in the interpolation inequality (2.15), we obtain

$$\|x\| \leq \|x\|_{-a}^{t/(t+a)} \|x\|_t^{a/(t+a)} \leq \left(\frac{\|Ax\|}{c_1} \right)^{t/(t+a)} \|x\|_t^{a/(t+a)}, \quad x \in H_t. \quad (6.5)$$

Therefore, for the set

$$M_{t,\rho} = \{x : \|x\|_t \leq \rho\} \quad (6.6)$$

with a fixed $t > 0$, $\rho > 0$,

$$e(M_{t,\rho}, \delta) \leq \left(\frac{\delta}{c_1} \right)^{t/(t+a)} \rho^{a/(t+a)}. \quad (6.7)$$

It is known that the above estimate for $e(M_{t,\rho}, \delta)$ is sharp (cf. Väinikko [21]). In view of the above observations, an algorithm is called an *optimal order yielding algorithm* with respect to $M_{t,\rho}$ and the assumption (2.18), if it yields an approximation \hat{x} corresponding to the data \hat{y} with $\|y - \hat{y}\| \leq \delta$ satisfying

$$\|\hat{x} - \tilde{x}\| = O(\delta^{t/(t+a)}), \quad x \in H_t. \quad (6.8)$$

Clearly, Corollary 5.8 shows that if $h = O(\delta)$ and if $\max\{0, 1 - a\} < t \leq s$ and τ is large enough such that

$$y + \frac{s}{2} \left[1 - \frac{1}{t+a} \right] \geq \frac{1}{\tau+1}, \quad (6.9)$$

then we obtain the optimal order.

7. Applications. For $r \geq 2$, denote by S_h the space of r th-order splines on the uniform mesh of width $h = 1/n$, that is, S_h consists of functions in $C^{r-1}[0, 1]$ which are piecewise polynomials of degree $r - 2$. For positive integers s , let H^s denote the Sobolev space of functions $u \in C^{s-1}[0, 1]$ with u^{s-1} absolutely continuous and the norm $\|u\|_{H^s}$ defined by

$$\|u\|_{H^s} = \left(\sum_{i=1}^s \|u^{(i)}\| \right)^{1/2}, \quad u \in H^s. \quad (7.1)$$

Then S_h is a finite-dimensional subspace of H^{r-1} which has the following well-known approximation property (cf. [1]): for $u \in H^s$, $s \in \mathbb{N}$, there is a constant κ (independent of h) such that

$$\inf_{\varphi \in S_h} \|u - \varphi\|_{H^j} \leq \kappa h^{\min\{s,r\}-j} \|u\|_{H^s}, \quad u \in H^s, \quad j \in \{0, 1\}, \tag{7.2}$$

so that assumption (2.32) is satisfied. We take L as in (2.19), that is,

$$Lx := \sum_{j=1}^{\infty} j^2 \langle x, u_j \rangle u_j, \tag{7.3}$$

where $u_j(t) := \sqrt{2} \sin(j\pi t)$, $j \in \mathbb{N}$, with domain of L as

$$D(L) := \left\{ x \in L^2[0, 1] : \sum_{j=1}^{\infty} j^4 |\langle x, u_j \rangle|^2 < \infty \right\}. \tag{7.4}$$

In this case, $(H_t)_{t \in \mathbb{R}}$ is given as in (2.21). It can be seen that

$$\begin{aligned} H_t &= \left\{ x \in L^2[0, 1] : \sum_{j=1}^{\infty} j^{4t} |\langle x, u_j \rangle|^2 < \infty \right\} \\ &= \left\{ x \in H^{2t}(0, 1) : x^{(2l)}(0) = x^{(2l)}(1) = 0, \quad l = 0, 1, \dots, \left\lceil t - \frac{1}{4} \right\rceil \right\}, \end{aligned} \tag{7.5}$$

where $\lceil p \rceil$ denotes the greatest integer less than or equal to p . We observe that $H^0 = L^2[0, 1]$, and for $s \in \mathbb{N}$, $H_s \subset H^s$.

Now, let $A : L^2[0, 1] \rightarrow L^2[0, 1]$ be a positive selfadjoint operator. Then we have

$$\|A(I - P_h)\| = \|(I - P)A\| = \sup_{\|u\| \leq 1} \inf_{\varphi \in S_h} \|Au - \varphi\|. \tag{7.6}$$

Hence, by (7.2),

$$\|A(I - P_h)\| \leq \kappa h^{\min(s,r)} \sup_{\|u\| \leq 1} \|Au\|_{H^s}. \tag{7.7}$$

From the above inequality it is clear that if $Au \in H_s$ for every $u \in L^2[0, 1]$, and if $A : L^2[0, 1] \rightarrow H_s$ is a bounded operator, then there exists a constant \hat{c} such that

$$\|A(I - P_h)\| \leq \kappa \hat{c} h^{\min(s,r)} \tag{7.8}$$

so that (5.2) is satisfied.

Now, we consider the case of an integral operator, namely (2.4), having all its eigenvalues nonnegative, and $k(\xi, t) = k(t, \xi)$ for all $(\xi, t) \in [0, 1] \times [0, 1]$ is such that it is differentiable s times with respect to the variable ξ with its s th derivative lying in $L^2[0, 1]$. For example, the integral operator may be the one associated with the *backward heat equation* problem considered in Section 2.

Now,

$$\frac{d^i}{d\xi^i} (Au)(\xi) = \int_0^1 \frac{\partial^i k(\xi, t)}{\partial \xi^i} u(t) dt \tag{7.9}$$

so that

$$\|Au\|_{H^s} \leq \|k\|_{0,s} \|u\| \quad \text{with } \|k\|_{0,s} = \sum_{i=0}^s \int_0^1 \int_0^1 \left| \frac{\partial^i k(t, \xi)}{\partial \xi^i} \right|^2 dt d\xi. \quad (7.10)$$

Thus we get (7.8) with $\hat{c} = \|k\|_{0,s}$.

REFERENCES

- [1] I. Babuška and A. K. Aziz, *Survey lectures on the mathematical foundations of the finite element method*, The Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations (Proc. Sympos., Univ. Maryland, Baltimore, Md, 1972) (A. K. Aziz, ed.), Academic Press, New York, 1972, pp. 1-359.
- [2] H. W. Engl, M. Hanke, and A. Neubauer, *Regularization of Inverse Problems*, Mathematics and Its Applications, vol. 375, Kluwer Academic Publishers, Dordrecht, 1996.
- [3] H. W. Engl and A. Neubauer, *Convergence rates for Tikhonov regularization in finite-dimensional subspaces of Hilbert scales*, Proc. Amer. Math. Soc. **102** (1988), no. 3, 587-592.
- [4] S. George and M. T. Nair, *Error bounds and parameter choice strategies for simplified regularization in Hilbert scales*, Integral Equations Operator Theory **29** (1997), no. 2, 231-242.
- [5] ———, *An optimal order yielding discrepancy principle for simplified regularization of ill-posed problems in Hilbert scales*, Int. J. Math. Math. Sci. (2003), no. 39, 2487-2499.
- [6] C. W. Groetsch, *The Theory of Tikhonov Regularization for Fredholm Equations of the First Kind*, Research Notes in Mathematics, vol. 105, Pitman, Massachusetts, 1984.
- [7] M. Hegland, *An optimal order regularization method which does not use additional smoothness assumptions*, SIAM J. Numer. Anal. **29** (1992), no. 5, 1446-1461.
- [8] A. Kirsch, *An Introduction to the Mathematical Theory of Inverse Problems*, Applied Mathematical Sciences, vol. 120, Springer-Verlag, New York, 1996.
- [9] S. G. Krein and J. I. Petunin, *Scales of Banach spaces*, Russian Math. Surveys **21** (1966), 85-160.
- [10] B. A. Mair, *Tikhonov regularization for finitely and infinitely smoothing operators*, SIAM J. Math. Anal. **25** (1994), no. 1, 135-147.
- [11] A. A. Melkman and C. A. Micchelli, *Optimal estimation of linear operators in Hilbert spaces from inaccurate data*, SIAM J. Numer. Anal. **16** (1979), no. 1, 87-105.
- [12] C. A. Micchelli and T. J. Rivlin, *A survey of optimal recovery*, Optimal Estimation in Approximation Theory (C. A. Micchelli and T. J. Rivlin, eds.), Plenum Press, New York, 1977, pp. 1-54.
- [13] M. T. Nair, *On Morozov's method for Tikhonov regularization as an optimal order yielding algorithm*, Z. Anal. Anwendungen **18** (1999), no. 1, 37-46.
- [14] ———, *Functional Analysis: A First Course*, Prentice Hall of India Private Limited, New Delhi, 2002.
- [15] ———, *Optimal order results for a class of regularization methods using unbounded operators*, Integral Equations Operator Theory **44** (2002), no. 1, 79-92.
- [16] M. T. Nair, M. Hegland, and R. S. Anderssen, *The trade-off between regularity and stability in Tikhonov regularization*, Math. Comp. **66** (1997), no. 217, 193-206.
- [17] F. Natterer, *Error bounds for Tikhonov regularization in Hilbert scales*, Applicable Anal. **18** (1984), no. 1-2, 29-37.
- [18] A. Neubauer, *An a posteriori parameter choice for Tikhonov regularization in Hilbert scales leading to optimal convergence rates*, SIAM J. Numer. Anal. **25** (1988), no. 6, 1313-1326.
- [19] E. Schock, *Ritz-regularization versus least-square-regularization. Solution methods for integral equations of the first kind*, Z. Anal. Anwendungen **4** (1985), no. 3, 277-284.

- [20] T. Schröter and U. Tautenhahn, *Error estimates for Tikhonov regularization in Hilbert scales*, Numer. Funct. Anal. Optim. **15** (1994), no. 1-2, 155-168.
- [21] G. Vainikko, *On the optimality of methods for ill-posed problems*, Z. Anal. Anwendungen **6** (1987), no. 4, 351-362.
- [22] G. M. Vainikko and A. Y. Veretennikov, *Iteration Procedures in Ill-Posed Problems*, 1st ed., Nauka, Moscow, 1986.
- [23] H. F. Weinberger, *A First Course in Partial Differential Equations with Complex Variables and Transform Methods*, Blaisdell Publishing, New York, 1965.

Santhosh George: Department of Mathematics, Government College of Arts, Science and Commerce, Sanquelim, Goa 403505, India

E-mail address: santhoshsq1729@yahoo.co.in

M. Thamban Nair: Department of Mathematics, Indian Institute of Technology Madras, Chennai 600036, India

E-mail address: mtnair@iitm.ac.in

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be