

SEMICOMPATIBILITY AND FIXED POINT THEOREMS IN FUZZY METRIC SPACE USING IMPLICIT RELATION

BIJENDRA SINGH AND SHISHIR JAIN

Received 27 January 2005 and in revised form 29 May 2005

The concept of semicompatibility has been introduced in fuzzy metric space and it has been applied to prove results on existence of unique common fixed point of four self-maps satisfying an implicit relation. Recently, Popa (2002) has employed a similar but not the same implicit relation to obtain a fixed point theorem for d -complete topological spaces. All the results of this paper are new.

1. Introduction

Cho et al. [2] introduced the notion of semicompatible maps in a d -topological space. They define a pair of self-maps (S, T) to be semicompatible if conditions (i) $Sy = Ty$ implies that $STy = TSy$; (ii) for sequence $\{x_n\}$ in X and $x \in X$, whenever $\{Sx_n\} \rightarrow x$, $\{Tx_n\} \rightarrow x$, then $STx_n \rightarrow Tx$, as $n \rightarrow \infty$, hold. However, in Fuzzy metric space (ii) implies (i), taking $x_n = y$ for all n and $x = Ty = Sy$. So, we define a semicompatible pair of self-maps in fuzzy metric space by condition (ii) only. Saliga [9] and Sharma et. al [10] proved some interesting fixed point results using implicit real functions and semicompatibility in d -complete topological spaces. Recently, Popa in [8] used the family F_4 of implicit real functions to find the fixed points of two pairs of semicompatible maps in a d -complete topological space. Here, F_4 denotes the family of all real continuous functions $F : (\mathbb{R}^+)^4 \rightarrow \mathbb{R}$ satisfying the following properties.

(F_h) There exists $h \geq 1$ such that for every $u \geq 0$, $v \geq 0$ with $F(u, v, u, v) \geq 0$ or $F(u, v, v, u) \geq 0$, we have $u \geq hv$.

(F_u) $F(u, u, 0, 0) < 0$, for all $u > 0$.

Jungck and Rhoades [6] (also Dhage [3]) termed a pair of self-maps to be coincidentally commuting or equivalently weak compatible if they commute at their coincidence points. This concept is most general among all the commutativity concepts in this field as every pair of commuting self-maps is R -weakly commuting, each pair of R -weakly commuting self-maps is compatible and each pair of compatible self-maps is weak compatible but the reverse is not always true. Similarly, every semicompatible pair of self-maps is weak compatible but the reverse is not true always. The main object of this paper is to obtain some fixed point theorems in the setting of fuzzy metric space using weak compatibility,

seicompatibility, and an implicit relation. In the sequel, we derive a characterization of this implicit relation if it is in linear form and use the same for obtaining some results in fixed points. For the sake of completeness, following Kramosil and Michálek [7] and Grabiec [5], we recall some definitions and known results in fuzzy metric space.

2. Preliminaries

Definition 2.1. A binary operation $* : [0,1]^2 \rightarrow [0,1]$ is called a continuous t -norm if $([0,1], *)$ is an abelian topological monoid with unit 1 such that $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$ for all a, b, c , and $d \in [0,1]$.

Examples of t -norm are $a * b = ab$ and $a * b = \min\{a, b\}$.

Definition 2.2 (Kramosil and Michálek [7]). The 3-tuple $(X, M, *)$ is called a fuzzy metric space if X is an arbitrary set, $*$ is a continuous t -norm, and M is a fuzzy set in $X^2 \times [0, \infty)$ satisfying the following conditions for all $x, y, z \in X$ and $s, t > 0$:

- (FM-1) $M(x, y, 0) = 0$;
- (FM-2) $M(x, y, t) = 1$, for all $t > 0$ if and only if $x = y$;
- (FM-3) $M(x, y, t) = M(y, x, t)$;
- (FM-4) $M(x, y, t) * M(y, z, s) \geq M(x, z, t + s)$;
- (FM-5) $M(x, y, \cdot) : [0, \infty) \rightarrow [0, 1]$ is left continuous.

Note that $M(x, y, t)$ can be thought of as the degree of nearness between x and y with respect to t . We identify $x = y$ with $M(x, y, t) = 1$ for all $t > 0$. The following example shows that every metric space induces a fuzzy metric space.

Example 2.3 (George and Veeramani [4]). Let (X, d) be a metric space. Define $a * b = \min\{a, b\}$ and for all $a, b \in X$,

$$M(x, y, t) = \frac{t}{t + d(x, y)}, \quad \forall t > 0, M(x, y, 0) = 0. \quad (2.1)$$

Then $(X, M, *)$ is a fuzzy metric space. It is called the fuzzy metric space induced by the metric d .

LEMMA 2.4 (Grabiec [5]). *For all $x, y \in X, M(x, y, \cdot)$ is a nondecreasing function.*

Definition 2.5 (Grabiec [5]). Let $(X, M, *)$ be a fuzzy metric space. A sequence $\{x_n\}$ in X is said to be convergent to a point $x \in X$ if $\lim_{n \rightarrow \infty} M(x_n, x, t) = 1$ for all $t > 0$. Further, the sequence $\{x_n\}$ is said to be a Cauchy sequence in X if $\lim_{n \rightarrow \infty} M(x_n, x_{n+p}, t) = 1$ for all $t > 0$ and $p > 0$. The space is said to be complete if every Cauchy sequence in it converges to a point of it.

Remark 2.6. Since $*$ is continuous, it follows from (FM-4) that the limit of a sequence in a fuzzy metric space is unique.

In this paper, $(X, M, *)$ is considered to be the fuzzy metric space with condition (FM-6) $\lim_{t \rightarrow \infty} M(x, y, t) = 1$, for all $x, y \in X$.

LEMMA 2.7 (Cho [1]). *Let $\{y_n\}$ be a sequence in a fuzzy metric space $(X, M, *)$ with the condition (FM-6). If there exists a number $k \in (0, 1)$ such that $M(y_{n+2}, y_{n+1}, kt) \geq M(y_{n+1}, y_n, t)$, for all $t > 0$, then $\{y_n\}$ is a Cauchy sequence in X .*

LEMMA 2.8. *Let A and B be two self-maps on a complete fuzzy metric space $(X, M, *)$ such that for some $k \in (0, 1)$, for all $x, y \in X$ and $t > 0$,*

$$M(Ax, By, kt) \geq \text{Min} \{M(x, y, t), M(Ax, x, t)\}. \quad (2.2)$$

Then A and B have a unique common fixed point in X .

Proof. Let $p \in X$. Taking $x_0 = p$, define sequence $\{x_n\}$ in X by $Ax_{2n} = x_{2n+1}$ and $Bx_{2n+1} = x_{2n+2}$. By taking $x = x_{2n}$, $y = x_{2n+1}$ and $x = x_{2n}$, $y = x_{2n-1}$, respectively, in the contractive condition, we obtain that

$$M(x_{n+1}, x_n, kt) \geq M(x_n, x_{n-1}, t), \quad \forall t > 0, \forall n. \quad (2.3)$$

Therefore by Lemma 2.7, $\{x_n\}$ is a Cauchy sequence in X , which is complete. Hence, $\{x_n\}$ converges to some u in X . Taking $x = x_{2n}$ and $y = u$ and letting $n \rightarrow \infty$ in the contractive condition, we get $Bu = u$. Similarly, by putting $x = u$ and $y = x_{2n+1}$, we get $Au = u$. Therefore, u is the common fixed point of the maps A and B . The uniqueness of the common fixed point follows from the contractive condition. \square

Definition 2.9. Let A and S be mappings from a fuzzy metric space $(X, M, *)$ into itself. The mappings are said to be weak compatible if they commute at their coincidence points, that is, $Ax = Sx$ implies that $ASx = SAx$.

Definition 2.10. Let A and S be mappings from a fuzzy metric space $(X, M, *)$ into itself. Then the mappings are said to be compatible if

$$\lim_{n \rightarrow \infty} M(ASx_n, SAx_n, t) = 1, \quad \forall t > 0, \quad (2.4)$$

whenever $\{x_n\}$ is a sequence in X such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = x \in X. \quad (2.5)$$

*PROPOSITION 2.11 [12]. Self-mappings A and S of a fuzzy metric space $(X, M, *)$ are compatible, then they are weak compatible.*

The converse is not true as seen in Example 2.16.

Definition 2.12. Let A and S be mappings from a fuzzy metric space $(X, M, *)$ into itself. Then the mappings are said to be semicompatible if

$$\lim_{n \rightarrow \infty} M(ASx_n, Sx, t) = 1, \quad \forall t > 0, \quad (2.6)$$

whenever $\{x_n\}$ is a sequence in X such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = x \in X. \quad (2.7)$$

It follows that if (A, S) is semicompatible and $Ay = Sy$, then $ASy = SAy$. Thus if the pair (A, S) is semicompatible, then it is weak compatible. The converse is not true as seen in Example 2.14.

PROPOSITION 2.13 [13]. *Let A and S be self-maps on a fuzzy metric space $(X, M, *)$. If S is continuous, then (A, S) is semicompatible if and only if (A, S) is compatible.*

The following is an example of a pair of self-maps (A, S) which is compatible but not semicompatible. Further, it is also seen here that the semicompatibility of the pair (A, S) need not imply the semicompatibility of (S, A) .

Example 2.14. Let $X = [0, 1]$ and let $(X, M, *)$ be the fuzzy metric space with $M(x, y, t) = [\exp|x - y|/t]^{-1}$, for all $x, y \in X, t > 0$. Define self-map S as follows:

$$Sx = \begin{cases} x & \text{if } 0 \leq x < \frac{1}{2}, \\ 1 & \text{if } x \geq \frac{1}{2}. \end{cases} \quad (2.8)$$

Let I be the identity map on X and $x_n = 1/2 - 1/n$. Then, $\{Ix_n\} = \{x_n\} \rightarrow 1/2$ and $\{Sx_n\} = \{x_n\} \rightarrow 1/2$. Thus, $\{ISx_n\} = \{Sx_n\} \rightarrow 1/2 \neq S(1/2)$. Hence (I, S) is not semicompatible. Again as (I, S) is commuting, it is compatible. Further, for any sequence $\{x_n\}$ in X such that $\{x_n\} \rightarrow x$ and $\{Sx_n\} \rightarrow x$, we have $\{Sx_n\} = \{Sx_n\} \rightarrow x = Ix$. Hence (S, I) is always semicompatible.

Remark 2.15. The above example gives an important aspect of semicompatibility as the pair of self-maps (I, S) is commuting, hence it is weakly commuting, compatible, and weak compatible yet it is not semicompatible. Further, it is to be noted that the pair (S, I) is semicompatible but (I, S) is not semicompatible here.

The following is an example of a pair of self-maps (A, S) which is semicompatible but not compatible.

Example 2.16. Let $(X, M, *)$ be a fuzzy metric space, where $X = [0, 2]$, with t -norm defined by $a * b = \min\{a, b\}$, for all $a, b \in [0, 1]$ and $M(x, y, t) = t/(t + d(x, y))$ for all $t > 0$ and $M(x, y, 0) = 0$, for all $x, y \in X$. Define self-maps A and S on X as follows:

$$Ax = \begin{cases} 2 & \text{if } 0 \leq x \leq 1, \\ \frac{x}{2} & \text{if } 1 < x \leq 2, \end{cases} \quad Sx = \begin{cases} 2 & \text{if } x = 1, \\ \frac{x+3}{5} & \text{otherwise,} \end{cases} \quad (2.9)$$

and $x_n = 2 - 1/(2n)$. Then we have $S(1) = A(1) = 2$ and $S(2) = A(2) = 1$. Also $SA(1) = AS(1) = 1$ and $SA(2) = AS(2) = 2$. Thus (A, S) is weak compatible. Again,

$$Ax_n = 1 - \frac{1}{4n}, \quad Sx_n = 1 - \frac{1}{10n}. \quad (2.10)$$

Thus,

$$Ax_n \rightarrow 1, \quad Sx_n \rightarrow 1. \quad (2.11)$$

Hence $u = 1$.

Further,

$$SAx_n = \frac{4}{5} - \frac{1}{20n}, \quad ASx_n = 2. \quad (2.12)$$

Now,

$$\begin{aligned} \lim_{n \rightarrow \infty} M(ASx_n, Su, t) &= M(2, 2, t) = 1, \\ \lim_{n \rightarrow \infty} M(ASx_n, SAx_n, t) &= \lim_{n \rightarrow \infty} M\left(2, \frac{4}{5} - \frac{1}{20n}, t\right) = \frac{t}{t + 6/5} < 1, \quad \forall t > 0. \end{aligned} \quad (2.13)$$

Hence (A, S) is semicompatible but it is not compatible.

For a detailed discussion of semicompatibility, we refer to [11, 13, 14].

2.1. A class of implicit relation. Let Φ be the set of all real continuous functions $\phi : (\mathbb{R}^+)^4 \rightarrow \mathbb{R}$, nondecreasing in first argument and satisfying the following conditions.

- (i) For $u, v \geq 0$, $\phi(u, v, v, u) \geq 0$ or $\phi(u, v, u, v) \geq 0$ implies that $u \geq v$.
- (ii) $\phi(u, u, 1, 1) \geq 0$ implies that $u \geq 1$.

Example 2.17. Define $\phi(t_1, t_2, t_3, t_4) = 15t_1 - 13t_2 + 5t_3 - 7t_4$. Then $\phi \in \Phi$.

3. Main results

THEOREM 3.1. Let A, B, S , and T be self-mappings of a complete fuzzy metric space $(X, M, *)$ satisfying that

$$A(X) \subseteq T(X), \quad B(X) \subseteq S(X); \quad (3.1)$$

the pair (A, S) is semicompatible and (B, T) is weak compatible; (3.2)

one of A or S is continuous; (3.3)

For some $\phi \in \Phi$, there exists $k \in (0, 1)$ such that for all $x, y \in X$ and $t > 0$,

$$\phi(M(Ax, By, kt), M(Sx, Ty, t), M(Ax, Sx, t), M(By, Ty, kt)) \geq 0, \quad (3.4)$$

$$\phi(M(Ax, By, kt), M(Sx, Ty, t), M(Ax, Sx, kt), M(By, Ty, t)) \geq 0. \quad (3.5)$$

Then A, B, S , and T have unique common fixed point in X .

Proof. Let $x_0 \in X$ be any arbitrary point as $A(X) \subseteq T(X)$ and $B(X) \subseteq S(X)$, there exist $x_1, x_2 \in X$ such that $Ax_0 = Tx_1, Bx_1 = Sx_2$. Inductively, construct sequences $\{y_n\}$ and $\{x_n\}$ in X such that $y_{2n+1} = Ax_{2n} = Tx_{2n+1}, y_{2n+2} = Bx_{2n+1} = Sx_{2n+2}$, for $n = 0, 1, 2, \dots$. Now using (3.4) with $x = x_{2n}, y = x_{2n+1}$, we get

$$\phi\left(\begin{array}{l} M(Ax_{2n}, Bx_{2n+1}, kt), M(Sx_{2n}, Tx_{2n+1}, t), \\ M(Ax_{2n}, Sx_{2n}, t), M(Bx_{2n+1}, Tx_{2n+1}, kt) \end{array}\right) \geq 0, \quad (3.6)$$

that is,

$$\phi\left(\begin{array}{l} M(y_{2n+1}, y_{2n+2}, kt), M(y_{2n}, y_{2n+1}, t), \\ M(y_{2n+1}, y_{2n}, t), M(y_{2n+2}, y_{2n+1}, kt) \end{array}\right) \geq 0. \quad (3.7)$$

Using (i), we get

$$M(y_{2n+2}, y_{2n+1}, kt) \geq M(y_{2n+1}, y_{2n}, t). \quad (3.8)$$

Similarly, by putting $x = x_{2n+2}$ and $y = x_{2n+1}$ in (3.5), we have

$$\phi \left(\begin{array}{l} M(y_{2n+3}, y_{2n+2}, kt), M(y_{2n+1}, y_{2n+2}, t), \\ M(y_{2n+3}, y_{2n+2}, kt), M(y_{2n+1}, y_{2n+2}, t) \end{array} \right) \geq 0. \quad (3.9)$$

Using (i), we get

$$M(y_{2n+3}, y_{2n+2}, kt) \geq M(y_{2n+1}, y_{2n+2}, t). \quad (3.10)$$

Thus, for any n and t , we have

$$M(y_n, y_{n+1}, kt) \geq M(y_{n-1}, y_n, t). \quad (3.11)$$

Hence by Lemma 2.7, $\{y_n\}$ is a Cauchy sequence in X , which is complete. Therefore $\{y_n\}$ converges to $u \in X$. Its subsequences $\{Ax_{2n}\}$, $\{Bx_{2n+1}\}$, $\{Sx_{2n}\}$, $\{Tx_{2n+1}\}$ also converge to u , that is,

$$\{Ax_{2n}\} \rightarrow u, \quad \{Bx_{2n+1}\} \rightarrow u. \quad (3.12)$$

$$\{Sx_{2n}\} \rightarrow u, \quad \{Tx_{2n+1}\} \rightarrow u. \quad (3.13)$$

Case I (S is continuous). In this case, we have

$$SAx_{2n} \rightarrow Su, \quad S^2x_{2n} \rightarrow Su. \quad (3.14)$$

The semicompatibility of the pair (A, S) gives

$$\lim_{n \rightarrow \infty} ASx_{2n} = Su. \quad (3.15)$$

Step 1. By putting $x = Sx_{2n}$, $y = x_{2n+1}$ in (3.4), we obtain that

$$\phi \left(\begin{array}{l} M(ASx_{2n}, Bx_{2n+1}, kt), M(SSx_{2n}, Tx_{2n+1}, t), \\ M(ASx_{2n}, SSx_{2n}, t), M(Bx_{2n+1}, Tx_{2n+1}, kt) \end{array} \right) \geq 0. \quad (3.16)$$

Letting $n \rightarrow \infty$, using (3.12), (3.14), (3.15), and the continuity of the t -norm $*$, we have

$$\phi(M(Su, u, kt), M(Su, u, t), M(Su, Su, t), M(u, u, kt)) \geq 0, \quad (3.17)$$

that is, $\phi(M(Su, u, kt), M(Su, u, t), 1, 1) \geq 0$.

As ϕ is nondecreasing in the first argument, we have

$$\phi(M(Su, u, t), M(Su, u, t), 1, 1) \geq 0. \quad (3.18)$$

Using (ii), we get $M(Su, u, t) \geq 1$, for all $t > 0$, which gives $M(Su, u, t) = 1$, that is,

$$Su = u. \quad (3.19)$$

Step 2. By putting $x = u$, $y = x_{2n+1}$ in (3.4), we obtain that

$$\phi \left(\begin{array}{l} M(Au, Bx_{2n+1}, kt), M(Su, Tx_{2n+1}, t), \\ M(Au, Su, t), M(Bx_{2n+1}, Tx_{2n+1}, kt) \end{array} \right) \geq 0. \quad (3.20)$$

Taking limit as $n \rightarrow \infty$ and using (3.12) and (3.19), we get

$$\phi(M(Au, u, kt), 1, M(Au, u, t), 1) \geq 0. \quad (3.21)$$

As ϕ is nondecreasing in the first argument, we have

$$\phi(M(Au, u, t), 1, M(Au, u, t), 1) \geq 0. \quad (3.22)$$

Using (i), we have $M(Au, u, t) \geq 1$, for all $t > 0$, which gives $u = Au$. Hence,

$$Au = u = Su. \quad (3.23)$$

Step 3. As $A(X) \subseteq T(X)$, there exists $w \in X$ such that $Au = Su = u = Tw$. By putting $x = x_{2n}$, $y = w$ in (3.4), we obtain that

$$\phi(M(Ax_{2n}, Bw, kt), M(Sx_{2n}, Tw, t), M(Ax_{2n}, Sx_{2n}, t), M(Bw, Tw, kt)) \geq 0. \quad (3.24)$$

Taking limit as $n \rightarrow \infty$ and using (3.12), we get

$$\phi(M(u, Bw, kt), 1, 1, M(Bw, u, kt)) \geq 0. \quad (3.25)$$

Using (i), we have $M(u, Bu, kt) \geq 1$, for all $t > 0$. Hence, $M(u, Bu, t) = 1$. Thus, $u = Bu$. Therefore $Bw = Tw = u$. Since (B, T) is weak compatible, we get that $TBw = BTw$, that is,

$$Bu = Tu. \quad (3.26)$$

Step 4. By putting $x = u$, $y = u$ in condition (3.4) and using (3.23) and (3.26), we obtain that

$$\phi(M(Au, Bu, kt), M(Su, Tu, t), M(Au, Su, t), M(Bu, Tu, kt)) \geq 0, \quad (3.27)$$

that is, $\phi(M(Au, Bu, kt), M(Au, Bu, t), 1, 1) \geq 0$.

As ϕ is nondecreasing in the first argument, we have

$$\phi(M(Au, Bu, t), M(Au, Bu, t), 1, 1) \geq 0. \quad (3.28)$$

Using (ii), we have $M(Au, Bu, t) \geq 1$, for all $t > 0$.

Thus, $M(Au, Bu, t) = 1$, we have, $Bu = Au$.

Therefore, $u = Au = Su = Bu = Tu$, that is, u is a common fixed point of A, B, S , and T .

Case II (A is continuous). In this case, we have

$$ASx_{2n} \longrightarrow Au. \quad (3.29)$$

The semicompatibility of the pair (A, S) gives

$$ASx_{2n} \longrightarrow Su. \quad (3.30)$$

By uniqueness of limit in fuzzy metric space, we obtain that $Au = Su$.

Step 5. By putting $x = u$, $y = x_{2n+1}$ in condition (3.4), we obtain that

$$\phi \left(\frac{M(Au, Bx_{2n+1}, kt), M(Su, Tx_{2n+1}, t)}{M(Au, Su, t), M(Bx_{2n+1}, Tx_{2n+1}, kt)} \right) \geq 0. \quad (3.31)$$

Taking limit as $n \rightarrow \infty$ and using (3.12) and (3.19), we get

$$\phi(M(Au, u, kt), 1, M(Au, u, t), 1) \geq 0. \quad (3.32)$$

As ϕ is nondecreasing in the first argument, we have

$$\phi(M(Au, u, t), 1, M(Au, u, t), 1) \geq 0. \quad (3.33)$$

Using (i), we have $M(Au, u, t) \geq 1$, for all $t > 0$, which gives $u = Au$ and the rest of the proof follows from Step 3 onwards of the previous case.

Uniqueness. Let z be another common fixed point of A, B, S , and T .

Then $z = Az = Bz = Sz = Tz$.

Putting $x = u$ and $y = z$ in (3.4), we get

$$\phi(M(Au, Bz, kt), M(Su, Tz, t), M(Au, Su, t), M(Bz, Tz, kt)) \geq 0, \quad (3.34)$$

that is, $\phi(M(u, z, kt), M(u, z, t), 1, 1) \geq 0$.

As ϕ is nondecreasing in the first argument, we have

$$\phi(M(u, z, t), M(u, z, t), 1, 1) \geq 0. \quad (3.35)$$

Using (i), we have $M(u, z, t) \geq 1$, for all $t > 0$.

Hence $M(u, z, t) = 1$, that is, $u = z$. Therefore, u is the unique common fixed point of the self-maps A, B, S , and T . \square

COROLLARY 3.2. *Let A, B, S , and T be self-mappings of a complete fuzzy metric space $(X, M, *)$ satisfying conditions (3.1), (3.4), (3.5), and that*

$$\begin{aligned} & \text{the pairs } (A, S) \text{ and } (B, T) \text{ are semicompatible;} \\ & \text{one of } A, B, S, \text{ or } T \text{ is continuous.} \end{aligned} \quad (3.36)$$

Then A, B, S , and T have a unique common fixed point in X .

Proof. As semicompatibility implies weak compatibility, the proof follows from Theorem 3.1. \square

On taking $A = B$ in Theorem 3.1, we have the following corollary.

COROLLARY 3.3. *Let A , S , and T be self-mappings of a complete fuzzy metric space $(X, M, *)$ satisfying that*

$$\begin{aligned} A(X) &\subseteq T(X) \cap S(X); \\ \text{the pair } (A, S) &\text{ is semicompatible and } (A, T) \text{ is weak compatible;} \\ \text{one of } A \text{ or } S &\text{ is continuous.} \end{aligned} \quad (3.37)$$

For some $\phi \in \Phi$, there exists $k \in (0, 1)$ such that for all $x, y \in X$ and $t > 0$,

$$\begin{aligned} \phi(M(Ax, Ay, kt), M(Sx, Ty, t), M(Ax, Sx, t), M(Ay, Ty, kt)) &\geq 0, \\ \phi(M(Ax, Ay, kt), M(Sx, Ty, t), M(Ax, Sx, kt), M(Ay, Ty, t)) &\geq 0. \end{aligned} \quad (3.38)$$

Then A , S , and T have a unique common fixed point in X .

Now, taking $S = I$ and $T = I$ in Theorem 3.1, the conditions (3.1), (3.2), (3.3) are satisfied trivially, and we get the following corollary.

COROLLARY 3.4. *Let A and B be self-mappings of a complete fuzzy metric space $(X, M, *)$ such that for some $\phi \in \Phi$, there exists some $k \in (0, 1)$ such that for all $x, y \in X$ and for all $t > 0$,*

$$\phi(M(Ax, By, kt), M(x, y, t), M(Ax, x, t), M(By, y, kt)) \geq 0, \quad (3.39)$$

$$\phi(M(Ax, By, kt), M(x, y, t), M(Ax, x, kt), M(By, y, t)) \geq 0. \quad (3.40)$$

Then A and B have a unique common fixed point in X .

THEOREM 3.5. *Let A , B , S , and T be self-mappings of a complete fuzzy metric space $(X, M, *)$ satisfying conditions (3.1), (3.3), (3.4), (3.5), and that*

$$\text{the pair } (A, S) \text{ is compatible and } (B, T) \text{ is weak compatible.} \quad (3.41)$$

Then A , B , S , and T have a unique common fixed point in X .

Proof. In view of Proposition 2.13 and Theorem 3.1, it suffices to prove the result when A is continuous. As in the proof of Theorem 3.1, the sequence $\{y_n\} \rightarrow u \in X$ and (3.12) and (3.13) are satisfied. As A is continuous, we have

$$ASx_{2n} \rightarrow Au, \quad AAx_{2n} \rightarrow Au. \quad (3.42)$$

The compatibility of (A, S) gives

$$\lim_{n \rightarrow \infty} ASx_{2n} = Au = \lim_{n \rightarrow \infty} SAx_{2n}. \quad (3.43)$$

Step I. By putting $x = Ax_{2n}$ and $y = x_{2n+1}$ in condition (3.4), we get that

$$\phi \left(\begin{aligned} M(AAx_{2n}, Bx_{2n+1}, kt), M(SAx_{2n}, Tx_{2n+1}, t), \\ M(AAx_{2n}, SAx_{2n}, t), M(Bx_{2n+1}, Tx_{2n+1}, kt) \end{aligned} \right) \geq 0. \quad (3.44)$$

Letting $n \rightarrow \infty$, using (3.12) and (3.13), we obtain that

$$\phi(M(Au, u, kt), M(Au, u, t), M(Au, Au, t), M(u, u, kt)) \geq 0, \quad (3.45)$$

that is, $\phi(M(Au, u, kt), M(Au, u, t), 1, 1) \geq 0$, that is, $\phi(M(Au, u, t), M(Au, u, t), 1, 1) \geq 0$.

Using (ii), we have $M(Au, u, t) \geq 1$, $t > 0$. Therefore $u = Au$.

Step II. As $A(X) \subseteq T(X)$, there exists $w \in X$ such that $Au = Su = u = Tw$. By putting $x = x_{2n}$, $y = w$ in (3.4), we obtain that

$$\phi(M(Ax_{2n}, Bw, kt), M(Sx_{2n}, Tw, t), M(Ax_{2n}, Sx_{2n}, t), M(Bw, Tw, kt)) \geq 0. \quad (3.46)$$

Taking limit as $n \rightarrow \infty$ and using (3.12), we get

$$\phi(M(u, Bw, kt), 1, 1, M(Bw, u, kt)) \geq 0. \quad (3.47)$$

Using (i), we have $M(u, Bu, kt) \geq 1$, for all $t > 0$. Hence, $M(u, Bu, t) = 1$. Thus, $u = Bw$. Therefore $Bw = Tw = u$. As (B, T) is weak compatible, we have $TBw = BTw$, that is, $Bu = Tu$.

Step III. Again as $u = Bw$ and $B(X) \subseteq S(X)$, there exists $v \in X$ such that $u = Bw = Sv$. By putting $x = v$, $y = w$ in (3.4), we have

$$\phi(M(Av, Bw, kt), M(Sv, Tw, t), M(Av, Sv, t), M(Bw, Tw, kt)) \geq 0, \quad (3.48)$$

that is, $\phi(M(Av, Sv, kt), 1, M(Av, Sv, t), 1) \geq 0$, that is, $\phi(M(Av, Sv, t), 1, M(Av, Sv, t), 1) \geq 0$.

Using (i), we have $M(Av, Sv, t) \geq 1$, for all $t > 0$.

This gives $Av = Sv$. As (A, S) is compatible, we have $ASv = SAv$ or $Au = Su = u$. Also $Au = Bu$ follows from Step 4 in the proof of Theorem 3.1 and it follows that u is a common fixed point of the four maps A , B , S , and T . The uniqueness follows as in the proof of Theorem 3.1. \square

COROLLARY 3.6. *Let A , B , S , and T be self-mappings of a complete fuzzy metric space $(X, M, *)$ satisfying conditions (3.1), (3.4), (3.5), and that*

$$\begin{aligned} &\text{the pairs } (A, S) \text{ and } (B, T) \text{ are compatible;} \\ &\text{one of } A, B, S, \text{ or } T \text{ is continuous.} \end{aligned} \quad (3.49)$$

Then A , B , S , and T have a unique common fixed point in X .

Proof. As compatibility implies weak compatibility, the proof follows from Theorem 3.5. \square

If we take $A = I$, the identity map on X in Theorem 3.5, we have the following result for three self-maps, none of which is continuous and just a pair of them is needed to be weak compatible.

COROLLARY 3.7. *Let B , S , and T be self-mappings of a complete fuzzy metric space $(X, M, *)$, satisfying that*

$$\begin{aligned} &\text{the pair } (B, T) \text{ is weak compatible;} \\ &T \text{ is surjective.} \end{aligned} \quad (3.50)$$

For some $\phi \in \Phi$, there exists $k \in (0, 1)$ such that for all $x, y \in X$ and $t > 0$,

$$\begin{aligned} &\phi(M(x, By, kt), M(Sx, Ty, t), M(x, Sx, t), M(By, Ty, kt)) \geq 0, \\ &\phi(M(x, By, kt), M(Sx, Ty, t), M(x, Sx, kt), M(By, Ty, t)) \geq 0. \end{aligned} \quad (3.51)$$

Then B , S , and T have a unique common fixed point in X .

Now, taking $A = I$ and $B = I$ in Corollary 3.6, the condition (3.49) is satisfied trivially and we get an important result for surjective maps as follows.

COROLLARY 3.8. *Let S and T be two surjective self-mappings of a complete fuzzy metric space $(X, M, *)$ such that for some $\phi \in \Phi$, there exists some $k \in (0, 1)$ satisfying*

$$\begin{aligned} &\phi(M(x, y, kt), M(Sx, Ty, t), M(x, Sx, t), M(y, Ty, kt)) \geq 0, \\ &\phi(M(x, y, kt), M(Sx, Ty, t), M(x, Sx, kt), M(y, Ty, t)) \geq 0, \end{aligned} \quad (3.52)$$

for all $x, y \in X$ and for all $t > 0$.

Then S and T have a unique common fixed point in X .

3.1. A characterization of Φ in linear form. Define $\phi(t_1, t_2, t_3, t_4) = at_1 + bt_2 + ct_3 + dt_4$, where $a, b, c, d \in \mathbb{R}$ with $a + b + c + d = 0$, $a > 0$, $a + c > 0$, $a + b > 0$, and $a + d > 0$. Then $\phi \in \Phi$.

Proof. For $u, v \geq 0$ and $\phi(u, v, v, u) \geq 0$, we have

$$(a + d)u + (b + c)v \geq 0, \quad (3.53)$$

that is, $(a + d)u \geq (a + d)v$. Hence $u \geq v$, since $a + d > 0$.

Again,

$$\phi(u, v, u, v) \geq 0 \quad (3.54)$$

gives

$$(a + c)u + (b + d)v \geq 0, \quad (3.55)$$

that is, $(a + c)u - (a + c)v \geq 0$. Hence, $u \geq v$ as $(a + c) > 0$.

Also, $\phi(u, u, 1, 1) \geq 0$ gives

$$(a + b)u + (c + d) \geq 0, \quad (3.56)$$

that is, $(a + b)u \geq -(c + d)$, that is, $(a + b)u \geq (a + b)$, as $a + b + c + d = 0$. Hence, $u \geq 1$.

As $a > 0$, ϕ is nondecreasing in the first argument and the result follows. \square

COROLLARY 3.9. *Let A and B be self-mappings of a complete fuzzy metric space $(X, M, *)$ such that there exists some $k \in (0, 1)$ satisfying*

$$aM(Ax, By, kt) + bM(x, y, t) + cM(Ax, x, t) + dM(By, y, kt) \geq 0, \quad (3.57)$$

$$aM(Ax, By, kt) + bM(x, y, t) + cM(Ax, x, kt) + dM(By, y, t) \geq 0, \quad (3.58)$$

for all $x, y \in X$, for all $t > 0$, and for some fixed $a, b, c, d \in \mathbb{R}$ such that $a > 0$, $a + b > 0$, $a + c > 0$, $a + d > 0$, and $a + b + c + d = 0$.

Then A and B have a unique common fixed point in X .

Proof. Using the characterization of Φ in Corollary 3.4, the result follows. \square

COROLLARY 3.10. *Let A and B be self-mappings of a complete fuzzy metric space $(X, M, *)$ such that there exists some $k \in (0, 1)$ satisfying*

$$M(Ax, By, kt) \geq b_0M(x, y, t) + c_0M(Ax, x, t), \quad \forall x, y \in X, \forall t > 0, \quad (3.59)$$

where $b_0, c_0 \in (0, 1)$ with $b_0 + c_0 = 1$. Then A and B have a unique common fixed point in X .

Proof. Choosing $a = 1$, $d = 0$, $b = -b_0$, and $c = -c_0$, $c_0 > 0$, in Corollary 3.9 and using the fact that $M(x, y, \cdot)$ is a nondecreasing function, the second condition of Corollary 3.9 is trivially satisfied and the result follows. \square

Combining Corollary 3.10 and Lemma 2.8, we have the following important result.

COROLLARY 3.11. *Let A and B be self-mappings of a complete fuzzy metric space $(X, M, *)$ such that there exists some $k \in (0, 1)$ satisfying*

$$M(Ax, By, kt) \geq b_0M(x, y, t) + c_0M(Ax, x, t), \quad \forall x, y \in X, \forall t > 0, \quad (3.60)$$

where $b_0, c_0 \in [0, 1]$ with $b_0 + c_0 = 1$. Then A and B have a unique common fixed point in X .

COROLLARY 3.12. *Let A be self-mapping of a complete fuzzy metric space $(X, M, *)$ such that there exists some $k \in (0, 1)$ satisfying*

$$M(Ax, Ay, kt) \geq b_0M(x, y, t) + c_0M(Ax, x, t), \quad \forall x, y \in X, \forall t > 0, \quad (3.61)$$

where $b_0, c_0 \in [0, 1]$ with $b_0 + c_0 = 1$. Then A has a unique fixed point in X .

Proof. Taking $A = B$ in Corollary 3.11, the result follows. \square

Remark 3.13. If we take $b_0 = 1$ and $c_0 = 0$ in Corollary 3.12, we get the Banach contraction principle in the setting of fuzzy metric space as given by Grabiec in [5].

Acknowledgment

Authors thank Shobha Jain, Department of Mathematics, MB Khalsa College, Devi Ahilya University, Indore, for her cooperation in the preparation of this paper.

References

- [1] Y. J. Cho, *Fixed points in fuzzy metric spaces*, J. Fuzzy Math. **5** (1997), no. 4, 949–962.
- [2] Y. J. Cho, B. K. Sharma, and D. R. Sahu, *Semi-compatibility and fixed points*, Math. Japon. **42** (1995), no. 1, 91–98.
- [3] B. C. Dhage, *On common fixed points of pairs of coincidentally commuting mappings in D-metric spaces*, Indian J. Pure Appl. Math. **30** (1999), no. 4, 395–406.
- [4] A. George and P. Veeramani, *On some results in fuzzy metric spaces*, Fuzzy Sets and Systems **64** (1994), no. 3, 395–399.
- [5] M. Grabiec, *Fixed points in fuzzy metric spaces*, Fuzzy Sets and Systems **27** (1988), no. 3, 385–389.
- [6] G. Jungck and B. E. Rhoades, *Fixed points for set valued functions without continuity*, Indian J. Pure Appl. Math. **29** (1998), no. 3, 227–238.
- [7] I. Kramosil and J. Michálek, *Fuzzy metrics and statistical metric spaces*, Kybernetika (Prague) **11** (1975), no. 5, 336–344.
- [8] V. Popa, *Fixed points for non-surjective expansion mappings satisfying an implicit relation*, Bul. Ştiinț. Univ. Baia Mare Ser. B Fasc. Mat.-Inform. **18** (2002), no. 1, 105–108.
- [9] L. M. Saliga, *Fixed point theorems for non-self maps in d-complete topological spaces*, Int. J. Math. Math. Sci. **19** (1996), no. 1, 103–110.
- [10] B. K. Sharma, D. R. Sahu, M. Bounias, and A. Bonaly, *Fixed points for non-surjective expansion mappings*, Int. J. Math. Math. Sci. **21** (1998), no. 2, 277–288.
- [11] B. Singh and S. Jain, *Semi-compatibility and fixed point theorems in Menger space*, Journal of the Chungcheong Mathematical Society **17** (2004), no. 1, 1–17.
- [12] ———, *A fixed point theorem in Menger space through weak compatibility*, J. Math. Anal. Appl. **301** (2005), no. 2, 439–448.
- [13] ———, *Semi-compatibility, compatibility and fixed point theorems in fuzzy metric space*, Journal of the Chungcheong Mathematical Society **18** (2005), no. 1, 1–23.
- [14] B. Singh, S. Jain, and S. Jain, *Semi-compatibility and fixed point theorems in an unbounded D-metric space*, Int. J. Math. Math. Sci. **2005** (2005), no. 5, 789–801.

Bijendra Singh: School of Studies in Mathematics, Vikram University, University Road Ujjain, 456010 Madhya Pradesh, India

E-mail address: bijendrasingh@yahoo.com

Shishir Jain: Shri Vaishnav Institute of Technology and Science, Indore 453331, Madhya Pradesh, India

E-mail address: jainshishir11@rediffmail.com

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be