

BOUNDARY VALUE PROBLEMS FOR THE DIFFUSION EQUATION WITH PIECEWISE CONTINUOUS TIME DELAY

JOSEPH WIENER

Department of Mathematics
University of Texas - Pan American
Edinburg, Texas 78539, U.S.A

LOKENATH DEBNATH

Department of Mathematics
University of Central Florida
Orlando, Florida 32816, U S A

(Received November 30, 1995 and in revised form January 20, 1996)

ABSTRACT. A study is made of partial differential equations with piecewise constant arguments. Boundary value problems for three types of equations are discussed: delayed; alternately of advanced and retarded type; and most importantly, an equation of neutral type (that is, including the derivative at different values of time t).

KEY WORDS AND PHRASES: Partial differential equations, piecewise constant arguments, oscillations, and stability.

1991 AMS SUBJECT CLASSIFICATION CODES: 35A05, 35B25, 35L10, 34K25.

1. INTRODUCTION

Functional differential equations (FDE) with delay provide a mathematical model for a physical or biological system in which the rate of change of the system depends upon its past history. The theory of FDE with continuous arguments is well developed and has numerous applications in natural and engineering sciences. This paper continues our earlier work in an attempt to extend this theory to differential equations with discontinuous argument deviations. In articles [1-5], ordinary differential equations with arguments having intervals of constancy have been studied. Such equations represent a hybrid of continuous and discrete dynamical systems and combine properties of both differential and difference equations. They include as particular cases loaded and impulsive equations, hence their importance in control theory and in certain biomedical models. Continuity of a solution at a point joining any two consecutive intervals implies recursion relations for the values of the solution at such points. Therefore, differential equations with piecewise continuous argument (EPCA) are intrinsically closer to difference rather than to differential equations. In [6] boundary value problems for some linear EPCA in partial derivatives were considered and the behavior of their solutions studied. The results were also extended to equations with positive definite operators in Hilbert spaces. In [7] initial value problems were studied for EPCA in partial derivatives. A class of loaded equations that arise in solving certain inverse problems was explored within the general framework of differential equations with piecewise continuous delay. The purpose of the present note is to investigate the asymptotic behavior of the solutions, especially their oscillatory properties, of a boundary value problem for some EPCA of parabolic type. For a rather comprehensive addition to the growing body of literature on EPCA the reader is referred to [8].

2. A COMPARISON OF TWO EPCA

The equation

$$u_t(x, t) = a^2 u_{xx}(x, t) - bu(x, t) \quad (2.1)$$

describes heat flow in a rod with both diffusion $a^2 u_{xx}$ along the rod and heat loss (or gain) across the lateral sides of the rod. Measuring the lateral heat change at discrete moments of time leads to the equation with piecewise continuous delay

$$u_t(x, t) = a^2 u_{xx}(x, t) - bu(x, [t]), \quad (2.2)$$

which was investigated in [6]. Here $[\cdot]$ designates the greatest integer function and $(x, t) \in [0, 1] \times [0, \infty)$. The problem posed in [6] for Eq. (2.2) consists of the boundary conditions

$$u(0, t) = 0, \quad u(1, t) = 0 \quad (2.3)$$

and the initial condition

$$u(x, 0) = u_0(x), \quad (2.4)$$

and the solution is sought in the form

$$u(x, t) = X(x)T(t). \quad (2.5)$$

Then separation of variables leads to the BVP

$$X' + \lambda^2 X = 0, \quad X(0) = X(1) = 0, \quad (2.6)$$

with the orthonormal set of solutions

$$X_j(x) = \sqrt{2} \sin(\pi jx) \quad (2.7)$$

on $[0, 1]$, and to the equation

$$T'_j(t) = -a^2 \pi^2 j^2 T_j(t) - bT_j([t]). \quad (2.8)$$

Let $T_{nj}(t)$ denote a solution of (2.8) on the interval $n \leq t < n + 1$, where n is a nonnegative integer. Then

$$T'_{nj}(t) = -a^2 \pi^2 j^2 T_{nj}(t) - bT_{nj}(n), \quad (2.9)$$

and the general solution of this equation is

$$T_{nj}(t) = C_{nj} e^{-a^2 \pi^2 j^2 (t-n)} - \frac{b}{a^2 \pi^2 j^2} T_{nj}(n). \quad (2.10)$$

We put here $t = n$ and get

$$C_{nj} = \left(1 + \frac{b}{a^2 \pi^2 j^2} \right) T_{nj}(n),$$

that is,

$$T_{nj}(t) = E_j(t-n) T_{nj}(n), \quad (2.11)$$

where

$$E_j(t) = e^{-a^2 \pi^2 j^2 t} - \frac{b}{a^2 \pi^2 j^2} \left(1 - e^{-a^2 \pi^2 j^2 t} \right). \quad (2.12)$$

At $t = n + 1$ we have

$$T_{nj}(n+1) = E_j(1) T_{nj}(n)$$

and since

$$T_{nj}(n+1) = T_{n+1,j}(n+1),$$

then

$$T_{n+1,j}(n+1) = E_j(1)T_{nj}(n)$$

and

$$T_{nj}(n) = E_n^n(1)T_{0,j}(0). \quad (2.13)$$

Therefore,

$$T_{nj}(t) = E_j(t-n)E_j^n(1)T_{0,j}(0) \quad (2.14)$$

and

$$u_n(x, t) = \sum_{j=1}^{\infty} \sqrt{2} E_j^n(1)T_{0,j}(0)E_j(t-n)\sin(\pi jx), \quad (2.15)$$

where $u_n(x, t)$ designates the solution of BVP (2.2), (2.3), (2.4) in $[0, 1] \times [n, n+1]$. Putting $t = 0$, $n = 0$ gives

$$u_0(x) = \sum_{j=1}^{\infty} T_{0,j}(0)\sqrt{2} \sin(\pi jx)$$

and

$$T_{0,j}(0) = \sqrt{2} \int_0^1 u_0(x)\sin(\pi jx)dx.$$

Along with Eq. (2.2), we study the equation

$$u_t(x, t) = a^2 u_{xx}(x, t) - bu\left(x, \left[t + \frac{1}{2}\right]\right) \quad (2.16)$$

under conditions (2.3), (2.4)

DEFINITION. A function $u(x, t)$ is said to be a solution of the above BVP if it satisfies the conditions (i) $u(x, t)$ is continuous in $G = [0, 1] \times [0, \infty)$; (ii) u_t and u_{xx} exist and are continuous in G , with the possible exception of the points $(x, n + \frac{1}{2})$, where one-sided derivatives exist ($n = 0, 1, 2, \dots$); (ii) $u(x, t)$ satisfies Eq. (2.16) in G , with the possible exception of the points $(x, n + \frac{1}{2})$, and conditions (2.3), (2.4).

Again, the solution is sought in form (2.5), and separation of variables generates the eigenfunctions (2.7) and leads to the EPCA

$$T'_j(t) = -a^2 \pi^2 j^2 T_j(t) - bT_j\left(\left[t + \frac{1}{2}\right]\right). \quad (2.17)$$

Eq. (2.17) is of considerable interest, since the argument deviation

$$\tau(t) = t - \left[t + \frac{1}{2}\right]$$

changes the sign in each interval $(n - \frac{1}{2}, n + \frac{1}{2})$, with integer n . Indeed, $\tau(t) < 0$ for $n - \frac{1}{2} \leq t < n$ and $\tau(t) > 0$ for $n < t < n + \frac{1}{2}$, which means that Eq. (2.17) is alternately of advanced and retarded type. Assume that $T_{nj}(t)$ is a solution of (2.17) on the interval $[n - \frac{1}{2}, n + \frac{1}{2}]$. Then Eq. (2.17) changes to (2.9), with the solution (2.11). At $t = n + \frac{1}{2}$ we have

$$T_{nj}\left(n + \frac{1}{2}\right) = E_j\left(\frac{1}{2}\right)T_{nj}(n)$$

and since

$$T_{n+1,j}(t) = E_j(t - n - 1)T_{n+1,j}(n + 1),$$

then

$$T_{n+1,j}\left(n + \frac{1}{2}\right) = E_j\left(-\frac{1}{2}\right)T_{n+1,j}(n + 1).$$

Furthermore, continuity of the solution $T_j(t)$ implies

$$T_{n+1,j}\left(n + \frac{1}{2}\right) = T_{nj}\left(n + \frac{1}{2}\right),$$

and therefore

$$E_j\left(-\frac{1}{2}\right)T_{n+1,j}(n + 1) = E_j\left(\frac{1}{2}\right)T_{nj}(n),$$

whence

$$T_{n+1,j}(n + 1) = \frac{E_j(1/2)}{E_j(-1/2)} T_{nj}(n).$$

From here,

$$T_{nj}(n) = E_j^n\left(\frac{1}{2}\right)E_j^{-n}\left(-\frac{1}{2}\right)T_{0j}(0) \quad (2.18)$$

and

$$T_{nj}(t) = E_j(t - n)E_j^n\left(\frac{1}{2}\right)E_j^{-n}\left(-\frac{1}{2}\right)T_{0j}(0). \quad (2.19)$$

The solution $u_n(x, t)$ of BVP (2.17), (2.3), (2.4) in the region $[0, 1] \times [n - \frac{1}{2}, n + \frac{1}{2}]$ is given by the formula

$$u_n(x, t) = \sum_{j=1}^{\infty} \sqrt{2} E_j^n\left(\frac{1}{2}\right) E_j^{-n}\left(-\frac{1}{2}\right) T_{0j}(0) E_j(t - n) \sin(\pi j x), \quad (2.20)$$

where $E_j(t)$ is defined by (2.12).

THEOREM 2.1. For

$$-a^2\pi^2 < b < a^2\pi^2 \left(e^{a^2\pi^2} + 1 \right) / \left(e^{a^2\pi^2} - 1 \right) \quad (2.21)$$

the solution (2.15) of Eq. (2.2) tends to zero as $t \rightarrow \infty$, uniformly with respect to x .

PROOF. From (2.15) it follows that the assertion is true if $|E_j(1)| < 1$. Solving the inequalities

$$-1 < e^{-a^2\pi^2 j^2} - \frac{b}{a^2\pi^2 j^2} \left(1 - e^{-a^2\pi^2 j^2} \right) < 1$$

for b proves the proposition. Furthermore, $u(x, t)$ approaches zero with an exponential rate

THEOREM 2.2. For

$$b > -a^2\pi^2 \quad (2.22)$$

the solution (2.20) of Eq. (2.16) tends to zero as $t \rightarrow \infty$, uniformly with respect to x .

PROOF. From (2.20) we see that the assertion is valid if $|E_j(1/2)E_j^{-1}(-1/2)| < 1$, where $E_j(t)$ is given by (2.12). Considering the case $E_j(-1/2) > 0$ gives

$$b > -2\mu_j e^{\mu_j} / (e^{\mu_j} - 1), \quad (2.23)$$

with the notation

$$\mu_j = a^2 \pi^2 j^2 / 2. \quad (2.24)$$

From $E_j(1/2) < E_j(-1/2)$ we obtain the inequality

$$b > -2\mu_j, \quad (2.25)$$

which is stronger than (2.23), and so (2.23) should be omitted. Furthermore, $E_j(1/2) > -E_j(-1/2)$ implies that

$$b > -2\mu_j(e^{\mu_j} + e^{-\mu_j}) / (e^{\mu_j} + e^{-\mu_j} - 2),$$

and since this inequality is weaker than (2.25) it may be disregarded. On the other hand, the case $E_j(-1/2) < 0$ is equivalent to

$$b < -2\mu_j e^{\mu_j} / (e^{\mu_j} - 1), \quad (2.26)$$

and the inequality $E_j(1/2) > E_j(-1/2)$ leads to

$$b < -2\mu_j.$$

The latter inequality may be ignored because it is weaker than (2.26). From $E_j(1/2) < -E_j(-1/2)$ it follows that

$$b < -2\mu_j(e^{\mu_j} + e^{-\mu_j}) / (e^{\mu_j} + e^{-\mu_j} - 2).$$

The latter inequality should be omitted since (2.26) is more stringent. On the other hand, (2.26) cannot hold true for all values of j , and we must retain only (2.25). This inequality is valid for all j if (2.22) takes place. Comparing inequalities (2.21) and (2.22) shows that the stability interval for Eq. (2.17) is larger than the stability interval for Eq. (2.8).

THEOREM 2.3. Each solution of Eq. (2.8) has a zero in the interval $[n, n + 1]$ if

$$b > a^2 \pi^2 / (e^{a^2 \pi^2} - 1). \quad (2.27)$$

The same is true for Eq. (2.17) if

$$b > (a\pi)^2 / (e^{a^2 \pi^2/2} - 1). \quad (2.28)$$

PROOF. From Eq. (2.13) we have

$$T_{nj}(n + 1) = E_j^{n+1}(1)T_{0j}(0)$$

and

$$T_{nj}(n)T_{nj}(n + 1) = E_j^{2n+1}(1)T_{0j}^2(0).$$

Hence,

$$T_{nj}(n)T_{nj}(n + 1) < 0 \quad \text{if} \quad E_j(1) < 0.$$

The latter inequality is equivalent to

$$b > a^2 \pi^2 j^2 / (e^{a^2 \pi^2 j^2} - 1),$$

which holds true for all j under condition (2.27). On the other hand, (2.28) is derived by examining the inequality

$$E_j(1/2)E_j(-1/2) < 0.$$

Inequalities (2.27) and (2.28) represent sufficient conditions of oscillation for Eqs (2.8) and (2.17), for all values of j simultaneously

THEOREM 2.4. Each solution of Eq. (2.8) is nonoscillatory, for all j , if $b < 0$. For any $b > 0$ and sufficiently large j , the functions $T_{nj}(t)$ are oscillatory.

PROOF. The inequality $E_j(1) > 0$ is equivalent to

$$b < a^2 \pi^2 j^2 / (e^{a^2 \pi^2 j^2} - 1)$$

and holds true for all j only if $b < 0$. Since its right-hand side tends to zero as $j \rightarrow \infty$, the above inequality breaks down for $b > 0$ and all sufficiently large j . Therefore, in this case the solutions $T_{nj}(t)$ of (2.8) oscillate, in sharp contrast to the functions $T_j(t)$ in the Fourier expansion for the solution of the equation $u_t = a^2 u_{xx} - bu$ without time delay.

THEOREM 2.5. Each solution of Eq. (2.17) is nonoscillatory, for all j , if

$$-a^2 \pi^2 / (e^{a^2 \pi^2/2} - 1) < b < 0. \quad (2.29)$$

For any $b > 0$ and sufficiently large j , the functions $T_{nj}(t)$ are oscillatory. Furthermore, if

$$b < -a^2 \pi^2 m^2 / (e^{a^2 \pi^2 m^2/2} - 1)$$

then the functions $T_{n1}(t), \dots, T_{nm}(t)$ oscillate.

PROOF. From the inequalities $E_j(-1/2) > 0$ and $E_j(1/2) > 0$ it follows that

$$-2\mu_j e^{\mu_j} / (e^{\mu_j} - 1) < b < 2\mu_j / (e^{\mu_j} - 1),$$

where μ_j is given by (2.24). The left of these inequalities holds true for all j if

$$b > -a^2 \pi^2 / (e^{a^2 \pi^2/2} - 1),$$

and the right one is satisfied for all j if $b < 0$. The inequalities $E_j(-1/2) < 0$ and $E_j(1/2) < 0$ are contradictory. On the other hand, the right side of the inequality

$$b < 2\mu_j / (e^{\mu_j} - 1)$$

tends to zero as $j \rightarrow \infty$, and therefore the inequality breaks down for any given $b > 0$ and all sufficiently large j . Moreover, the left side of the inequality

$$-2\mu_j e^{\mu_j} / (e^{\mu_j} - 1) < b$$

monotonically decreases to $-\infty$ as $j \rightarrow \infty$. Hence, if this inequality fails for $j = m$, the same is true also for $j < m$.

THEOREM 2.6. If, for some integer $m \geq 1$,

$$b = a^2 \pi^2 m^2 (e^{a^2 \pi^2 m^2} + 1) / (e^{a^2 \pi^2 m^2} - 1), \quad (2.30)$$

then the solution $T_m(t)$ of Eq. (2.8) is a periodic function with period 2. The same is true for the solution of Eq. (2.17) if

$$b = 2\mu_m (e^{\mu_m} + e^{-\mu_m}) / (2 - e^{\mu_m} - e^{-\mu_m}), \quad (2.31)$$

where μ_m is given in (2.24).

PROOF. Since the initial value problem for Eq. (2.8) has a unique solution on each unit interval with integral endpoints, we have only to show that

$$T_m(n+2) = T_m(n), \quad n = 0, 1, 2, \dots.$$

Consider solution (2.14) for $j = m$ on the intervals $[n, n+1]$ and $[n+2, n+3]$, that is,

$$T_m(t) = E_m(t-n)E_m^n(1)T_{0m}(0), \quad n \leq t \leq n+1$$

and

$$T_m(t) = E_m(t-n-2)E_m^{n+2}(1)T_{0m}(0), \quad n+2 \leq t \leq n+3.$$

At $t = n$ and $t = n+2$ these formulas respectively yield

$$T_m(n) = E_m^n(1)T_{0m}(0) \quad \text{and} \quad T_m(n+2) = E_m^{n+2}(1)T_{0m}(0).$$

Since hypothesis (2.30) implies $E_m(1) = 1$, then $T_m(n+2) = T_m(n)$, which proves the theorem for Eq. (2.8). The proof for Eq. (2.17) is analogous. Condition (2.31) is equivalent to $E_m(1/2) = -E_m(-1/2)$. It remains to note that the value of b in (2.30) is positive, and the value of b in (2.31) is negative.

3. A PARABOLIC EPCA OF NEUTRAL TYPE

Consider the boundary value problem (BVP) consisting of the equation

$$u_t(x, t) = a^2 u_{xx}(x, t) + bu_t(x, [t]) \quad (3.1)$$

and conditions (2.3) and (2.4). This equation is of neutral type since it includes the derivative u_t at different values of t .

Let $u_n(x, t)$ be the solution of the given problem on the interval $n \leq t \leq n+1$. Then $u_n(x, t)$ satisfies the equation

$$\frac{\partial u_n(x, t)}{\partial t} = a^2 \frac{\partial^2 u_n(x, t)}{\partial x^2} + b \frac{\partial^2 u_n}{\partial x^2}(x, n^+) \quad (3.2)$$

with boundary conditions (2.3) and the initial condition

$$u_n(x, n^+) = u_n(x),$$

where $u_n(x)$ is yet unknown. The solution of (3.2) is sought in form (2.5), and separation of variables generates BVP (2.6) with eigenfunctions (2.7) and the ODE

$$T'_{nj} = -a^2 \lambda_j^2 T_{nj}(t) + b T'_{nj}(n^+), \quad \lambda_j = \pi j. \quad (3.3)$$

Then the series

$$u_n(x, t) = \sum_{j=1}^{\infty} X_j(x) T_{nj}(t) \quad (3.4)$$

represents a formal solution of problem (3.2), (2.3). Turning to Eq. (3.3), its general solution on the interval $n \leq t < n+1$ is

$$T_{nj}(t) = C_{nj} e^{-a^2 \lambda_j^2 (t-n)} + \frac{b}{a^2 \lambda_j^2} T'_{nj}(n^+).$$

At $t = n^+$, Eq. (3.3) gives

$$T'_{nj}(n^+) = -a^2 \lambda_j^2 T_{nj}(n) + b T'_{nj}(n^+)$$

whence

$$T'_{nj}(n^+) = \frac{a^2 \lambda_j^2}{b-1} T_{nj}(n), \quad b \neq 1.$$

Therefore,

$$T_{nj}(t) = C_{nj} e^{a^2 \lambda_j^2 (t-n)} + \frac{b}{b-1} T_{nj}(n).$$

At $t = n$, we have

$$T_{nj}(n) = C_{nj} + \frac{b}{b-1} T_{nj}(n),$$

that is,

$$C_{nj} = \frac{1}{1-b} T_{nj}(n)$$

and

$$T_{nj}(t) = \frac{1}{b-1} \left(b - e^{-a^2 \lambda_j^2 (t-n)} \right) T_{nj}(n).$$

Denote

$$F_j(t) = \frac{b - e^{-a^2 \lambda_j^2 t}}{b-1}. \quad (3.5)$$

At $t = n + 1$, we have

$$T_{nj}(n+1) = F_j(1) T_{nj}(n)$$

and since, by hypothesis (i) of the above definition,

$$T_{nj}(n+1) = T_{n+1,j}(n+1),$$

then

$$T_{n+1,j}(n+1) = F_j(1) T_{nj}(n)$$

and

$$T_{nj}(n) = F_j^n(1) T_{0,j}(0). \quad (3.6)$$

Consequently,

$$T_{nj}(t) = F_j(t-n) F_j^n(1) T_{0,j}(0) \quad (3.7)$$

and

$$u_n(x, t) = \sum_{j=1}^{\infty} \sqrt{2} F_j^n(1) T_{0,j}(0) F_j(t-n) \sin(\pi j x). \quad (3.8)$$

The following theorems illustrate the far more complicated solution structure of Eq. (3.1) comparing with the diffusion equation without time delay.

THEOREM 3.1. For $b > 1$, each function $T_{nj}(t)$ is monotone unbounded as $t \rightarrow +\infty$.

PROOF. If $b > 1$, then (3.5) implies that $F_j(t) > 1$, for all $t > 0$. Hence, $F_j^n(1)$ grows exponentially as $t \rightarrow +\infty$, and the proof follows from (3.7).

THEOREM 3.2. For

$$\frac{1}{2} \left(1 + e^{-a^2 \lambda_1^2} \right) < b < 1, \quad (3.9)$$

each function $T_{nj}(t)$ is unbounded and oscillating.

PROOF. From inequalities (3.9) it follows that $F_j(1) < -1$, and therefore $F_j^{[t]}(1)$ is unbounded and has a zero in each unit interval with integer endpoints.

THEOREM 3.3. For $b < 0$, each function $T_{nj}(t)$ monotonically tends to zero as $t \rightarrow +\infty$.

PROOF. The inequalities

$$0 < F_j(1) < 1 \quad (3.10)$$

imply that solution (3.7) is asymptotically stable and nonoscillatory. From (3.10) it follows that

$$b < e^{-a^2 \lambda_j^2}. \quad (3.11)$$

Clearly the right-hand side of (3.11) vanishes as $j \rightarrow \infty$, and therefore the condition $b < 0$ is necessary and sufficient for all $T_{nj}(t)$ to approach zero monotonically.

THEOREM 3.4. If

$$a^2 \lambda_1^2 > \ln 2 \quad \text{and} \quad e^{-a^2 \lambda_1^2} < b < \frac{1}{2}, \quad (3.12)$$

then every solution (3.7) tends to zero and oscillates as $t \rightarrow +\infty$.

PROOF. From the inequalities

$$-1 < F_j(1) < 0, \quad (3.13)$$

which imply asymptotic stability and oscillatory behavior of solution (3.7), it follows that

$$e^{-a^2 \lambda_j^2} < b < \frac{1}{2} (1 + e^{-a^2 \lambda_j^2}).$$

These inequalities hold true for all j simultaneously with (3.12).

THEOREM 3.5. For any coefficient b that satisfies the inequalities

$$\frac{1}{2} < b < 1,$$

there exist infinitely many solutions $T_{nj}(t)$ which are unbounded and oscillating.

PROOF. For $b < 1$, the inequality $F_j(1) < -1$ which implies unboundedness and oscillation of $T_{nj}(t)$, is equivalent to

$$b > \frac{1}{2} + \frac{1}{2} e^{-a^2 \lambda_j^2}.$$

Clearly the right-hand side approaches $1/2$ as $j \rightarrow \infty$.

THEOREM 3.6. If $b < e^{-a^2 \lambda_m^2}$, then the solutions $T_1(t), \dots, T_m(t)$ of Eq. (3.3) monotonically tend to zero as $t \rightarrow +\infty$.

ACKNOWLEDGEMENT. This research was partially supported by NASA Grant NAG9-553

REFERENCES

- [1] AFTABIZADEH, A.R. and WIENER, J., Oscillatory properties of first order linear functional differential equations, *Applicable Anal.* **20** (1985), 165-187.
- [2] AFTABIZADEH, A.R., WIENER, J. and XU, J.-M., Oscillatory and periodic solutions of delay differential equations with piecewise constant argument, *Proc. Amer. Math. Soc.* **99** (1987), 673-679.
- [3] COOKE, K.L. and WIENER, J., Retarded differential equations with piecewise constant delays, *J. Math. Anal. App.* **99** (1984), 265-297.
- [4] COOKE, K.L. and WIENER, J., An equation alternately of retarded and advanced type, *Proc. Amer. Math. Soc.* **99** (1987), 726-732.
- [5] WIENER, J. and COOKE, K.L., Oscillations in systems of differential equations with piecewise constant argument, *J. Math. Anal. Appl.* **137** (1989), 221-239.
- [6] WIENER, J., Boundary-value problems for partial differential equations with piecewise constant delay, *Internat. J. Math. and Math. Sci.* **14** (1991), 301-321.
- [7] WIENER, J. and DEBNATH, L., Partial differential equations with piecewise constant delay, *Internat. J. Math. and Math. Sci.* **14** (1991), 485-496.
- [8] WIENER, J., *Generalized Solutions of Functional Differential Equations*, World Scientific, Singapore, 1993

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br