

UNIQUENESS OF WEAK SOLUTION FOR NONLINEAR ELLIPTIC EQUATIONS IN DIVERGENCE FORM

XU ZHANG

(Received 3 April 1997 and in revised form 21 June 1997)

ABSTRACT. We study the uniqueness of weak solutions for quasilinear elliptic equations in divergence form. Some counterexamples are given to show that our uniqueness result cannot be improved in the general case.

Keywords and phrases. Uniqueness, weak solution, quasilinear elliptic equation, divergence form, comparison theorem.

2000 Mathematics Subject Classification. Primary 35A05, 35B05, 35B50.

1. Introduction. In this paper, we demonstrate the uniqueness of weak solution of the Dirichlet problem for divergence structure elliptic equations of the form

$$L[u] \equiv - \sum_{i=1}^n \frac{\partial}{\partial x_i} (a_i(x, u, \nabla u)) + b(x, u) = 0 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0, \quad (1.1)$$

where Ω is a bounded domain of \mathbb{R}^n . In [1, 2, 3, 7, 8, 5], the uniqueness of classical solutions of problem (1.1) is treated under various hypotheses. Here, we consider the same problem for weak solutions. Especially, we give some counterexamples to show that our result cannot be improved in the general case.

To conclude this section, we would like to point out that after this paper had been submitted for publication, it came to our attention that a similar (uniqueness) result had been given in [4]. However, there is no further discussion in [4] as we do in Section 4.

2. Statement of the main results. Suppose that, for any $(x, z_1, z_2, \eta) \in \Omega \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}^n$,

$$\sum_{i,j} \frac{\partial(a_i(x, z, \eta))}{\partial \eta_j} \xi_i \xi_j \geq |\xi|^2, \quad \text{for all } \xi \in \mathbb{R}^n, \quad (2.1)$$

$$(b(x, z_1) - b(x, z_2))(z_1 - z_2) \geq 0, \quad (2.2)$$

$$|a_i(x, z_1, \eta) - a_i(x, z_2, \eta)| \leq |a(z_1, z_2)| |z_1 - z_2| (1 + |\eta|), \quad (2.3)$$

where $a \in L_{\text{loc}}^\infty(\mathbb{R} \times \mathbb{R})$.

REMARK 2.1. Unlike the previous works (cf. [1, 5, 7, 8] and so on), we do not assume that $b(\cdot, \cdot)$ is Lipschitz continuous in its second argument.

We need the following two definitions, which can be found in many references (cf. [3] and so on).

DEFINITION 2.2. $u \in H^1(\Omega) \cap L^\infty(\Omega)$ is said to satisfy $L[u] \geq (\leq)0$ in Ω in the weak sense if

$$\int_{\Omega} \left\{ \sum_{i=1}^n a_i(x, u, \nabla u) \frac{\partial w}{\partial x_i} + b(x, u) w \right\} dx \geq (\leq)0, \quad (2.4)$$

for any $w \in H_0^1(\Omega)$ such that $w \geq 0$ a.e. in Ω .

DEFINITION 2.3. $u, v \in H^1(\Omega)$ are said to satisfy $u \leq v$ on $\partial\Omega$ in the weak sense if $(u - v)^+ \equiv \max\{u - v, 0\} \in H_0^1(\Omega)$.

Now, we can state our main results.

THEOREM 2.4 (Comparison theorem). *Let the hypotheses (2.1), (2.2), and (2.3) hold, and let $u_1, u_2 \in H^1(\Omega) \cap L^\infty(\Omega)$ satisfy*

$$L[u_1] \geq 0, \quad L[u_2] \leq 0 \quad \text{in } \Omega, \quad u_2 \leq u_1 \quad \text{on } \partial\Omega \quad (2.5)$$

in the weak sense. Then we have $u_1 \geq u_2$ a.e. in Ω .

If, furthermore, $a \in L^\infty(\mathbb{R} \times \mathbb{R})$ and $u_1, u_2 \in H^1(\Omega)$ satisfy condition (2.5), then the same conclusion holds.

THEOREM 2.5 (Uniqueness Theorem). *Let the hypotheses (2.1), (2.2), (2.3), and (2.4) hold. Then the problem (1.1) admits at most one weak solution $u \in H_0^1(\Omega) \cap L^\infty(\Omega)$.*

If, furthermore, $a \in L^\infty(\mathbb{R} \times \mathbb{R})$, then (1.1) admits at most one weak solution $u \in H_0^1(\Omega)$.

3. Proof of the main results

PROOF OF THEOREM 2.4. Assume that $u_1, u_2 \in H^1(\Omega) \cap L^\infty(\Omega)$ satisfy condition (2.5) in the weak sense, that is

$$\int_{\Omega} \left\{ \sum_{i=1}^n a_i(x, u_1, \nabla u_1) \frac{\partial w}{\partial x_i} + b(x, u_1) w \right\} dx \geq 0 \quad (3.1)$$

and

$$\int_{\Omega} \left\{ \sum_{i=1}^n a_i(x, u_2, \nabla u_2) \frac{\partial w}{\partial x_i} + b(x, u_2) w \right\} dx \leq 0, \quad (3.2)$$

for any $w \in H_0^1(\Omega)$ such that $w \geq 0$, a.e. in Ω .

Put

$$\Omega_1 = \{x \in \Omega; u_1(x) < u_2(x)\}. \quad (3.3)$$

We assert that $|\Omega_1| = 0$ ($|\Omega_1|$ denotes the Lebesgue measure of Ω_1). In fact, for any $\varepsilon > 0$, we write

$$E_\varepsilon = \{x \in \Omega_1; u_2 - u_1 > \varepsilon\}, \quad v_\varepsilon = \min(\varepsilon, (u_2 - u_1)^+). \quad (3.4)$$

Note that $v_\varepsilon = 0$ and $\nabla v_\varepsilon = 0$, a.e. in $\Omega \setminus \Omega_1$, and that $v_\varepsilon = \varepsilon$ and $\nabla v_\varepsilon = 0$, a.e. in E_ε (see, e.g., [9]).

LEMMA 3.1. *Let $u \in H_0^1(\Omega)$. Then $|u|, u^+, \min(\varepsilon, u) \in H_0^1(\Omega)$ for any nonnegative constant ε .*

PROOF. Let $u_n \in C_0^\infty(\Omega)$ be such that $u_n \rightarrow u$ in $H_0^1(\Omega)$. Set

$$v_n = \left(u_n^2 + \frac{1}{n^2} \right)^{1/2} - \frac{1}{n}. \quad (3.5)$$

It is easy to see that $v_n \in C_0^\infty(\Omega)$. By a direct calculation, we can check that $\{v_n\}$ is a Cauchy sequence in $H_0^1(\Omega)$. Thus, there is a function $v \in H_0^1(\Omega)$ such that $v_n \rightarrow v$ in $H_0^1(\Omega)$. On the other hand, it is obvious that for an appropriate subsequence (still denoted by itself) $u_n \rightarrow u$, a.e. in Ω and, hence,

$$v_n \rightarrow |u|, \quad \text{a.e. in } \Omega. \quad (3.6)$$

So, we obtain that $|u| = v \in H_0^1(\Omega)$. Similarly, we can prove that $u^+, \min(\varepsilon, u) \in H_0^1(\Omega)$ for any nonnegative constant ε . This completes the proof of the lemma. \square

Now, we can return to the proof of Theorem 2.4. Since $u_1, u_2 \in H^1(\Omega)$ and $u_2 \leq u_1$ on $\partial\Omega$ in the weak sense, we have $(u_2 - u_1)^+ \in H_0^1(\Omega)$ by Definition 2.3 and $v_\varepsilon \in H_0^1(\Omega)$ by the above lemma. By (3.1) and (3.2), we then obtain

$$\begin{aligned} 0 &\geq \int_{\Omega} \sum_i (a_i(x, u_2, \nabla u_2) - a_i(x, u_1, \nabla u_1)) \frac{\partial w}{\partial x_i} dx \\ &\quad + \int_{\Omega} (b(x, u_2) - b(x, u_1)) w dx \\ &= \int_{\Omega} \sum_{i,j} \int_0^1 \frac{\partial a_i(x, u_2, \nabla u_1 + s(\nabla u_2 - \nabla u_1))}{\partial \eta_j} ds \frac{\partial (u_2 - u_1)}{\partial x_j} \frac{\partial w}{\partial x_i} dx \quad (3.7) \\ &\quad + \int_{\Omega} \sum_i (a_i(x, u_2, \nabla u_1) - a_i(x, u_1, \nabla u_1)) \frac{\partial w}{\partial x_i} dx \\ &\quad + \int_{\Omega} (b(x, u_2) - b(x, u_1)) w dx, \end{aligned}$$

for any $w \in H_0^1(\Omega)$ satisfying $w \geq 0$ a.e. in Ω . Take $w = v_\varepsilon$ in the above inequality. Note that $v_\varepsilon \geq 0$ and that $u_2 > u_1$ whenever $v_\varepsilon > 0$. By condition (2.2), we see that $b(x, u_2) - b(x, u_1) \geq 0$ whenever $v_\varepsilon > 0$. Thus,

$$\int_{\Omega} (b(x, u_2) - b(x, u_1)) v_\varepsilon dx \geq 0. \quad (3.8)$$

So, we have

$$\begin{aligned} 0 &\geq \int_{\Omega} \sum_{i,j} \int_0^1 \frac{\partial a_i(x, u_2, \nabla u_1 + s(\nabla u_2 - \nabla u_1))}{\partial \eta_j} ds \frac{\partial(u_2 - u_1)}{\partial x_j} \frac{\partial v_{\varepsilon}}{\partial x_i} dx \\ &\quad + \int_{\Omega} \sum_i (a_i(x, u_2, \nabla u_1) - a_i(x, u_1, \nabla u_1)) \frac{\partial v_{\varepsilon}}{\partial x_i} dx. \end{aligned} \quad (3.9)$$

Since $v_{\varepsilon} = u_2 - u_1$ a.e. in $\Omega_1 \setminus E_{\varepsilon}$ and $\nabla v_{\varepsilon} = 0$ a.e. in $\Omega \setminus \{\Omega_1 \setminus E_{\varepsilon}\}$, by the above inequality, we get

$$\begin{aligned} 0 &\geq \int_{\Omega_1 \setminus E_{\varepsilon}} \sum_{i,j} \int_0^1 \frac{\partial a_i(x, u_2, \nabla u_1 + s(\nabla u_2 - \nabla u_1))}{\partial \eta_j} ds \frac{\partial v_{\varepsilon}}{\partial x_j} \frac{\partial v_{\varepsilon}}{\partial x_i} dx \\ &\quad + \int_{\Omega_1 \setminus E_{\varepsilon}} \sum_i (a_i(x, u_2, \nabla u_1) - a_i(x, u_1, \nabla u_1)) \frac{\partial v_{\varepsilon}}{\partial x_i} dx \\ &\geq \int_{\Omega_1 \setminus E_{\varepsilon}} |\nabla v_{\varepsilon}|^2 dx + \int_{\Omega_1 \setminus E_{\varepsilon}} \sum_i (a_i(x, u_2, \nabla u_1) - a_i(x, u_1, \nabla u_1)) \frac{\partial v_{\varepsilon}}{\partial x_i} dx \\ &= \int_{\Omega} |\nabla v_{\varepsilon}|^2 dx + \int_{\Omega_1 \setminus E_{\varepsilon}} \sum_i (a_i(x, u_2, \nabla u_1) - a_i(x, u_1, \nabla u_1)) \frac{\partial v_{\varepsilon}}{\partial x_i} dx, \end{aligned} \quad (3.10)$$

where we have used condition (2.1) at the second step. By (2.3) and the above inequality, we get

$$\begin{aligned} \int_{\Omega} |\nabla v_{\varepsilon}|^2 dx &\leq \left| \int_{\Omega_1 \setminus E_{\varepsilon}} \sum_i (a_i(x, u_1, \nabla u_1) - a_i(x, u_2, \nabla u_1)) \frac{\partial v_{\varepsilon}}{\partial x_i} dx \right| \\ &\leq \varepsilon C_1 \left(\int_{\Omega_1 \setminus E_{\varepsilon}} (1 + |\nabla u_1|)^2 dx \right)^{1/2} \|\nabla v_{\varepsilon}\|_{L^2(\Omega)} \end{aligned} \quad (3.11)$$

for some constant C_1 (since u_1, u_2 are bounded). Thus,

$$\|\nabla v_{\varepsilon}\|_{L^2(\Omega)} \leq \varepsilon C_1 \left(\int_{\Omega_1 \setminus E_{\varepsilon}} (1 + |\nabla u_1|)^2 dx \right)^{1/2}. \quad (3.12)$$

But also, using the Sobolev-Poincaré inequality on Ω , we obtain

$$\begin{aligned} |E_{\varepsilon}| &= \varepsilon^{-2} \int_{E_{\varepsilon}} \varepsilon^2 dx \leq \varepsilon^{-2} \int_{\Omega} |\nabla v_{\varepsilon}|^2 dx \\ &\leq C \varepsilon^{-2} \int_{\Omega} |\nabla v_{\varepsilon}|^2 dx \leq C C_1^2 \int_{\Omega_1 \setminus E_{\varepsilon}} (1 + |\nabla u_1|)^2 dx. \end{aligned} \quad (3.13)$$

But $0 < u_2 - u_1 \leq \varepsilon$ in $\Omega_1 \setminus E_{\varepsilon}$. So, obviously,

$$\lim_{\varepsilon \rightarrow 0} (\Omega_1 \setminus E_{\varepsilon}) = \emptyset. \quad (3.14)$$

Thus, $|E_{\varepsilon}| \rightarrow 0$ as $\varepsilon \rightarrow 0$. On the other hand, clearly, $|E_{\varepsilon}|$ is nondecreasing. Thus, $|E_{\varepsilon}| = 0$ for any $\varepsilon > 0$ and in turn $u_2 \leq u_1$, a.e. in Ω .

We omit the details of the proof of the second statement since it is similar to that of the first part. \square

PROOF OF THEOREM 2.5. Assume that $u_1, u_2 \in H_0^1(\Omega) \cap L^\infty(\Omega)$ are two weak solutions of problem (1.1). By Theorem 2.4, we must have

$$u_1 \geq u_2, \quad u_2 \geq u_1, \quad \text{a.e. in } \Omega. \quad (3.15)$$

So, $u_1 = u_2$ a.e. in Ω , which proves the first conclusion.

We omit the details of the proof of the second statement since it is similar to that of the first part. \square

4. Some remarks and counterexamples

REMARK 4.1. Uniqueness does not hold for equations of nondivergence form,

$$-\sum_{i,j=1}^n a_{ij}(x, u, \nabla u) \frac{\partial^2 u}{\partial x_i \partial x_j} + b(x, u) = 0 \quad \text{in } \Omega, \quad (4.1)$$

even when $b \equiv 0$ and a_{ij} is independent of ∇u , as shown by Meyers [6]. In particular, he gave an example of a nondivergence equation with analytic coefficients, which is uniformly elliptic and which has nonunique analytic solutions in a bounded domain with analytic boundary.

REMARK 4.2. Condition (2.3) essentially says that $a_i(x, z, \eta)$ is locally Lipschitz continuous with respect to its second argument and, furthermore, that the Lipschitz constant is independent of x . This condition also cannot be removed. In fact, we have the following counterexamples.

(1) Consider the equation

$$-\sum \frac{\partial}{\partial x_i} \left(a(x, u) \delta_{ij} \frac{\partial u}{\partial x_j} \right) = 1 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0, \quad (4.2)$$

where $a(x, u) = 2 - \sin\{\pi \operatorname{sgn}(u - u_1)\}$ and where $u_1 \in H_0^1(\Omega)$ solves the problem

$$-2\Delta u = 1 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0. \quad (4.3)$$

It is easy to check that $u = u_1$ and $u = 2u_1$ are both solutions for problem (4.2), and that the minimum principle holds for (4.2) (for all of the following examples, maximum/minimum principle also holds). (Note that $a(x, u)$ is not continuous in its second argument.)

(2) In the previous example, we take

$$a(x, u) = \begin{cases} 2 - \sin(u - 2u_1) \sin((u - u_1)^{1/2}) & \text{if } u \geq u_1, \\ 2 & \text{if } u < u_1. \end{cases} \quad (4.4)$$

It is easy to check that both $u = u_1$ and $u = 2u_1$ are solutions of problem (4.2). (Note that $a(x, u)$ is not locally Lipschitz continuous in its second argument.)

(3) Take

$$a(x, u) = 2 - \sin \left\{ \frac{\pi}{2} \frac{u(u - u_1)}{u_1^2} \right\} \quad (4.5)$$

in the previous example. It is easy to check that $u = u_1$ and $u = 2u_1$ are again solutions of problem (4.2).

REMARK 4.3. If $b(x, u)$ is replaced by $b(x, u, \nabla u)$ in (1.1), we cannot obtain any satisfactory uniqueness result except for special cases such as $\partial b(\cdot, \cdot, \cdot)/\partial z \gg 0$ and where other restrictive conditions are assumed. In particular, consider the equation

$$\frac{d^2u}{dx^2} + \left| x \frac{du}{dx} \right| = 0 \quad \text{in } \Omega = (-1, 1), \quad u(-1) = u(1) = 0. \quad (4.6)$$

We can check directly that $u = 0$ and

$$u(x) = \int_x^1 e^{s^2/2} ds \chi_{[0,1]}(x) + \int_{-1}^x e^{s^2/2} ds \chi_{[-1,0)}(x) \quad (4.7)$$

are both weak solutions of this problem.

ACKNOWLEDGEMENTS. This work is partially supported by Chinese State Education Commission Science Foundation. The author gratefully acknowledges Professor James Serrin for his careful comments and useful help.

REFERENCES

- [1] J. Douglas, Jr., T. Dupont, and J. Serrin, *Uniqueness and comparison theorems for nonlinear elliptic equations in divergence form*, Arch. Rational Mech. Anal. **42** (1971), 157–168. MR 52 14637. Zbl 222.35017.
- [2] A. Friedman, *Uniqueness properties in the theory of differential operators of elliptic type*, J. Math. Mech. **7** (1958), 61–67. MR 20#174. Zbl 080.30702.
- [3] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Grundlehren der Mathematischen Wissenschaften, vol. 224, Springer-Verlag, Berlin-New York, 1977. MR 57 13109. Zbl 361.35003.
- [4] I. Hlaváček, M. Křížek, and J. Mály, *On Galerkin approximations of a quasilinear nonpotential elliptic problem of a nonmonotone type*, J. Math. Anal. Appl. **184** (1994), no. 1, 168–189. MR 95c:35102. Zbl 802.65113.
- [5] A. McNabb, *Strong comparison theorems for elliptic equations of second order*, J. Math. Mech. **10** (1961), 431–440. MR 26#448. Zbl 106.29903.
- [6] N. G. Meyers, *An example of non-uniqueness in the theory of quasi-linear elliptic equations of second order*, Arch. Rational Mech. Anal. **14** (1963), 177–179. MR 27#6021. Zbl 135.15501.
- [7] J. Serrin, *The problem of Dirichlet for quasilinear elliptic differential equations with many independent variables*, Philos. Trans. Roy. Soc. London Ser. A **264** (1969), 413–496. MR 43#7772. Zbl 181.38003.
- [8] ———, *On the strong maximum principle for quasilinear second order differential inequalities*, J. Functional Analysis **5** (1970), 184–193. MR 41#3966. Zbl 188.41701.
- [9] W. P. Ziemer, *Weakly differentiable functions*, Graduate Texts in Mathematics, vol. 120, Springer-Verlag, New York, 1989, Sobolev spaces and functions of bounded variation. MR 91e:46046. Zbl 692.46022.

ZHANG: INSTITUTE OF MATHEMATICS, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA
E-mail address: xuzhang@fudan.edu

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru