

COMBINATORICS OF GEOMETRICALLY DISTRIBUTED RANDOM VARIABLES: NEW q -TANGENT AND q -SECANT NUMBERS

HELMUT PRODINGER

(Received 1 November 1999)

ABSTRACT. Up-down permutations are counted by tangent (respectively, secant) numbers. Considering words instead, where the letters are produced by independent geometric distributions, there are several ways of introducing this concept; in the limit they all coincide with the classical version. In this way, we get some new q -tangent and q -secant functions. Some of them also have nice continued fraction expansions; in one particular case, we could not find a proof for it. Divisibility results à la Andrews, Foata, Gessel are also discussed.

Keywords and phrases. Tangent numbers, secant numbers, alternating permutations, geometric distribution, q -analogues.

2000 Mathematics Subject Classification. Primary 05A15, 05A30.

1. Introduction. Permutations $\pi = \pi_1 \pi_2 \cdots \pi_n$ are called up-down permutations if $\pi_1 < \pi_2 > \pi_3 < \pi_4 > \pi_5 \cdots$. For odd n , their number is given by $n![z^n] \tan z$, and for even n by $n![z^n] \sec z$ (cf. [8, 9]).

Instead of speaking about exponential generating functions, we prefer to think of the coefficients of $\tan z$ and $\sec z$ as probabilities.

This paper introduces several q -analogues of the functions $\tan z$ and $\sec z$. Some are classical (cf. [8, 12]), but the others seem to be new.

If we consider words $a_1 a_2 \cdots a_n$ with letters in $\{1, 2, \dots\}$ with probabilities (weights) p, pq, pq^2, \dots , where $p+q=1$ (independent geometric probabilities), then there are several ways to introduce this concept. We can use $<$ or \leq for “up,” $>$ or \geq for “down,” which gives four possibilities. Also, it makes a difference to consider “up-down” versus “down-up.” That gives in principle eight versions for q -tangent and q -secant numbers. However, reading the word from right to left, the instance “ $\leq > \leq > \cdots$ ” coincides with “ $< \geq < \geq < \cdots$,” and similarly for “ $\geq < \geq < \cdots$ ” and “ $> \leq > \leq \cdots$,” which gives us six q -tangent numbers (probabilities, to be more precise). In the instance of even length (secant numbers), there are more symmetries, and we have only four q -secant numbers.

By general principles, the limit $q \rightarrow 1$ reduces all the instances to the classical quantities.

Originally, the idea to use geometric probabilities for words came from some combinatorial problems in computer science (cf. [14]).

This paper aims by no means to offer a complete theory of these new numbers. Some results are given, others conjectured, and it is hoped that the papers stimulates other people to do more related research.

A few definitions from q -analysis are needed; see [1, 2]:

$$\begin{aligned} [n]_q &:= \frac{1-q^n}{1-q}, & [n]_q! &:= [1]_q [2]_q \cdots [n]_q, \\ (x;q)_n &:= (1-x)(1-xq)(1-xq^2) \cdots (1-xq^{n-1}). \end{aligned} \quad (1.1)$$

2. Recursions. The classical book [8] offers a general framework to deal with words and pattern (“pattern algebra”). However, we have decided to use a different approach that works particularly well in the present context, and is completely elementary. It is called the “adding a new slice” technique. In doing so, recursions are obtained; by iterating them, the generating functions of interest come out almost effortlessly.

We introduce the functions

$$T_n^{\leq\geq}(u), \quad (2.1)$$

where the coefficient of u^i in it is the probability that a word of length n satisfies the $\leq\geq\leq\geq \cdots$ condition and ends with the letter i . Also, we define

$$\tau_n^{\leq\geq} = T_n^{\leq\geq}(1), \quad (2.2)$$

which drops the technical condition about the last letter.

Furthermore, we introduce the generating functions

$$F^{\leq\geq}(z, u) = \sum_{n \geq 0} T_n^{\leq\geq}(u) z^n, \quad f^{\leq\geq}(z) = F^{\leq\geq}(z, 1). \quad (2.3)$$

Quantities like $F^{\leq>}(z, u)$ etc. are defined in an obvious way.

For the instance of secant numbers, similar quantities will be defined, but the letters S, σ, G, g are used instead of T, τ, F, f .

Obviously, there are only nonzero contributions for odd n in the tangent case and for even n in the secant case.

The reason to operate with a variable u that controls the last letter is the technique of “adding a new slice,” that was applied with success in [6] and, more recently, in [13].

THEOREM 2.1. *The functions $T_{2n+1}^{\nabla\Delta}(u)$ satisfy the following recurrences:*

$$\begin{aligned} T_{2n+1}^{\geq\leq}(u) &= \frac{p^2 u}{(1-qu)(1-q^2u)} T_{2n-1}^{\geq\leq}(1) - \frac{p^2 u}{(1-qu)(1-q^2u)} T_{2n-1}^{\geq\leq}(q^2u), \\ T_1^{\geq\leq}(u) &= \frac{pu}{1-qu}, \end{aligned} \quad (2.4)$$

$$\begin{aligned} T_{2n+1}^{\geq<}(u) &= \frac{p^2 qu^2}{(1-qu)(1-q^2u)} T_{2n-1}^{\geq<}(1) - \frac{p^2 qu^2}{(1-qu)(1-q^2u)} T_{2n-1}^{\geq<}(q^2u), \\ T_1^{\geq<}(u) &= \frac{pu}{1-qu}, \end{aligned} \quad (2.5)$$

$$\begin{aligned} T_{2n+1}^{><}(u) &= \frac{p^2 qu^2}{(1-qu)(1-q^2u)} T_{2n-1}^{><}(1) - \frac{p^2 u}{q(1-qu)(1-q^2u)} T_{2n-1}^{><}(q^2u), \\ T_1^{><}(u) &= \frac{pu}{1-qu}, \end{aligned} \quad (2.6)$$

$$\begin{aligned} T_{2n+1}^{\leq\geq}(u) &= \frac{pu}{q(1-qu)} T_{2n-1}^{\leq\geq}(q) - \frac{p^2 u}{q(1-qu)(1-q^2u)} T_{2n-1}^{\leq\geq}(q^2u), \\ T_1^{\leq\geq}(u) &= \frac{pu}{1-qu}, \end{aligned} \quad (2.7)$$

$$\begin{aligned} T_{2n+1}^{\leq>} (u) &= \frac{pu}{q(1-qu)} T_{2n-1}^{\leq>} (q) - \frac{p^2}{q^2(1-qu)(1-q^2u)} T_{2n-1}^{\leq>} (q^2u), \\ T_1^{\leq>} (u) &= \frac{pu}{1-qu}, \end{aligned} \quad (2.8)$$

$$\begin{aligned} T_{2n+1}^{<>} (u) &= \frac{pu}{1-qu} T_{2n-1}^{<>} (q) - \frac{p^2u}{(1-qu)(1-q^2u)} T_{2n-1}^{<>} (q^2u), \\ T_1^{<>} (u) &= \frac{pu}{1-qu}. \end{aligned} \quad (2.9)$$

PROOF. Since the technique is the same for all the instances, it is enough to discuss, for example, the “ $\geq\leq$ ” case. Adding a new slice means adding a pair (k, j) with $1 \leq k \leq i$, $j \geq k$, replacing u^i by 1 and providing the factor u^j . But

$$\sum_{k=1}^i pq^{k-1} \sum_{j \geq k} pq^{j-1} u^j = \frac{p^2u}{(1-qu)(1-q^2u)} - \frac{p^2u}{(1-qu)(1-q^2u)} (q^2u)^i, \quad (2.10)$$

which explains the recursion. The starting value is just

$$\sum_{j \geq 1} pq^{j-1} u^j = \frac{pu}{1-qu}. \quad (2.11)$$

(Readers who feel uncomfortable with this technique can write down a recursion for $P_{2n+1,j}$, the probability that a down-up composition of length $2n+1$ ends with j , namely,

$$P_{2n+1,j} = pq^{j-1} (1-q^j) \sum_{j \geq 1} P_{2n-1,j} - \sum_{k=1}^{j-1} pq^{j-k-1} (1-q^{j-k}) q^{2k} P_{2n-1,k}, \quad (2.12)$$

and translate it *afterwards* into (2.4).) \square

THEOREM 2.2. *The numbers $\tau_{2n+1}^{\nabla\Delta}$ have the generating functions $f^{\nabla\Delta}(z) = \tan_q(z) = \sin_q(z)/\cos_q(z)$:*

$$\begin{aligned} f^{\geq\leq}(z) &= \frac{\sum_{n \geq 0} ((-1)^n z^{2n+1}/[2n+1]_q!) q^{n(n+1)}}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!) q^{n(n-1)}}, \\ f^{\geq<}(z) &= \frac{\sum_{n \geq 0} ((-1)^n z^{2n+1}/[2n+1]_q!)}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!)}, \\ f^{><}(z) &= \frac{\sum_{n \geq 0} ((-1)^n z^{2n+1}/[2n+1]_q!) q^{n^2}}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!) q^{n^2}}, \\ f^{\leq\geq}(z) &= \frac{\sum_{n \geq 0} ((-1)^n z^{2n+1}/[2n+1]_q!) q^{n^2}}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!) q^{n(n-1)}}, \\ f^{\leq>}(z) &= \frac{\sum_{n \geq 0} ((-1)^n z^{2n+1}/[2n+1]_q!)}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!)}, \\ f^{<>}(z) &= \frac{\sum_{n \geq 0} ((-1)^n z^{2n+1}/[2n+1]_q!) q^{n(n+1)}}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!) q^{n^2}}. \end{aligned} \quad (2.13)$$

PROOF. The proofs of the first three relations are very similar, and we only sketch the first instance. Summing up we find

$$F^{\geq\leq}(z, u) = \frac{puz}{1-qu} + \frac{p^2uz^2}{(1-qu)(1-q^2u)} F^{\geq\leq}(z, 1) - \frac{p^2uz^2}{(1-qu)(1-q^2u)} F^{\geq\leq}(z, q^2u). \quad (2.14)$$

Iterating that we find for $f(z) = f^{\geq\leq}(z)$:

$$\begin{aligned} f(z) &= \frac{pz}{1-q} + \frac{p^2z^2}{(1-q)(1-q^2)} f(z) - \frac{p^2q^2z^3}{(1-q)(1-q^2)(1-q^3)} \\ &\quad - \frac{p^4q^2z^4}{(1-q)(1-q^2)(1-q^3)(1-q^4)} f(z) + \dots \end{aligned} \quad (2.15)$$

from which the announced formula follows by solving for $f(z)$.

The three others are trickier, because of a term $T_{2n-1}^{\nabla\Delta}(q)$. Again, one case is discussed. Observe that

$$T_{2n-1}^{\leq\geq}(q) = q^{-1} S_{2n}^{\leq\geq}(1), \quad (2.16)$$

because one more “up” step should replace u^i by $\sum_{k\geq i} pq^{k-1} = q^{i-1}$. Now the generating function $g^{\leq\geq}(z)$ of the quantities $S_{2n}^{\leq\geq}(1)$ (upcoming) is obtained independently, whence we get

$$F^{\leq\geq}(z, u) = \frac{puz}{q^2(1-qz)} g^{\leq\geq}(z) - \frac{p^2uz^2}{q(1-qu)(1-q^2u)} F^{\leq\geq}(z, q^2u). \quad (2.17)$$

Now iteration as usual derives the desired result. \square

THEOREM 2.3. *The functions $S_{2n}^{\nabla\Delta}(u)$ satisfy the following recurrences:*

$$\begin{aligned} S_{2n+2}^{\leq\geq}(u) &= \frac{p^2u}{(1-qu)(1-q^2u)} S_{2n}^{\leq\geq}(1) - \frac{p^2u}{(1-qu)(1-q^2u)} S_{2n}^{\leq\geq}(q^2u), \\ S_2^{\leq\geq}(u) &= \frac{p^2u}{(1-qu)(1-q^2u)}, \end{aligned} \quad (2.18)$$

$$\begin{aligned} S_{2n+2}^{\leq>}(u) &= \frac{p^2u}{(1-qu)(1-q^2u)} S_{2n}^{\leq>}(1) - \frac{p^2}{q^2(1-qu)(1-q^2u)} S_{2n}^{\leq>}(q^2u), \\ S_2^{\leq>}(u) &= \frac{p^2u}{(1-qu)(1-q^2u)}, \end{aligned} \quad (2.19)$$

$$\begin{aligned} S_{2n+2}^{\geq\geq}(u) &= \frac{p^2qu^2}{(1-qu)(1-q^2u)} S_{2n}^{\geq\geq}(1) - \frac{p^2qu^2}{(1-qu)(1-q^2u)} S_{2n}^{\geq\geq}(q^2u), \\ S_2^{\geq\geq}(u) &= \frac{p^2qu^2}{(1-qu)(1-q^2u)}, \end{aligned} \quad (2.20)$$

$$\begin{aligned} S_{2n+2}^{<>} (u) &= \frac{p^2 qu^2}{(1-qu)(1-q^2u)} S_{2n}^{<>} (1) - \frac{p^2 u}{q(1-qu)(1-q^2u)} S_{2n}^{<>} (q^2 u), \\ S_2^{<>} (u) &= \frac{p^2 qu^2}{(1-qu)(1-q^2u)}. \end{aligned} \quad (2.21)$$

PROOF. The proof works as in the easy cases of the tangent recursions and is omitted. For the starting value, the first pair of numbers must be considered. \square

THEOREM 2.4. *The numbers $\sigma_{2n}^{\nabla\Delta}$ have the generating functions $g^{\nabla\Delta}(z) = 1/\cos_q(z)$:*

$$\begin{aligned} g^{\leq\geq}(z) &= \frac{1}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!) q^{n(n-1)}}, \\ g^{\leq>}(z) &= \frac{1}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!)}, \\ g^{<\geq}(z) &= \frac{1}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!) q^{n(2n-1)}}, \\ g^{<>}(z) &= \frac{1}{\sum_{n \geq 0} ((-1)^n z^{2n}/[2n]_q!) q^{n^2}}. \end{aligned} \quad (2.22)$$

PROOF. The proofs are quite similar as before; however, iteration must be done for the function $G^{\nabla\Delta}(z, u) - 1$, and 1 must be added at the end. \square

3. Jackson's *q*-sine and *q*-cosine functions. In this section, we are considering a general class of *q*-sine and *q*-cosine functions and sort out those that satisfy a natural condition. This condition is even more natural, as all the previously encountered *q*-sine and *q*-cosine functions satisfy them.

Jackson in [12] has introduced the functions

$$\sin_q(z) = \sum_{n \geq 0} \frac{(-1)^n z^{2n+1}}{[2n+1]_q!}, \quad \cos_q(z) = \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n]_q!} \quad (3.1)$$

and proved the relation

$$\sin_q(z) \sin_{1/q}(z) + \cos_q(z) \cos_{1/q}(z) = 1. \quad (3.2)$$

Since we have here several *q*-sine and *q*-cosine functions, we call them a *q*-sine-cosine pair, if relation (3.2) holds.

THEOREM 3.1. *For the functions*

$$\sin_q(z) := \sum_{n \geq 0} \frac{(-1)^n z^{2n+1}}{[2n+1]_q!} q^{An^2+Bn}, \quad \cos_q(z) := \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n]_q!} q^{Cn^2+Dn} \quad (3.3)$$

*exactly the twelve pairs in Table 3.1 are *q*-sine-cosine pairs:*

TABLE 3.1.

A	B	C	D
0	0	0	0
2	1	0	0
0	0	2	-1
2	1	2	-1
0	1	0	1
2	0	0	1
1	0	1	0
1	1	1	0
0	1	2	-2
2	0	2	-2
1	0	1	-1
1	1	1	-1

PROOF. The desired relation gives more and more restrictions for the (complex) numbers A, B, C, D when we look at the coefficients of z^{2n} for $n = 0, 1, 2, \dots$. By a tedious search that will not be reported here we find these twelve possibilities, and all others can be excluded. The proof that this indeed works is very similar for all of them, so we give just one, namely the instance $(1, 0, 1, 0)$.

Note the following expansions:

$$\sin_{1/q} z = \sum_{n \geq 0} \frac{(-1)^n z^{2n+1}}{[2n+1]_q!} q^{\binom{2n+1}{2} - n^2}, \quad \cos_{1/q} z = \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n]_q!} q^{\binom{2n}{2} - n^2}. \quad (3.4)$$

So we must prove that, for $n \geq 1$,

$$\sum_{k=0}^n \begin{bmatrix} 2n \\ 2k \end{bmatrix}_q q^{\binom{2k}{2} - k^2 + (n-k)^2} = \sum_{k=0}^{n-1} \begin{bmatrix} 2n \\ 2k+1 \end{bmatrix}_q q^{\binom{2k+1}{2} - k^2 + (n-k-1)^2} \quad (3.5)$$

or, reversing the order of summation in the second sum,

$$\sum_{k=0}^n \begin{bmatrix} 2n \\ 2k \end{bmatrix}_q q^{2k^2 - k - 2nk} = \sum_{k=0}^{n-1} \begin{bmatrix} 2n \\ 2k+1 \end{bmatrix}_q q^{2k^2 + k - n - 2nk}. \quad (3.6)$$

We rewrite this again as

$$\sum_{k \text{ even}} \begin{bmatrix} 2n \\ k \end{bmatrix}_q q^{-nk + \binom{k}{2}} = \sum_{k \text{ odd}} \begin{bmatrix} 2n \\ k \end{bmatrix}_q q^{-nk + \binom{k}{2}}. \quad (3.7)$$

Therefore, we have to prove that

$$\sum_{k=0}^{2n} \begin{bmatrix} 2n \\ k \end{bmatrix}_q (-1)^k q^{-nk + \binom{k}{2}} = 0. \quad (3.8)$$

We use the formula (10.0.9) in [2], see also [1],

$$\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q z^k q^{\binom{k}{2}} = \prod_{j=0}^{n-1} (1 + q^j z). \quad (3.9)$$

The desired result now follows by replacing n by $2n$ and plugging in $z = -q^{-n}$. \square

THEOREM 3.2. *The six $\tan_q(z)$ functions in Theorem 2.2 all involve q -sine-cosine pairs.*

REMARK 3.3. Replacing q by $1/q$ in the q -sine-cosine pairs and rewriting everything again in the q -notation means replacing the vector (A, B, C, D) of exponents by $(2 - A, 1 - B, 2 - C, -1 - D)$. This will be called “duality.”

TABLE 3.2.

A	B	C	D	A'	B'	C'	D'
0	0	0	0	2	1	2	-1
2	1	0	0	0	0	2	-1
0	1	0	1	2	0	2	-2
2	0	0	1	0	1	2	-2
1	0	1	0	1	1	1	-1
1	1	1	0	1	0	1	-1

This reduces the twelve pairs to six pairs.

4. Continued fractions. Experimenting with Maple, it was found that some of the twelve tangent functions have nice continued fraction expansions. Some could be proved, others not (yet).

A lot of work still needs to be done concerning a *combinatorial theory* of these continued fractions, perhaps in the style of [5].

THEOREM 4.1. *For $(A, B, C, D) = (0, 0, 0, 0)$ and $(A, B, C, D) = (2, 1, 2, -1)$ we have*

$$\begin{aligned} \tan_q(z) = & \frac{z}{z^2} \\ & [1]_q q^{-0} - \frac{z^2}{z^2} \\ & [3]_q q^{-1} - \frac{z^2}{z^2} \\ & [5]_q q^{-2} - \frac{z^2}{z^2} \\ & [7]_q q^{-3} - \frac{z^2}{\ddots} \end{aligned} \quad (4.1)$$

The two tangent functions coincide, which is classical, since Jackson [12] has shown that for his functions

$$\sin_q z \cos_{1/q} z - \sin_{1/q} z \cos_q z = 0 \quad (4.2)$$

holds.

PROOF. For the proof by induction we must do the following: set $a_n = [2n-1]_q q^{1-n}$ and

$$\begin{aligned} p_n(z) &= a_n p_{n-1}(z) - z^2 p_{n-2}(z), & p_0(z) &= 0, & p_1(z) &= z, \\ q_n(z) &= a_n q_{n-1}(z) - z^2 q_{n-2}(z), & q_0(z) &= 1, & q_1(z) &= a_1. \end{aligned} \quad (4.3)$$

We must show that

$$[z^k](p_n(z) \cos_q z - q_n(z) \sin_q z) = 0 \quad \text{for } k \leq 2n. \quad (4.4)$$

Now look at

$$[z^k]((a_n p_{n-1}(z) - z^2 p_{n-2}(z)) \cos_q z - (a_n q_{n-1}(z) - z^2 q_{n-2}(z)) \sin_q z). \quad (4.5)$$

By the induction hypothesis we only have to show that

$$[z^{2n-1}](p_n(z) \cos_q z - q_n(z) \sin_q z) = 0. \quad (4.6)$$

However, we can easily show by induction that

$$\begin{aligned} p_n(z) &= \sum_k z^{2k+1} (-1)^k \frac{[2n-2k-1]_q! q^{k(2k+1)-\binom{n}{2}}}{[n-2k-1]_q! [2k+1]_q! \prod_{i=1}^{n-1-2k} (1+q^i)}, \\ q_n(z) &= \sum_k z^{2k} (-1)^k \frac{[2n-2k]_q! q^{k(2k-1)-\binom{n}{2}}}{[n-2k]_q! [2k]_q! \prod_{i=1}^{n-2k} (1+q^i)}, \end{aligned} \quad (4.7)$$

holds (the hard part is to find these formulae). We have to prove that

$$\sum_{k \geq 0} [z^{2k+1}] p_n(z) [z^{2n-2k-2}] \cos_q z = \sum_{k \geq 0} [z^{2k}] q_n(z) [z^{2n-2k-1}] \sin_q z \quad (4.8)$$

or

$$\begin{aligned} &\sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \frac{[2n-2k-1]_q! q^{k(2k+1)}}{[n-2k-1]_q! [2k+1]_q! \prod_{i=1}^{n-1-2k} (1+q^i) [2n-2k-2]_q!} \\ &= \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{[2n-2k]_q! q^{k(2k-1)}}{[n-2k]_q! [2k]_q! \prod_{i=1}^{n-2k} (1+q^i) [2n-2k-1]_q!}. \end{aligned} \quad (4.9)$$

Thus we must prove

$$\sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \frac{(1-q^{2n-2k-1}) q^{k(2k+1)}}{[n-2k-1]_q! [2k+1]_q! \prod_{i=1}^{n-1-2k} (1+q^i)} = \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(1-q^{2n-2k}) q^{k(2k-1)}}{[n-2k]_q! [2k]_q! \prod_{i=1}^{n-2k} (1+q^i)} \quad (4.10)$$

or

$$\sum_{k=0}^n \frac{(1-q^{2n-k}) q^{\binom{k}{2}} (-1)^k}{[n-k]_q! [k]_q! \prod_{i=1}^{n-k} (1+q^i)} = 0, \quad (4.11)$$

or

$$\sum_{k=0}^n \frac{1}{(q;q)_k (q^2;q^2)_{n-k}} (1-q^{2n-k}) q^{\binom{k}{2}} (-1)^k = 0. \quad (4.12)$$

Now

$$\begin{aligned} \sum_{k=0}^n \frac{1}{(q;q)_k (q^2;q^2)_{n-k}} q^{\binom{k}{2}} (-1)^k &= [z^n] \sum_{k \geq 0} \frac{q^{\binom{k}{2}} (-1)^k z^k}{(q;q)_k} \sum_{k \geq 0} \frac{z^k}{(q^2;q^2)_k} \\ &= \frac{[z^n] \prod_{k \geq 0} (1-zq^k)}{\prod_{k \geq 0} (1-zq^{2k})} = [z^n] \prod_{k \geq 0} (1-zq^{2k+1}) \\ &= q^n [z^n] \prod_{k \geq 0} (1-zq^{2k}) = \frac{q^n q^{2\binom{n}{2}} (-1)^n}{(q^2;q^2)_n} = \frac{(-1)^n q^{n^2}}{(q^2;q^2)_n}. \end{aligned} \quad (4.13)$$

Similarly,

$$\begin{aligned} \sum_{k=0}^n \frac{q^{2n-k}}{(q;q)_k (q^2;q^2)_{n-k}} q^{\binom{k}{2}} (-1)^k &= q^{2n} [z^n] \sum_{k \geq 0} \frac{q^{\binom{k}{2}} (-1)^k (z/q)^k}{(q;q)_k} \sum_{k \geq 0} \frac{z^k}{(q^2;q^2)_k} \\ &= \frac{q^{2n} [z^n] \prod_{k \geq 0} (1-zq^{k-1})}{\prod_{k \geq 0} (1-zq^{2k})} = q^{2n} [z^n] \prod_{k \geq 0} (1-zq^{2k-1}) \\ &= q^{2n} q^{-n} [z^n] \prod_{k \geq 0} (1-zq^{2k}) = \frac{q^n q^{2\binom{n}{2}} (-1)^n}{(q^2;q^2)_n} = \frac{(-1)^n q^{n^2}}{(q^2;q^2)_n}. \end{aligned} \quad (4.14)$$

This finishes the proof.

The continued fraction for $(2, 1, 2, -1)$ follows by replacing q by $1/q$. \square

THEOREM 4.2. *For $(A, B, C, D) = (0, 1, 0, 1)$ we have*

$$\begin{aligned} \tan_q(z) &= \frac{z}{[1]_q q^{-0} - \frac{z^2}{[3]_q q^{-2} - \frac{z^2}{[5]_q q^{-2} - \frac{z^2}{[7]_q q^{-4} - \frac{z^2}{\ddots}}}}. \end{aligned} \quad (4.15)$$

The negative powers of q go like $0, 2, 2, 4, 4, 6, 6, 8, 8, \dots$

PROOF. The proof follows the same lines; this time the polynomials (continuants) are

$$\begin{aligned} p_n(z) &= \sum_k z^{2k+1} (-1)^k \frac{[2n-2k-1]_q! q^{2k(k+1)-\binom{n}{2}-[n/2]}}{[n-2k-1]_q! [2k+1]_q! \prod_{i=1}^{n-1-2k} (1+q^i)}, \\ q_n(z) &= \sum_k z^{2k} (-1)^k \frac{[2n-2k]_q! q^{2k^2-\binom{n}{2}-[n/2]}}{[n-2k]_q! [2k]_q! \prod_{i=1}^{n-2k} (1+q^i)}. \end{aligned} \quad (4.16)$$

Hence we have to prove that

$$\sum_{k=0}^{[(n-1)/2]} \frac{(1-q^{2n-2k-1})q^{2k(k+1)-k}}{[n-2k-1]_q![2k+1]_q!\prod_{i=1}^{n-1-2k} (1+q^i)} = \sum_{k=0}^{[n/2]} \frac{(1-q^{2n-2k})q^{2k^2-k}}{[n-2k]_q![2k]_q!\prod_{i=1}^{n-2k} (1+q^i)}; \quad (4.17)$$

from here on we can use the previous proof.

An alternative proof is by noting that

$$\tan_q^{(0,1,0,1)}(z) = \frac{1}{\sqrt{q}} \tan_q^{(0,0,0,0)}(z\sqrt{q}) \quad (4.18)$$

and using the previous result. \square

THEOREM 4.3. *For $(A, B, C, D) = (2, 0, 2, -2)$ we have*

$$\tan_q(z) = \frac{z}{[1]_q q^{-0} - \frac{z^2}{[3]_q q^{-0} - \frac{z^2}{[5]_q q^{-2} - \frac{z^2}{[7]_q q^{-2} - \frac{z^2}{\ddots}}}}}. \quad (4.19)$$

The negative powers of q go like $0, 0, 2, 2, 4, 4, 6, 6, 8, 8, \dots$

PROOF. This follows from the previous theorem by replacing q by $1/q$. \square

CONJECTURE 4.4. *For $(A, B, C, D) = (1, 0, 1, 0)$ we have*

$$\tan_q(z) = \frac{z}{[1]_q q^0 - \frac{z^2}{[3]_q q^{-2} - \frac{z^2}{[5]_q q^1 - \frac{z^2}{[7]_q q^{-9} - \frac{z^2}{\ddots}}}}}. \quad (4.20)$$

The positive powers of q go like $0, 1, 6, 15, \dots (k(2k-1))$.

The negative powers of q go like $2, 9, 20, 35, \dots ((k+1)(2k-1))$.

COMMENT. It might be useful to rewrite the continued fraction as

$$\frac{z}{1 - \frac{z^2 b_1}{1 - \frac{z^2 b_2}{1 - \frac{z^2 b_3}{1 - \frac{z^2 b_4}{\ddots}}}}}. \quad (4.21)$$

with

$$\begin{aligned} b_k &= \frac{1}{[k]_q[k+1]_q} q^{-k+(-1)^k(2k-1)} \\ &= \frac{1}{[k]_q[k+1]_q} \left[\frac{1}{2} q^{-3k+1} (1+q^{4k-2}) - \frac{(-1)^k}{2} q^{-3k+1} (1-q^{4k-2}) \right]. \end{aligned} \quad (4.22)$$

The recursions for the continuants are now

$$\begin{aligned} p_n(z) &= p_{n-1}(z) - b_{n-1} z^2 p_{n-2}(z), & p_0(z) &= 0, & p_1(z) &= z, \\ q_n(z) &= q_{n-1}(z) - b_{n-1} z^2 q_{n-2}(z), & q_0(z) &= 1, & q_1(z) &= 1. \end{aligned} \quad (4.23)$$

Unfortunately, even with this form, I am currently unable to guess the coefficients of these polynomials, whence I must leave this expansion as an open problem.

NOTE. Added in proof (September 2000): Markus Fulmek has established that recently.

CONJECTURE 4.5. *For $(A, B, C, D) = (1, 1, 1, -1)$ we have*

$$\begin{aligned} \tan_q(z) &= \frac{z}{[1]_q q^{-0} - \frac{z^2}{[3]_q q^0 - \frac{z^2}{[5]_q q^{-5} - \frac{z^2}{[7]_q q^3 - \frac{z^2}{\ddots}}}}} \end{aligned} \quad (4.24)$$

The positive powers of q go like $0, 3, 10, 21, \dots$ ($(k-1)(2k-1)$).

The negative powers of q go like $0, 5, 14, 27, \dots$ ($(k-1)(2k+1)$).

COMMENT. This would be a corollary of the previous expansion.

REMARK 4.6. Normally, as for example in [5, 10], the continued fraction expansions of the ordinary generating function of the tangent and secant numbers are considered, whereas we stick here to the exponential (or probability) generating functions.

5. Divisibility

THEOREM 5.1. *The coefficient*

$$[2n+1]_q [z^{2n+1}] \tan_q(z) \quad (5.1)$$

is divisible by

$$(1+q)(1+q^2) \cdots (1+q^n) \quad (5.2)$$

for the vectors of exponents $(0, 0, 0, 0)$, $(2, 1, 2, -1)$, $(0, 1, 0, 1)$, $(2, 0, 2, -2)$, $(1, 0, 1, 0)$, and $(1, 1, 1, -1)$.

PROOF. The proof of [4] covers the first four instances, since we note that

$$\tan_q^{(0,1,0,1)}(z) = \frac{1}{\sqrt{q}} \tan_q^{(0,0,0,0)}(z\sqrt{q}). \quad (5.3)$$

The only open case is thus $(1, 0, 1, 0)$, as the remaining one would follow from duality. Thus, let us now consider

$$\sin_q z = \sum_{n \geq 0} \frac{(-1)^n z^{2n+1}}{[2n+1]_q!} q^{n^2}, \quad \cos_q z = \sum_{n \geq 0} \frac{(-1)^n z^{2n}}{[2n]_q!} q^{n^2}, \quad (5.4)$$

and $\tan_q z = \sin_q z / \cos_q z$.

We need the following computation that is akin to the one in Theorem 3.1,

$$\begin{aligned} [z^{2n+1}] \sin_{1/q} z \cos_q z &= \sum_{k=0}^n \frac{(-1)^k q^{k(k+1)}}{[2k+1]_q!} \frac{(-1)^{n-k} q^{(n-k)^2}}{[2n-2k]_q!} \\ &= \frac{q^{n^2} (-1)^n}{[2n+1]_q!} \sum_{k=0}^n \begin{bmatrix} 2n+1 \\ 2k+1 \end{bmatrix}_q q^{2k^2+k-2nk} \\ &= \frac{q^{n(n+1)} (-1)^n}{[2n+1]_q!} \sum_{k \text{ odd}} \begin{bmatrix} 2n+1 \\ k \end{bmatrix}_q q^{\binom{k}{2}-nk} \\ &= \frac{q^{n(n+1)} (-1)^n}{[2n+1]_q!} \frac{1}{2} \sum_{k=0}^{2n+1} \begin{bmatrix} 2n+1 \\ k \end{bmatrix}_q q^{\binom{k}{2}-nk} \\ &= \frac{q^{n(n+1)} (-1)^n}{[2n+1]_q!} \frac{1}{2} \prod_{j=0}^{2n} (1 + q^j z) \Big|_{z=q^{-n}} \\ &= \frac{q^{\binom{n+1}{2}} (-1)^n}{[2n+1]_q!} \prod_{i=1}^n (1 + q^i)^2. \end{aligned} \quad (5.5)$$

We also mention the dual formula

$$[z^{2n+1}] \sin_q z \cos_{1/q} z = \frac{q^{\binom{n}{2}} (-1)^n}{[2n+1]_q!} \prod_{i=1}^n (1 + q^i)^2. \quad (5.6)$$

A similar computation gives the result ($n \geq 1$),

$$[z^{2n}] \cos_q z \cos_{1/q} z = -[z^{2n}] \sin_q z \sin_{1/q} z = \frac{q^{\binom{n}{2}} (-1)^n}{[2n]_q!} \prod_{i=1}^{n-1} (1 + q^i)^2 (1 + q^n). \quad (5.7)$$

Now we write $\tan_q z = \sin_q z \cos_{1/q} z / \cos_q z \cos_{1/q} z$ and thus

$$\cos_q z \cos_{1/q} z \sum_{n \geq 0} \frac{T_{2n+1}(q)}{[2n+1]_q!} z^{2n+1} = \sin_q z \cos_{1/q} z. \quad (5.8)$$

Comparing coefficients, we find

$$T_{2n+1}(q) + \sum_{k=1}^n \begin{bmatrix} 2n+1 \\ 2k \end{bmatrix}_q q^{\binom{k}{2}} (-1)^k \prod_{i=1}^{k-1} (1 + q^i)^2 (1 + q^k) T_{2n+1-2k}(q) = q^{\binom{n}{2}} (-1)^n \prod_{i=1}^n (1 + q^i)^2. \quad (5.9)$$

The induction argument is as in [4]; $T_{2n+1-2k}(q)$ has a factor $\prod_{i=1}^{n-k} (1+q^i)$ and, according again to [4],

$$\begin{bmatrix} 2n+1 \\ 2k \end{bmatrix}_q \frac{\prod_{i=1}^k (1+q^i)}{\prod_{i=n-k+1}^n (1+q^i)} \quad (5.10)$$

is still a polynomial.

The two factors $\prod_{i=n-k+1}^n (1+q^i)$ and $\prod_{i=1}^{n-k} (1+q^i)$ mean that everything in (5.9) must be divisible by $\prod_{i=1}^n (1+q^i)$, and this finishes the proof.

It is likely that stronger results as in [7] hold, but we have not investigated that. \square

The new q -secant numbers do not enjoy any divisibility results that are worthwhile to report; for the classical ones (see [3]).

REMARK 5.2. The paper [11] has a q -exponential function

$$\mathcal{E}_q := \sum_{n \geq 0} \frac{q^{n^2/4} z^n}{(q;q)_n}. \quad (5.11)$$

Plugging in $iz(1-q)$ for z and taking real parts would result in the q -cosine with factor q^{n^2} . To get the corresponding q -sine, replace z by $izq(1-q)$, take the imaginary part and multiply by $q^{1/4}$. We consider that merely to be a curiosity, not being of much help.

ACKNOWLEDGEMENT. Parts of this research were conducted while the author was a visitor of the Technical University of Graz where he was supported by the start project Y96-MAT.

I want to thank Dominique Foata for several pointers to the literature and an anonymous referee for valuable hints.

REFERENCES

- [1] G. E. Andrews, *The Theory of Partitions*, Encyclopedia of Mathematics and its Applications, vol. 2, Addison-Wesley Publishing Co., Reading, Mass., London, Amsterdam, 1976. MR 58#27738. Zbl 371.10001.
- [2] G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, Cambridge, 1999. MR 2000g:33001. Zbl 920.33001.
- [3] G. E. Andrews and D. Foata, *Congruences for the q -secant numbers*, European J. Combin. **1** (1980), no. 4, 283–287. MR 82d:05018. Zbl 455.10006.
- [4] G. E. Andrews and I. Gessel, *Divisibility properties of the q -tangent numbers*, Proc. Amer. Math. Soc. **68** (1978), no. 3, 380–384. MR 57#2925. Zbl 401.10020.
- [5] P. Flajolet, *Combinatorial aspects of continued fractions*, Discrete Applied Mathematics **39** (1992), 207–229.
- [6] P. Flajolet and H. Prodinger, *Level number sequences for trees*, Discrete Math. **65** (1987), no. 2, 149–156. MR 88e:05030. Zbl 634.05021.
- [7] D. Foata, *Further divisibility properties of the q -tangent numbers*, Proc. Amer. Math. Soc. **81** (1981), no. 1, 143–148. MR 81k:05005. Zbl 485.05007.
- [8] I. P. Goulden and D. M. Jackson, *Combinatorial Enumeration*, with a foreword by Gian-Carlo Rota. A Wiley-Interscience Publication. Wiley-Interscience Series in Discrete Mathematics, John Wiley & Sons Inc., New York, 1983. MR 84m:05002. Zbl 519.05001.

- [9] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics*, A foundation for computer science, 2nd ed., Addison-Wesley Publishing Company, Reading, MA, 1994. MR 97d:68003. Zbl 836.00001.
- [10] G.-N. Han, A. Randrianarivony, and J. Zeng, *Un autre q -analogue des nombres d'Euler [Another q -analogue of the Euler numbers]* *The Andrews Festschrift (Maratea, 1998)*, Sémin. Lothar. Combin. **42** (1999), Art. B42e, 22 pp. (electronic). MR 2000g:11013. Zbl 990.28747.
- [11] M. E. H. Ismail, M. Rahman, and D. Stanton, *Quadratic q -exponentials and connection coefficient problems*, Proc. Amer. Math. Soc. **127** (1999), no. 10, 2931–2941. MR 2000a:33027. Zbl 934.33025.
- [12] F. H. Jackson, *A basic-sine and cosine with symbolic solutions of certain differential equations*, Proc. Edinburg Math. Soc. **22** (1904), 28–39.
- [13] A. Knopfmacher and H. Prodinger, *On Carlitz compositions*, European J. Combin. **19** (1998), no. 5, 579–589. MR 99j:05006. Zbl 902.05004.
- [14] H. Prodinger, *Combinatorics of geometrically distributed random variables: left-to-right maxima*. Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics (Florence, 1993), Discrete Math. **153** (1996), no. 1-3, 253–270. MR 97c:05012. Zbl 853.60006.

HELMUT PRODINGER: THE JOHN KNOPFMACHER CENTRE FOR APPLICABLE ANALYSIS AND NUMBER THEORY, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, P.O. WITS, 2050 JOHANNESBURG, SOUTH AFRICA

E-mail address: helmut@gauss.cam.wits.ac.za

URL: <http://www.wits.ac.za/helmut/index.htm>

Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

Lead Guest Editor

Juan J. Nieto, Departamento de Análisis Matemático,
Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;
juanjose.nieto.roig@usc.es

Guest Editor

Donal O'Regan, Department of Mathematics, National
University of Ireland, Galway, Ireland;
donal.oregan@nuigalway.ie