

DYNAMICS OF A CERTAIN SEQUENCE OF POWERS

ROMAN SZNAJDER and KANCHAN BASNYAT

(Received 14 September 1998 and in revised form 15 January 1999)

ABSTRACT. For any nonzero complex number z we define a sequence $a_1(z) = z$, $a_2(z) = z^{a_1(z)}$, \dots , $a_{n+1}(z) = z^{a_n(z)}$, $n \in \mathbb{N}$. We attempt to describe the set of these z for which the sequence $\{a_n(z)\}$ is convergent. While it is almost impossible to characterize this convergence set in the complex plane \mathbb{C} , we achieved it for positive reals. We also discussed some connection to the Euler's functional equation.

Keywords and phrases. Power sequence, dynamics, convergence.

2000 Mathematics Subject Classification. Primary 26A18, 39B12; Secondary 11B37, 37A45.

1. Introduction. In this paper, for any $z \in \mathbb{C} \setminus \{0\}$, we consider the sequence

$$a_1(z) = z, \quad a_2(z) = z^z = z^{a_1(z)}, \dots, a_{n+1}(z) = z^{a_n(z)}, \quad n = 1, 2, \dots \quad (1.1)$$

We wish to characterize the convergence set $L := \{z \in \mathbb{C} \setminus \{0\} : \{a_n(z)\} \text{ is convergent}\}$. Since $1 \in L$, this set is nonempty. For a complex value of z , the mapping $z \rightarrow a_n(z)$ is a multivalued operation and $a_n(z)$ usually represents an infinite set. If we agree to use the main branch of the complex logarithm (as it is done, e.g., in MATLAB), the above exponentiation becomes a single-valued operation. For arbitrary z , it is very easy to run out of accuracy of any available computational software in a few iterations. Thus, it is practically impossible to describe the convergence set L . Also, one can show that the sequence $\{a_n(z)\}$ cannot be obtained as a functional iteration, i.e., there exists no function $f : \mathbb{C} \rightarrow \mathbb{C}$ such that $a_n(z) = f^{[n]}(z) = f \circ f \circ \dots \circ f(z)$ (n -times). This sequence has nasty numerical features. Nevertheless, the situation is not hopeless when it comes to considering real values of $a_n(z)$.

2. Main result. Our main goal is to demonstrate the following.

THEOREM 2.1. *We have $L \cap \mathbb{R}_+ = [e^{-e}, e^{1/e}]$.*

Before we proceed with the proof of Theorem 2.1, the following lemma is in order.

LEMMA 2.2. *For the function $f(t) = \ln(t)/t$ defined on $\mathbb{R}_+ = (0, +\infty)$, the following hold true:*

- (a) $\text{range}(f) = (-\infty, 1/e]$,
- (b) $\text{for any } 0 < c < 1/e, \text{ the equation } f(t) = c \text{ has exactly two solutions, while for } c = 1/e \text{ exactly one, namely } t^* = e.$

PROOF. The first derivative of the function $f(t)$ is $f'(t) = (1 - \ln(t))/t^2$, so that $f'(t) = 0$ if and only if $t = e$, f is increasing on $(0, e]$ and decreasing on $[e, +\infty)$. Since

$\lim_{t \rightarrow 0^+} f(t) = -\infty$ and $\lim_{t \rightarrow +\infty} f(t) = 0$, we obtain statement (a). Since $t^* = e$ is the only maximizer of $f(t)$ in $(0, +\infty)$, (b) follows from the intermediate value theorem and the fact that $\lim_{t \rightarrow +\infty} f(t) = 0$. The proof is complete. \square




FIGURE 2.1.

Note that if $t_c^{(1)}$ and $t_c^{(2)}$ denote two distinct solutions of the equation $f(t) = c$, then $t_c^{(1)} \rightarrow 1^+$ and $t_c^{(2)} \rightarrow +\infty$ as $c \rightarrow 0^+$. This remark follows from the piecewise monotonicity of the function $f(t)$ and the fact that $\lim_{t \rightarrow 1} f(t) = 0$ and $\lim_{t \rightarrow +\infty} f(t) = 0$. By putting $c = \ln(x)$ in Lemma 2.2, we get the following.

COROLLARY 2.3. *For any given $x \in (1, e^{1/e})$ there exist exactly two solutions of the equation $x^\alpha = \alpha$, while for $x = e^{1/e}$, exactly one.*

LEMMA 2.4. *Let $K(x) := \{0 \leq t \leq 1 : h_x(t) = 0\}$, where $h_x(t) = x^{x^t} - t$ and $x > 0$. The following statements hold true:*

- (a) *For $x \in (0, e^{-e})$, $K(x)$ consists of three distinct elements.*
- (b) *For $x \in [e^{-e}, 1)$, $K(x)$ is a singleton.*

PROOF. Instead of giving a rigorous analytic proof similar to Bomberger [1], we rather illustrate our point with an aid of appropriate graphs.

CASE 1 ($x \in (0, e^{-e})$). See Figure 2.2.

CASE 2 ($x = e^{-e}$). See Figure 2.3.

In fact, $h_x'(t) = x^t x^{x^t} (\ln(x))^2 - 1$, so $h_{e^{-e}}'(1/e) = 0$. Also, $h_{e^{-e}}(1/e) = 0$.

CASE 3 ($x > e^{-e}$). See Figure 2.4.

This completes our proof. \square

PROOF OF THEOREM 2.1

CASE 1 ($0 < x < e^{-e}$). By Lemma 2.4 (see Figure 2.2), there exist numbers $0 < \alpha_1 <$

$\alpha_2 < \alpha_3 < 1$ such that

$$x^{x^{\alpha_i}} = \alpha_i \quad \text{for } i = 1, 2, 3. \quad (2.1)$$

While α_2 is the fixed point of the function $g_x(t) = x^t$, α_1 and α_3 are points of period 2 ($x^{\alpha_1} = \alpha_3$ and $x^{\alpha_3} = \alpha_1$).

Generating the sequence $\{a_n(x)\}$ by putting $a_1(x) := x = g_x(1)$ and $a_{n+1}(x) = g_x(a_n(x))$ for $n \geq 1$, as illustrated by Figure 2.5, we get

$$a_1(x) < a_3(x) < \dots < a_{2n-1}(x) \nearrow \alpha_1, \quad a_2(x) > a_4(x) > \dots > a_{2n}(x) \searrow \alpha_3. \quad (2.2)$$

Thus, $\{a_n(x)\}$ diverges but consists of two complementary convergent subsequences.

CASE 2 ($x = e^{-e}$). In this case, $\{a_n(x)\}$ is convergent and the limit is $1/e$. Indeed, the sequence $\{a_n(x)\}$ consists of two complementary subsequences and each of them is convergent to a point of period 2. But in this case, $\alpha_1 = \alpha_2 = \alpha_3 = 1/e$, and the two points of period 2 collapse to the fixed point of the function $g_{e^{-e}}(t)$. Hence, $\{a_n(x)\}$ converges to $1/e$. However its convergence is very slow (e.g., $a_{1,000,000}(e^{-e}) \approx 0.36697888108297$ and $|a_{1,000,000}(e^{-e}) - 1/e| \approx 0.00180$).

CASE 3 ($e^{-e} < x \leq 1$). For $x < 1$ we reason the same way as in Case 2, so the sequence $\{a_n(x)\}$ converges to a unique fixed point of the function $g_x(t)$. For $x = 1$, the statement is trivial.

CASE 4 ($1 < x < e^{1/e}$). If $\{a_n(x)\}$ is convergent to a limit α , $\{a_{n+1}(x)\}$ also converges to α , moreover, $a_{n+1}(x) = x^{a_n(x)}$ converges to x^α . Consequently, $x^\alpha = \alpha$. By Corollary 2.3, there are exactly two fixed points α_1 and α_2 ($\alpha_1 < \alpha_2$) of the function $g_x(t)$. Obviously, $1 < \alpha_1 < e < \alpha_2$. Since $g_x(t)$ is increasing, $1 < x < \alpha_1$, and we check by the math induction that $a_1(x) < a_2(x) < \dots < a_n(x) < a_1$, so the limit of the sequence $\{a_n(x)\}$ exists and is just a fixed point α_1 .




FIGURE 2.2.




FIGURE 2.3.

CASE 5 ($x = e^{1/e}$). As before, $1 < x < e$ and $\{a_n(x)\}$ is increasing and bounded by e , thus convergent to the only fixed point of $g_{e^{1/e}}(t)$, namely e itself.

CASE 6 ($x > e^{1/e}$). By Corollary 2.3, no fixed point of $g_x(t)$ exists, so the sequence $\{a_n(x)\}$ does not converge. Now, the proof is complete. \square

We note that in Case 4 of the above proof, α_1 is an attracting fixed point, while α_2 is repelling fixed point for $g_x(t)$. In the limiting Case 5, the point $\alpha = e$ is left




FIGURE 2.4.

attracting and right repelling. Our Corollary 2.3 and Lemma 2.4 have some interesting relations to certain functional equations, one of them considered already by Euler (see Sierpiński [2]):

(a) $0 < x < e^{-e}$. As we have already noticed, there exist exactly three zeros of the function $h_x(t)$, say, $\alpha_1 < \alpha_2 < \alpha_3$, so that $x^{x^{\alpha_i}} = \alpha_i$ for $i = 1, 2, 3$, see (2.1); α_1 and α_3 are points of period 2 of the function $g_x(t)$, i.e., $x^{\alpha_1} = \alpha_3$ and $x^{\alpha_3} = \alpha_1$; (α_2 is just the fixed point of the function $g_x(t)$). Thus,

$$\alpha_1^{\alpha_1} = (x^{\alpha_3})^{\alpha_1} = (x^{\alpha_1})^{\alpha_3} = \alpha_3^{\alpha_3}. \quad (2.3)$$

In other words, α_1 and α_3 satisfy the functional equation $u^u = v^v$. For a detailed account on solving this equation we refer to Bomberger [1]. Observe that all α_i 's are in $(0, 1)$.

(b) $1 < x < e^{1/e}$. As we noted after Lemma 2.2, there exist two distinct fixed points α_1 and α_2 of $g_x(t)$, and when $x \rightarrow 1^+$, $\alpha_1 \rightarrow 1^+$, and $\alpha_2 \rightarrow +\infty$. Since $x^{\alpha_1} = \alpha_1$ and $x^{\alpha_2} = \alpha_2$, we get

$$\alpha_1^{\alpha_2} = (x^{\alpha_1})^{\alpha_2} = (x^{\alpha_2})^{\alpha_1} = \alpha_2^{\alpha_1}, \quad (2.4)$$

thus, α_1 and α_2 are related by the equation $u^v = v^u$ ($u, v > 0$), called the Euler equation.

(c) $e^{-e} \leq x \leq 1$. Now, α_1 and α_3 considered in (a) collapse to the same point and both types of equations considered in (a) and (b) are automatically satisfied.




FIGURE 2.5.

For the sake of completeness of this exposition we describe the set of solutions of the Euler equation.

PROPOSITION 2.5. *The set of solutions of the Euler equation $u^v = v^u$, where $u \neq v$, $u, v > 0$ is given by a parametrized family $\{(u, v) = (\alpha^{1/(\alpha-1)}, \alpha^{\alpha/(\alpha-1)}) : \alpha > 1\}$.*

PROOF. We note that if $0 < u < v \leq 1$, then $u < u^v < u^u < v^u$, and a pair (u, v) cannot be a solution. Similarly, when $0 < u \leq 1 < v$. Observe that $\ln(u)/u = \ln(v)/v$, and if this common value is less than $1/e$, by Lemma 2.2 there exist exactly two solutions u and v of this equation and $1 < u < e < v$. We postulate a solution in the form $u = t$ and $v = t^\alpha$, where $t > 1$ and $\alpha > 0$ are parameters. Thus,

$$t^{t^\alpha} = (t^\alpha)^t = t^{\alpha t}, \quad (2.5)$$

so $t^\alpha = \alpha t$ and, in consequence, $t = \alpha^{1/(\alpha-1)}$. Hence, $(u, v) = (\alpha^{1/(\alpha-1)}, \alpha^{\alpha/(\alpha-1)})$. For any $1 \neq \alpha > 0$, $\alpha^{1/(\alpha-1)} > 1$, so in order to keep $u < v$, α must be greater than 1. We check that the pair (u, v) actually solves the equation:

$$u^v = (\alpha^{1/(\alpha-1)})^{\alpha^{\alpha/(\alpha-1)}} = \alpha^{1/(\alpha-1)\alpha^{\alpha/(\alpha-1)}} = \alpha^{1/(\alpha-1)\alpha^{1+1/(\alpha-1)}} = (\alpha^{\alpha/(\alpha-1)})^{\alpha^{1/(\alpha-1)}} = v^u. \quad (2.6)$$

Now, we show that this parametrized family exhausts all possible solutions of the Euler equation such that $u < v$. In fact, if $\alpha \rightarrow 1^+$, $\alpha^{1/(\alpha-1)} \rightarrow e$, and $\alpha^{\alpha/(\alpha-1)} \rightarrow e$; if $\alpha \rightarrow +\infty$, $\alpha^{1/(\alpha-1)} \rightarrow 1$, and $\alpha^{\alpha/(\alpha-1)} \rightarrow +\infty$. The result now follows from the intermediate value theorem. \square

ACKNOWLEDGEMENTS. We are very grateful to Dr. Denny Gulick from the University of Maryland, College Park and an anonymous referee whose suggestions and comments were invaluable for improving the presentation of this paper. This research was supported by the Fall 1997 MIE Faculty Grant.

REFERENCES

- [1] J. D. Bomberger, *On the solutions of $a^a = b^b$* , Pi Mu Epsilon J. **9** (1993), no. 9, 571-572. Zbl 875.26013.
- [2] W. Sierpiński, *Elementary Theory of Numbers*, 2nd ed., Elsevier Science Publ. Co., New York, 1973.

ROMAN SZNAJDER: DEPARTMENT OF MATHEMATICS, BOWIE STATE UNIVERSITY, BOWIE, MARYLAND 20715, USA

E-mail address: rsznajde@bowiestate.edu

KANCHAN BASNYAT: DEPARTMENT OF COMPUTER SCIENCE, BOWIE STATE UNIVERSITY, BOWIE, MARYLAND 20715, USA

E-mail address: kbasnyat@cs.bowiestate.edu

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.oter@usc.es