

## TRANSIENT PROBABILITIES FOR A SIMPLE BIRTH-DEATH-IMMIGRATION PROCESS UNDER THE INFLUENCE OF TOTAL CATASTROPHES

RANDALL J. SWIFT

(Received 27 August 2000)

**ABSTRACT.** The transient probabilities for a simple birth-death-immigration process are considered. Catastrophes occur at a constant rate, and when they occur, reduce the population to size zero.

2000 Mathematics Subject Classification. Primary 60J80.

**1. Introduction.** In this note, a simple birth-death-immigration process is considered, which is influenced by total catastrophes which, when they occur, reduce the population size to zero. Population processes under the influence of various types of catastrophes have been studied by Bartoszynski et al. [2], Brockwell et al. [3], and Kyriakidis [4, 5].

The process is formulated by letting  $N(t)$  represent the size of the population at time  $t$  and

$$P_n(t) = P[N(t) = n \mid N(0) = 0]. \quad (1.1)$$

As in the simple birth-death process, births and deaths occur proportional to the population size with a birth rate  $\lambda > 0$  and a death rate  $\mu > 0$ . Immigration will occur independent of population size with rate  $\alpha > 0$ . Further, the occurrence of a catastrophe is also independent of population size and will occur at a rate  $\gamma > 0$ . Thus, the process can be described by the following transition rates:

---

Transition	Rate	
$i \rightarrow i + 1$	$\lambda i + \alpha$	$(i \geq 0)$
$i \rightarrow i - 1$	$\mu i,$	$(i \geq 1)$
$i \rightarrow 0$	$\gamma$	$(i \geq 1)$

---

The special case of  $\gamma = 1$  was recently considered by Kyriakidis [4], who obtained the stationary probabilities for this process. The transient probabilities of a simple immigration-catastrophe process, where  $\lambda = 0$  and  $\mu = 0$ , was obtained by Swift [6]. In the next section, the transient probabilities for the general case are derived.

**2. The transient probabilities.** The standard argument using the forward Kolmogorov equations shows that  $P_n(t)$  satisfies

$$P'_n(t) = (n+1)\mu P_{n+1}(t) + ((n-1)\lambda + \alpha)P_{n-1}(t) - (n(\mu + \lambda) + \alpha + \gamma)P_n(t), \quad (2.1)$$

$$P'_0(t) = \gamma \sum_{i=1}^{\infty} P_i(t) + \mu P_1(t) - \alpha P_0(t). \quad (2.2)$$

Now

$$\sum_{i=1}^{\infty} P_i(t) = 1 - P_0(t) \quad (2.3)$$

so that

$$P'_0(t) = \gamma + \mu P_1(t) - (\alpha + \gamma)P_0(t). \quad (2.4)$$

Letting

$$\psi(s, t) = \sum_{k=0}^{\infty} P_k(t) s^k \quad (2.5)$$

be the probability generating function (PGF) for the system, it follows from the standard generating function method, that  $\psi(s, t)$  satisfies the partial differential equation

$$\frac{\partial \psi(s, t)}{\partial t} = (\lambda s - \mu)(s-1) \frac{\partial \psi(s, t)}{\partial s} + \alpha(s-1)\psi(s, t) + \gamma(1 - \psi(s, t)). \quad (2.6)$$

Since we are considering a process with immigrations, we can assume, for the sake of simplicity, the initial condition  $P_0(0) = 1$ . This gives  $\psi(s, 0) = 1$ .

As in most birth-death processes, the solution of the partial differential equation (2.6) for  $\psi(\cdot, \cdot)$  depends upon the values of the parameters  $\lambda$  and  $\mu$ . The PGF  $\psi(\cdot, \cdot)$  is given below for three important cases. The method of solution of (2.6) is standard and indeed, it is interesting to note that these solutions can also be obtained using a computer algebra system such as Mathematica.

**CASE 2.1** ( $\lambda \neq \mu$  and  $\lambda \neq 0$ ). The PDE (2.6) has solution

$$\begin{aligned} \psi(s, t) = & \left( \frac{1}{(\alpha + \gamma)\lambda - \alpha\mu} \right) \\ & \times \left[ \gamma \lambda \left( \frac{(1-s)\lambda}{\lambda - \mu} \right)^{\gamma/(\lambda - \mu)} {}_2F_1 \left( \frac{\alpha\lambda + \gamma\lambda - \alpha\mu}{\lambda^2 - \lambda\mu}, \frac{\lambda - \mu + \gamma}{\lambda - \mu}, \frac{(\lambda + \alpha)(\lambda - \mu) + \gamma\lambda}{\lambda(\lambda - \mu)}, \frac{\lambda s - \mu}{\lambda - \mu} \right) \right. \\ & + (s-1)^{\gamma/(\lambda - \mu)} (\lambda s - \mu)^{-\alpha/\lambda + \gamma/(\mu - \lambda)} \left( \frac{(\mu - \lambda)(\lambda s - \mu)}{\mu - \lambda s + \lambda(s-1)e^{(\lambda - \mu)t}} \right)^{\alpha/\lambda + \gamma/(\lambda - \mu)} \\ & \times \left( \frac{(\mu - \lambda)(s-1)}{\lambda(s-1) + (\mu - \lambda s)e^{-(\lambda - \mu)t}} \right)^{-\gamma/(\lambda - \mu)} \\ & \times \left( ((\alpha + \gamma)\lambda - \alpha\mu) - \gamma \lambda \left( \frac{\lambda}{\lambda(s-1) + (\mu - \lambda s)e^{-(\lambda - \mu)t}} \right)^{\gamma/(\lambda - \mu)} \right. \\ & \times \left. {}_2F_1 \left( \frac{\alpha\lambda + \gamma\lambda - \alpha\mu}{\lambda^2 - \lambda\mu}, \frac{\lambda - \mu + \gamma}{\lambda - \mu}, \frac{(\lambda + \alpha)(\lambda - \mu) + \gamma\lambda}{\lambda(\lambda - \mu)}, (\mu - \lambda s)e^{-(\lambda - \mu)t} \right) \right] \end{aligned} \quad (2.7)$$

where  ${}_2F_1(a, b; c, z)$  is the hypergeometric function defined by

$${}_2F_1(a, b; c, z) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k} \frac{z^k}{k!} \quad (2.8)$$

(cf. Agarwal [1]).

**CASE 2.2** ( $\lambda = \mu$  and  $\lambda \neq 0$ ). In this case, the PDE (2.6) has solution

$$\begin{aligned} \psi(s, t) &= \frac{\gamma e^{\gamma/\lambda(1-s)}}{\lambda(1-s)} \int_1^{\infty} z^{-\alpha/\lambda} e^{-\gamma z/\lambda(1-s)} dz - \frac{e^{-\gamma t}}{\lambda(1-s)} \left( \frac{1}{1 + (1-s)t\lambda} \right)^{\alpha/\lambda} \\ &\quad \times \left( \lambda(s-1) + \gamma(1 + \gamma\lambda t(1-s)) \int_1^{\infty} z^{-\alpha/\lambda} e^{-\gamma(t+1/\lambda(1-s))z} dz \right). \end{aligned} \quad (2.9)$$

**CASE 2.3** ( $\lambda = 0$  and  $\mu \neq 0$ ). In this case, the PGF is given as

$$\begin{aligned} \psi(s, t) &= \frac{1}{\mu} \left[ e^{-\alpha s/\mu} \left( \gamma e^{\alpha/\mu} \int_1^{\infty} z^{-(1+\gamma/\mu)} e^{-\alpha(1-s)z/\mu} dz \right. \right. \\ &\quad \left. \left. + e^{-\gamma t} \left( \mu e^{\alpha(1-(1-s)e^{\mu t})/\mu} - \gamma e^{\alpha/\mu} \int_1^{\infty} z^{-(1+\gamma/\mu)} e^{-\alpha(1-s)e^{\mu t}/\mu} dz \right) \right) \right]. \end{aligned} \quad (2.10)$$

Using the PGF, the probability of extinction,  $P_0(t)$ , can be obtained in each of the above cases.

**CASE 2.1** ( $\lambda \neq \mu$  and  $\lambda \neq 0$ ).

$$\begin{aligned} P_0(t) &= \psi(0, t) \\ &= \frac{1}{(\alpha+\gamma)\lambda-\alpha\mu} \left[ \left( \frac{\lambda-\mu}{\lambda e^{(\lambda-\mu)t}-\mu} \right)^{\alpha/\lambda+\gamma/(\lambda-\mu)} \left( \frac{\mu-\lambda}{\lambda-\mu e^{(-\lambda+\mu)t}} \right)^{\gamma/(\mu-\lambda)} \right. \\ &\quad \times \left( \alpha\mu - (\alpha+\gamma)\lambda + \gamma\lambda \left( \frac{\lambda}{\lambda-\mu e^{(-\lambda+\mu)t}} \right)^{\gamma/(\lambda-\mu)} \right. \\ &\quad \times {}_2F_1 \left( \frac{\alpha(\lambda-\mu)+\lambda\gamma}{\lambda(\lambda-\mu)}, \frac{\lambda-\mu+\gamma}{\lambda-\mu}, \frac{(\lambda+\alpha)(\lambda-\mu)+\gamma\lambda}{\lambda(\lambda-\mu)}, \frac{\mu}{\mu-\lambda e^{(\lambda-\mu)t}} \right) \\ &\quad \left. \left. + \gamma\lambda \left( \frac{\lambda}{\lambda-\mu} \right)^{\gamma/(\lambda-\mu)} \right. \right. \\ &\quad \times {}_2F_1 \left( \frac{\alpha(\lambda-\mu)+\lambda\gamma}{\lambda(\lambda-\mu)}, \frac{\lambda-\mu+\gamma}{\lambda-\mu}, \frac{(\lambda+\alpha)(\lambda-\mu)+\gamma\lambda}{\lambda(\lambda-\mu)}, \frac{\mu}{\mu-\lambda} \right) \right]. \end{aligned} \quad (2.11)$$

**CASE 2.2** ( $\lambda = \mu$  and  $\lambda \neq 0$ ).

$$\begin{aligned} P_0(t) &= e^{\gamma/\lambda} \left[ \left( \frac{1}{1+\lambda t} \right)^{\alpha/\lambda} \left( e^{-\gamma(1+\lambda t)/\lambda} - \frac{1}{\lambda} \left( (\gamma+\gamma\lambda t) \int_1^{\infty} z^{-\alpha/\lambda} e^{-\gamma(t+1/\lambda)z} dz \right) \right) \right. \\ &\quad \left. + \frac{\gamma}{\lambda} \int_1^{\infty} z^{-\alpha/\lambda} e^{-(\gamma/\lambda)z} dz \right]. \end{aligned} \quad (2.12)$$

**CASE 2.3** ( $\lambda = 0$  and  $\mu \neq 0$ ).

$$P_0(t) = \frac{e^{-yt}}{\mu} \left( \mu e^{-\alpha(e^{t\mu}-1)/\mu} - ye^{\alpha/\mu} \int_1^\infty z^{-(y+\mu)/\mu} e^{-(\alpha e^{t\mu}/\mu)z} dz \right) \\ + \frac{ye^{\alpha/\mu}}{\mu} \int_1^\infty z^{-(y+\mu)/\mu} e^{-(\alpha/\mu)z} dz. \quad (2.13)$$

We note here that these expressions for  $P_0(t)$ , as  $t \rightarrow \infty$ , with  $y = 1$ , reduce to the stationary probabilities obtained by Kyriakidis.

The probabilities  $P_n(t)$ , for  $n \geq 1$  can be obtained by expanding  $\psi(s, t)$  as a power series in  $s$ . However, the nature of the representations (2.7), (2.9), and (2.10) for  $\psi(s, t)$  makes this a formidable task. Alternatively,  $P_0(t)$  can be used in (2.4) to obtain  $P_1(t)$  then (2.1) can be used recursively, to obtain  $P_n(t)$  for  $n \geq 1$ .

## REFERENCES

- [1] R. P. Agarwal, *Generalized Hypergeometric Series*, Uttar Pradesh Scientific Research Committee, Allahabad, India, Asia Publishing House, Bombay, 1963. [MR 31#3636](#).
- [2] R. Bartoszynski, W. J. Buehler, W. Chan, and D. K. Pearl, *Population processes under the influence of disasters occurring independently of population size*, J. Math. Biol. **27** (1989), no. 2, 167–178. [MR 90h:92017](#). [Zbl 715.92022](#).
- [3] P. J. Brockwell, J. Gani, and S. I. Resnick, *Birth, immigration and catastrophe processes*, Adv. in Appl. Probab. **14** (1982), no. 4, 709–731. [MR 84d:60107](#). [Zbl 496.92007](#).
- [4] E. G. Kyriakidis, *Stationary probabilities for a simple immigration-birth-death process under the influence of total catastrophes*, Statist. Probab. Lett. **20** (1994), no. 3, 239–240. [CMP 1 294 110](#). [Zbl 801.60073](#).
- [5] E. G. Kyriakidis and A. Abakuks, *Optimal pest control through catastrophes*, J. Appl. Probab. **26** (1989), no. 4, 873–879. [MR 90m:92066](#). [Zbl 688.60069](#).
- [6] R. J. Swift, *A simple immigration-catastrophe process*, Math. Sci. **25** (2000), no. 1, 32–36. [CMP 1 771 175](#).

RANDALL J. SWIFT: DEPARTMENT OF MATHEMATICS, WESTERN KENTUCKY UNIVERSITY, BOWLING GREEN, KY 42101, USA

*E-mail address:* [randall.swift@wku.edu](mailto:randall.swift@wku.edu)

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

### Guest Editors

**Edson Denis Leonel**, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)