

ROUGH MARCINKIEWICZ INTEGRAL OPERATORS

HUSSAIN AL-QASSEM and AHMAD AL-SALMAN

(Received 16 January 2001)

ABSTRACT. We study the Marcinkiewicz integral operator $M_{\mathcal{P}}f(x) = (\int_{-\infty}^{\infty} |\int_{|y| \leq 2t} f(x - \mathcal{P}(y))(\Omega(y)/|y|^{n-1}) dy|^2 dt / 2^{2t})^{1/2}$, where \mathcal{P} is a polynomial mapping from \mathbb{R}^n into \mathbb{R}^d and Ω is a homogeneous function of degree zero on \mathbb{R}^n with mean value zero over the unit sphere \mathbb{S}^{n-1} . We prove an L^p boundedness result of $M_{\mathcal{P}}$ for rough Ω .

2000 Mathematics Subject Classification. 42B20, 42B15, 42B25.

1. Introduction. Let \mathbb{R}^n , $n \geq 2$ be the n -dimensional Euclidean space and \mathbb{S}^{n-1} be the unit sphere in \mathbb{R}^n equipped with the induced Lebesgue measure. Consider the Marcinkiewics integral operator

$$\mu f(x) = \left(\int_{-\infty}^{\infty} |\mathbf{F}_t(x)|^2 \frac{dt}{2^{2t}} \right)^{1/2}, \quad (1.1)$$

where

$$\mathbf{F}_t(x) = \int_{|x-y| \leq 2t} f(y) \frac{\Omega(x-y)}{|x-y|^{n-1}} dy, \quad (1.2)$$

and Ω is a homogeneous function of degree zero which has the following properties:

$$\Omega \in L^1(\mathbb{S}^{n-1}), \quad \int_{\mathbb{S}^{n-1}} \Omega(y') d\sigma(y') = 0. \quad (1.3)$$

When $\Omega \in \text{Lip}_{\alpha}(\mathbb{S}^{n-1})$, $(0 < \alpha \leq 1)$, Stein proved the L^p boundedness of $\mu(f)$ for all $1 < p \leq 2$. Subsequently, Benedek, Calderón, and Panzone proved the L^p boundedness of $\mu(f)$ for all $1 < p < \infty$ under the condition $\Omega \in C^1(\mathbb{S}^{n-1})$ (see [2]).

The authors of [3] were able to prove the following result for the more general class of operators

$$\mu_P f(x) = \left(\int_{-\infty}^{\infty} |\mathbf{F}_{P,t}(x)|^2 \frac{dt}{2^{2t}} \right)^{1/2}, \quad (1.4)$$

where

$$\mathbf{F}_{P,t}(x) = \int_{|y| \leq 2t} f(x - P(|y|)y') \frac{\Omega(y)}{|y|^{n-1}} dy \quad (1.5)$$

and P is a real-valued polynomial on \mathbb{R} and satisfies $P(0) = 0$.

THEOREM 1.1 (see [3]). *Let $\alpha > 0$, and $\Omega \in V_{\alpha}(n)$. Then the operator μ_P is bounded in $L^p(\mathbb{R}^n)$ for $(2\alpha+2)/(2\alpha+1) < p < 2+2\alpha$.*

In [1], Al-Salman and Pan studied the singular integral operator

$$\mathbf{T}_{\Omega, \mathcal{P}} f(x) = \text{p.v.} \int_{\mathbb{R}^n} f(x - \mathcal{P}(y)) \frac{\Omega(y')}{|y'|^n} dy, \quad (1.6)$$

where $\mathcal{P} = (P_1, \dots, P_d): \mathbb{R}^n \rightarrow \mathbb{R}^d$ is a polynomial mapping, $d \geq 1$, $n \geq 2$. The authors of [1] proved that $\mathbf{T}_{\Omega, \mathcal{P}}$ is bounded in $L^p(\mathbb{R}^d)$ whenever $(2+2\alpha)/(1+2\alpha) < p < 2+2\alpha$ and $\Omega \in W_\alpha(n)$. Here $W_\alpha(n)$ is a subspace of $L^1(\mathbf{S}^{n-1})$ and its definition as well as the definition of $V_\alpha(n)$ will be reviewed in Section 2. It was shown in [1] that $W_\alpha(n) = V_\alpha(n)$, if $n = 2$ and it is a proper subspace of $V_\alpha(n)$ if $n \geq 3$.

Our purpose in this paper is to study the L^p boundedness of the class of operators

$$M_{\mathcal{P}} f(x) = \left(\int_{-\infty}^{\infty} |\mathbf{F}_{\mathcal{P}, t}(x)|^2 \frac{dt}{2^{2t}} \right)^{1/2}, \quad (1.7)$$

where

$$\mathbf{F}_{\mathcal{P}, t}(x) = \int_{|y| \leq 2^t} f(x - \mathcal{P}(y)) \frac{\Omega(y)}{|y|^{n-1}} dy. \quad (1.8)$$

Our main result in this paper is the following theorem.

THEOREM 1.2. *Let $\alpha > 0$, and $\Omega \in W_\alpha(n)$. Then the operator $M_{\mathcal{P}}$ is bounded in $L^p(\mathbb{R}^d)$ for $(2\alpha+2)/(2\alpha+1) < p < 2+2\alpha$. The bound of $M_{\mathcal{P}} f$ is independent of the coefficients of $\{P_j\}$.*

By [1, Theorem 3.1] and Theorem 1.2 we have the following corollary.

COROLLARY 1.3. *Let $\alpha > 0$, $\Omega \in V_\alpha(2)$ and $\mathcal{P}: \mathbb{R}^2 \rightarrow \mathbb{R}^d$. Then $M_{\mathcal{P}}$ is bounded in $L^p(\mathbb{R}^d)$ for $(2\alpha+2)/(2\alpha+1) < p < 2+2\alpha$. The bound of $M_{\mathcal{P}} f$ is independent of the coefficients of $\{P_j\}$.*

2. Preparation. We start this section by recalling the following definition from [1].

DEFINITION 2.1. For $\alpha > 0$, $N \geq 1$, let $\tilde{\mathcal{V}}(n, N) = \bigcup_{m=1}^N \mathcal{V}(n, m)$ and let $W_\alpha(N, n)$ be the subspace of $L^1(\mathbf{S}^{n-1})$ defined by

$$W_\alpha(N, n) = \left\{ \Omega \in L^1(\mathbf{S}^{n-1}) : \int_{\mathbf{S}^{n-1}} \Omega(y') d\sigma(y') = 0, M_\alpha(N, n) < \infty \right\}, \quad (2.1)$$

where

$$\begin{aligned} M_\alpha(N, n) &= \max \left\{ \int_{\mathbf{S}^{n-1}} |\Omega(y')| \left(\log \frac{1}{|P(y')|} \right)^{1+\alpha} d\sigma(y') : P \in \tilde{\mathcal{V}}(n, N) \text{ with } \|P\| = 1 \right\}. \end{aligned} \quad (2.2)$$

For $\alpha > 0$, we define $W_\alpha(n)$ to be

$$W_\alpha(n) = \bigcap_{N=1}^{\infty} W_\alpha(N, n). \quad (2.3)$$

Also, for $\alpha > 0$, we define $V_\alpha(n)$ by $V_\alpha(n) = W_\alpha(1, n)$ (see [6]).

Here $\mathcal{V}(n, m)$ is the space of all real-valued homogeneous polynomials on \mathbb{R}^n with degree equal to m and with norm $\|\cdot\|$ defined by

$$\left\| \sum_{|\alpha|=m} a_\alpha y^\alpha \right\| = \sum_{|\alpha|=m} |a_\alpha|. \quad (2.4)$$

Now we need to recall the following results.

LEMMA 2.2 (see van der Corput [7]). *Suppose ϕ and ψ are real-valued and smooth in (a, b) , and that $|\phi^{(k)}(t)| \geq 1$ for all $t \in (a, b)$. Then the inequality*

$$\left| \int_a^b e^{-i\lambda\phi(t)} \psi(t) dt \right| \leq C_k |\lambda|^{-1/k} \left[|\psi(b)| + \int_a^b |\psi'(t)| dt \right], \quad (2.5)$$

holds when

- (i) $k \geq 2$, or
- (ii) $k = 1$ and ϕ' is monotonic.

The bound C_k is independent of a, b, ϕ , and λ .

LEMMA 2.3 (see [7]). *Let $\mathcal{P} = (P_1, \dots, P_d)$ be a polynomial mapping from \mathbb{R}^n into \mathbb{R}^d . Let $\deg(\mathcal{P}) = \max_{1 \leq j \leq d} \deg(P_j)$. Suppose $\Omega \in L^1(\mathbb{S}^{n-1})$ and*

$$\mu_{\Omega, \mathcal{P}} f(x) = \sup_{h>0} \left| \frac{1}{h^n} \int_{|y'| < h} f(x - \mathcal{P}(y')) \Omega(y') dy' \right|. \quad (2.6)$$

Then for every $1 < p \leq \infty$, there exists a constant $C_p > 0$ which is independent of Ω and the coefficients of $\{P_j\}$ such that

$$\|\mu_{\Omega, \mathcal{P}} f\|_p \leq C_p \|\Omega\|_{L^1(\mathbb{S}^{n-1})} \|f\|_p \quad (2.7)$$

for every $f \in L^p(\mathbb{R}^d)$.

To each polynomial mapping $\mathcal{P} = (P_1, \dots, P_d) : \mathbb{R}^n \rightarrow \mathbb{R}^d$ with

$$\deg \mathcal{P} = \max_{1 \leq j \leq d} \deg P_j = N, \quad d \geq 1, \quad n \geq 2, \quad (2.8)$$

we define a family of measures

$$\{\vartheta_t^l, \lambda_t^l : l = 0, 1, \dots, N, t \in \mathbb{R}\} \quad (2.9)$$

as follows.

For $1 \leq j \leq d$, $0 \leq l \leq N$ let $P_j = \sum_{|\alpha| \leq N} C_{j\alpha} y^\alpha$ and let $Q^l = (Q_1^l, \dots, Q_d^l)$ where $Q_j^l = \sum_{|\alpha| \leq l} C_{j\alpha} y^\alpha$.

Now for $0 \leq l \leq N$ and $t \in \mathbb{R}$, let ϑ_t^l and λ_t^l be the measures defined in the Fourier transform side by

$$\begin{aligned} (\vartheta_t^l)(\xi) &= \int_{|y| \leq 2^t} e^{-2\pi i \xi \cdot Q^l(y)} \frac{\Omega(y')}{|y|^{n-1}} \frac{dy}{2^t}, \\ (\lambda_t^l)(\xi) &= \int_{|y| \leq 2^t} e^{-2\pi i \xi \cdot Q^l(y)} \frac{|\Omega(y')|}{|y|^{n-1}} \frac{dy}{2^t}. \end{aligned} \quad (2.10)$$

The maximal functions $(\mathfrak{B}^l)^*$ defined by

$$(\mathfrak{B}^l)^*(f)(x) = \sup_{t \in \mathbb{R}} |\lambda_t^l * f(x)|, \quad (2.11)$$

for $l = 0, 1, \dots, N$.

For later purposes, we need the following definition.

DEFINITION 2.4. For each $1 \leq l \leq N$, let $N_l = |\{\alpha \in \mathbb{N}^n : |\alpha| = l\}|$ and let $\{\alpha \in \mathbb{N}^n : |\alpha| = l\} = \{\alpha_1, \dots, \alpha_{N_l}\}$. For each $1 \leq l \leq N$, define the linear transformations $L_l^{\alpha_j} : \mathbb{R}^d \rightarrow \mathbb{R}$ and $L_l : \mathbb{R}^d \rightarrow \mathbb{R}^{N_l}$ by

$$\begin{aligned} L_l^{\alpha_j}(\xi) &= \sum_{i=1}^d (C_{i,\alpha_j} \gamma^{\alpha_j}) \xi_i, \quad j = 1, \dots, N_l, \\ L_l(\xi) &= (L_l^{\alpha_1}(\xi), \dots, L_l^{\alpha_{N_l}}(\xi)). \end{aligned} \quad (2.12)$$

To simplify the proof of our result we need the following lemma.

LEMMA 2.5. Let $\{\sigma_t^l : l = 0, 1, \dots, N, t \in \mathbb{R}\}$ be a family of measures such that $\sigma_t^0 = 0$ for all $t \in \mathbb{R}$. Let $D_l : \mathbb{R}^n \rightarrow \mathbb{R}^d$, $l = 0, 1, \dots, N$ be linear transformations. Suppose that for all $t \in \mathbb{R}$ and $l = 0, 1, \dots, N$, then

$$\begin{aligned} \|\sigma_t^l\| &\leq C(l), \\ |(\sigma_t^l)\hat{(\xi)}| &\leq C \frac{M_\alpha}{(\log [c 2^{lt} |D_l(\xi)|])^{1+\alpha}}, \\ |(\sigma_t^l)\hat{(\xi)} - (\sigma_t^{l-1})\hat{(\xi)}| &\leq C 2^{lt} |D_l(\xi)|. \end{aligned} \quad (2.13)$$

Then there exists a family of measures $\{\nu_t^l : l = 1, \dots, N\}_{t \in \mathbb{R}}$ such that

$$\begin{aligned} \|\nu_t^l\| &\leq C(l), \\ |(\nu_t^l)\hat{(\xi)}| &\leq C \frac{M_\alpha}{(\log [c 2^{lt} |D_l(\xi)|])^{1+\alpha}}, \\ |(\nu_t^l)\hat{(\xi)}| &\leq C 2^{lt} |D_l(\xi)|, \\ \sigma_t^N &= \sum_{l=1}^N \nu_t^l. \end{aligned} \quad (2.14)$$

PROOF. By [5, Lemma 6.1], for each $l = 1, \dots, N$ choose two nonsingular linear transformations

$$A_l : \mathbb{R}^{r(l)} \rightarrow \mathbb{R}^d, \quad B_l : \mathbb{R}^d \rightarrow \mathbb{R}^{r(l)}, \quad (2.15)$$

such that

$$|A_l \pi_{r(l)}^d B_l(\xi)| \leq |D_l(\xi)| \leq N |A_l \pi_{r(l)}^d B_l(\xi)|, \quad \xi \in \mathbb{R}^d, \quad (2.16)$$

where $r(l) = \text{rank}(D_l)$ and $\pi_{r(l)}^d$ is the projection operator from \mathbb{R}^d into $\mathbb{R}^{r(l)}$.

Now choose $\eta \in C_0^\infty(\mathbb{R})$ such that $\eta(t) = 1$ for $|t| \leq 1/2$ and $\eta(t) = 0$ for $|t| \geq 1$. Let $\varphi(t) = \phi(t^2)$ and let

$$\begin{aligned} (\nu_t^l)(\xi) &= (\sigma_t^l)(\hat{\xi}) \prod_{l < j \leq N} \varphi(|2^{tj} A_j \pi_{r(j)}^d B_j(\xi)|) \\ &\quad - (\sigma_t^{l-1})(\hat{\xi}) \prod_{l-1 < j \leq N} \varphi(|2^{tj} A_j \pi_{r(j)}^d B_j(\xi)|) \end{aligned} \quad (2.17)$$

with the convention $\prod_{j \in \emptyset} a_j = 1$, $1 \leq l \leq N$.

Hence, one can easily see that $\{\sigma_t^l : l = 1, \dots, N, t \in \mathbb{R}\}$ is the desired family of measures. \square

Now for the boundedness of the maximal functions $(\vartheta^l)^*, l = 0, 1, \dots, N$, we have the following lemma whose proof is an easy consequence of [Lemma 2.3](#), polar coordinates and Hölder's inequality:

LEMMA 2.6. *For $l = 1, \dots, N$ and $p \in (1, \infty)$, there exists a constant $C_{p,l}$ which is independent of the coefficients of the polynomial components of the mapping Q^l such that*

$$\|(\vartheta^l)^* f\|_p \leq C_{p,l} \|f\|_p. \quad (2.18)$$

3. Boundedness of some square functions. For a nonnegative C^∞ radial function Φ on \mathbb{R}^n with

$$\text{supp}(\Phi) \subset \left\{ x \in \mathbb{R}^n : \frac{1}{2} \leq |x| \leq 2 \right\}, \quad \int_0^\infty \frac{\Phi(t)}{t} dt = 1, \quad (3.1)$$

and for a linear transformation $\mathbf{L} : \mathbb{R}^n \rightarrow \mathbb{R}^d$, define the functions $\psi_t, t \in \mathbb{R}$ by $\hat{\psi}_t(\gamma) = \Phi(2^t \mathbf{L}(\gamma))$.

For a family of measures $\{\sigma_t\}_{t \in \mathbb{R}}$, real number u and $l \in \mathbb{N}$, let $\mathbf{J}_u^l(f)$ be the square function defined by

$$\mathbf{J}_u^l(f)(x) = \left(\int_{-\infty}^\infty |\sigma_t * \psi_{l(t+u)} * f(x)|^2 dt \right)^{1/2}. \quad (3.2)$$

For such a square function we have the following theorem.

THEOREM 3.1. *If $\{\sigma_t\}_{t \in \mathbb{R}}$ is a family of measures such that the corresponding maximal function*

$$\sigma^*(f)(x) = \sup_{t \in \mathbb{R}} |\sigma_t * f(x)| \quad (3.3)$$

is bounded on $L^p(\mathbb{R}^d)$ for every $1 < p < \infty$, then

$$\|\mathbf{J}_u^l(f)\|_{L^p(\mathbb{R}^d)} \leq C_{p,l} \sqrt{\|\sigma^*\|_{(p/2)'}} \sup_{t \in \mathbb{R}} \|\sigma_t\| \|f\|_{L^p(\mathbb{R}^d)} \quad (3.4)$$

for every $1 < p < \infty$. Here $C_{p,l}$ is a constant that depends only on p and the dimension of the underlying space.

PROOF. If $\sup_{t \in \mathbb{R}} \|\sigma_t\| = \infty$, then the inequality holds trivially. Thus we may assume that $\sup_{t \in \mathbb{R}} \|\sigma_t\| < \infty$. In this case we follow a similar argument as in [4]. Let $p > 2$ and $q = (p/2)'$. Choose a nonnegative function $v \in L_+^q$ with $\|v\|_q = 1$ such that

$$\|\mathbf{J}_u^l(f)\|_p^2 = \int_{\mathbb{R}^d} \left(\int_{-\infty}^{\infty} |\sigma_t * \psi_{l(t+u)} * f(x)|^2 dt \right) v(x) dx. \quad (3.5)$$

Thus it is easy to see that

$$\begin{aligned} \|\mathbf{J}_u^l(f)\|_p^2 &\leq \sup_{t \in \mathbb{R}} \|\sigma_t\| \int_{-\infty}^{\infty} \int_{\mathbb{R}^d} |\psi_{l(t+u)} * f(z)|^2 \sigma^*(v)(-z) dz dt \\ &\leq \sup_{t \in \mathbb{R}} \|\sigma_t\| \int_{\mathbb{R}^d} [\mathcal{G}(f)]^2(z) \sigma^*(v)(-z) dz, \end{aligned} \quad (3.6)$$

where

$$\mathcal{G}(f)(x) = \left(\int_{-\infty}^{\infty} |\psi_{l(t+u)} * f(x)|^2 dt \right)^{1/2}. \quad (3.7)$$

Now since $\int_{\mathbb{R}^d} \psi_t(x) dx = 0$, it is well known that

$$\|\mathcal{G}(f)\|_p \leq C_p \|f\|_p \quad \forall 1 < p < \infty \quad (3.8)$$

with constant C_p that depends only on p and the dimension of the underlying space.

Thus by (3.6) and Hölder's inequality we have

$$\begin{aligned} \|\mathbf{J}_u^l(f)\|_p^2 &\leq \sup_{t \in \mathbb{R}} \|\sigma_t\| \|\mathcal{G}(f)\|_p^2 \|\sigma^*(u)\|_q \\ &\leq C_p^2 \sup_{t \in \mathbb{R}} \|\sigma_t\| \|\sigma^*\|_{(p/2)'} \|f\|_p^2. \end{aligned} \quad (3.9)$$

Hence our result follows by taking the square root on both sides. The case $p < 2$ follows by duality. \square

4. Proof of the main theorem. Let $\alpha > 0$, $\Omega \in W_\alpha(n)$. Let $\mathcal{P} = (P_1, \dots, P_d)$ be a polynomial mapping from \mathbb{R}^n into \mathbb{R}^d with $\deg \mathcal{P} = \max_{1 \leq j \leq d} \deg P_j = N$, where $d \geq 1$ and $n \geq 2$. For $0 \leq l \leq N$ let N_l , Q^l , ν_t^l , λ_t^l , and L_l be as in Section 3.

The first step in our proof is to show that each ϑ_t^l , $l = 1, \dots, N$ satisfies the hypotheses of Lemma 2.5, that is,

$$\|\vartheta_t^l\| \leq C(l), \quad (4.1)$$

$$|(\vartheta_t^l)(\xi)| \leq C \frac{M_\alpha}{(\log [c 2^{lt} |L_l(\xi)|])^{1+\alpha}}, \quad (4.2)$$

$$|(\vartheta_t^l)(\xi) - (\vartheta_t^{l-1})(\xi)| \leq C 2^{lt} |L_l(\xi)|. \quad (4.3)$$

One can easily see that (4.1) holds trivially. Using the cancellation property of Ω , it is easy to see that (4.3) holds. Thus, we need only to verify (4.2). To see that, we notice that

$$|(\vartheta_t^l)(\xi)| \leq \int_{S^{n-1}} |\Omega(y')| \left| \int_0^1 e^{-2\pi i \xi \cdot Q^l(2^t r y')} dr \right| d\sigma(y'). \quad (4.4)$$

Now the quantity $\xi \cdot Q^l(2^{tl}ry')$ can be written in the form

$$\xi \cdot Q^l(2^{tl}ry') = 2^{tl}r^l \lambda G^l(y') + \xi \cdot R(2^t r y'), \quad (4.5)$$

where Q^l is a homogeneous polynomial of degree l with $\|G^l\| = 1$, R is a polynomial of degree at most $l-1$ in the variable r ,

$$\lambda = \sum_{j=1}^{N_l} |L_l^{\alpha_j}(\xi)| \geq N_l |L_l(\xi)| \quad (4.6)$$

and $\alpha_1, \dots, \alpha_{N_l}$ are the constants that appeared in [Section 2](#). Thus by van der Corput lemma, we have

$$\left| \int_0^1 e^{-2\pi i \xi \cdot Q^l(2^t r y')} dr \right| \leq C \min \left\{ 1, (2^{tl} |L_l(\xi)| |G^l(y')|)^{-1/l} \right\} \quad (4.7)$$

and hence

$$\left| \int_0^1 e^{-2\pi i \xi \cdot Q^l(2^t r y')} dr \right| \leq C \frac{[\log(c |G^l(y')|^{-1})]^{1+\alpha}}{(\log[c 2^{tl} |L_l(\xi)|])^{1+\alpha}}, \quad (4.8)$$

where C is a constant independent of t and ξ . Since $\Omega \in W_\alpha(n)$, the estimate (4.2) follows.

By [Lemma 2.5](#), there exists a family of measures $\{\nu_t^l : l = 1, \dots, N, t \in \mathbb{R}\}$ such that

$$\|\nu_t^l\| \leq C(l), \quad (4.9)$$

$$|(\nu_t^l)(\xi)| \leq C \frac{M_\alpha}{(\log[c 2^{lt} |L_l(\xi)|])^{1+\alpha}}, \quad (4.10)$$

$$|(\nu_t^l)(\xi)| \leq C 2^{lt} |L_l(\xi)|, \quad (4.11)$$

$$\vartheta_t^N = \sum_{l=1}^N \nu_t^l. \quad (4.12)$$

Also by [Lemma 2.6](#) and the definition of ν_t^l (see the proof of [Lemma 2.5](#)), we have

$$\|(\nu^l)^* f\|_p \leq C_{p,l} \|f\|_p \quad \forall 1 < p < \infty. \quad (4.13)$$

Now one can easily see that

$$2^{-t} \mathbf{F}_{\vartheta,t}(x) = \vartheta_t^N * f(x) = \sum_{l=1}^N \nu_t^l * f(x). \quad (4.14)$$

Therefore,

$$\|M_{\vartheta} f\|_p \leq \sum_{l=1}^N \|M_{\vartheta}^l f\|_p, \quad (4.15)$$

where

$$M_{\vartheta}^l f(x) = \left(\int_{-\infty}^{\infty} |\nu_t^l * f(x)|^2 dt \right)^{1/2}. \quad (4.16)$$

Thus to show the boundedness of $M_{\mathcal{P}}f$, it suffices to show that

$$\|M_{\mathcal{P}}^l f\|_p \leq C_{p,l} \|f\|_p \quad (4.17)$$

for $p \in ((2+2\alpha)/(1+2\alpha), 2+2\alpha)$, and for all $l = 1, \dots, N$.

To show (4.17), we proceed as follows: let Φ and ψ_t be as in [Section 3](#). Then

$$\begin{aligned} M_{\mathcal{P}}^l f(x) &= \log 2^l \left(\int_{-\infty}^{\infty} \left| \int_{-\infty}^{\infty} v_t^l * \psi_{l(t+u)} * f(x) du \right|^2 dt \right)^{1/2} \\ &\leq \log 2^l \int_{-\infty}^{\infty} S_u^l f(x) du, \end{aligned} \quad (4.18)$$

where

$$S_u^l f(x) = \left(\int_{-\infty}^{\infty} |v_t^l * \psi_{l(t+u)} * f(x)|^2 dt \right)^{1/2}. \quad (4.19)$$

Now by (4.13) and [Theorem 3.1](#), we have

$$\|S_u^l f\|_p \leq C_p \|f\|_p \quad (4.20)$$

for all $p \in (1, \infty)$ and for $l = 1, \dots, N$ which in turn implies that

$$\int_{-1}^1 \|S_u^l f\|_p du \leq 2C_p \|f\|_p \quad \forall p \in (1, \infty). \quad (4.21)$$

On the other hand, if $u \geq 1$, by the estimate (4.11) we have

$$\begin{aligned} \|S_u^l f\|_2^2 &= \int_{\mathbb{R}^d} \int_{-\infty}^{\infty} |v_t^l * \psi_{l(t+u)} * f(x)|^2 dt dx \\ &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^d} (\Phi(2^{lt+lu} \mathbf{L}_l(\xi)))^2 |(\hat{v}_t^l)(\xi)|^2 |\hat{f}(\xi)|^2 d\xi dt \\ &\leq 2^{2l-2lu} \int_{\mathbb{R}^d} |\hat{f}(\xi)|^2 \left(\int_{\log(1/2^{l|\mathbf{L}_l(\xi)|})-u}^{\log(2^{l/|\mathbf{L}_l(\xi)|})-u} dt \right) d\xi \\ &= 2 \log 2^l 2^{l-2lu} \|f\|_2^2. \end{aligned} \quad (4.22)$$

Thus

$$\|S_u^l f\|_2 \leq \sqrt{2 \log 2^l} 2^{l-lu} \|f\|_2. \quad (4.23)$$

By interpolating between (4.20) and (4.23) we get

$$\|S_u^l f\|_p \leq C_{p,l} 2^{\theta l - \theta lu} \|f\|_p \quad (4.24)$$

for all $1 < p < \infty$ and for some $\theta = \theta(p) > 0$. Hence we have

$$\int_1^{\infty} \|S_u^l f\|_p du \leq C_p \|f\|_p \quad \text{for } p \in (1, \infty). \quad (4.25)$$

Finally, if $u < -1$, by the estimate (4.10) and similar argument as in the case of $u \leq 1$, we get

$$\|S_u^l f\|_2 \leq C_l(|u|)^{-1-\alpha} \|f\|_2. \quad (4.26)$$

By interpolating between (4.26) and any $p \in (1, \infty)$ in (4.20), we get that, if $p \in ((2+2\alpha)/(1+2\alpha), 2+2\alpha)$ there exists $\beta > 0$ such that

$$\|S_u^l f\|_p \leq C_p(|u|)^{-\beta} \|f\|_p, \quad (4.27)$$

which implies that

$$\int_{-\infty}^{-1} \|S_u^l f\|_p du \leq C_p \|f\|_p \quad (4.28)$$

for $p \in ((2+2\alpha)/(1+2\alpha), 2+2\alpha)$.

Hence by combining (4.18), (4.21), (4.25), and (4.28) we get (4.17). \square

ACKNOWLEDGEMENT. The publication of this paper was supported by Yarmouk University Research Council.

REFERENCES

- [1] A. Al-Salman and Y. Pan, *Singular integrals with rough kernels*, to appear in Canad. Math. Bull.
- [2] A. Benedek, A. P. Calderón, and R. Panzone, *Convolution operators on Banach space valued functions*, Proc. Nat. Acad. Sci. U.S.A. **48** (1962), 356–365. [MR 24#A3479](#). [Zbl 103.33402](#).
- [3] J. Chen, D. Fan, and Y. Pan, *A note on a Marcinkiewics integral operator*, to appear in Math. Nachr.
- [4] J. Duoandikoetxea and J. L. Rubio de Francia, *Maximal and singular integral operators via Fourier transform estimates*, Invent. Math. **84** (1986), no. 3, 541–561. [MR 87f:42046](#). [Zbl 568.42012](#).
- [5] D. Fan and Y. Pan, *Singular integral operators with rough kernels supported by subvarieties*, Amer. J. Math. **119** (1997), no. 4, 799–839. [MR 99c:42029](#). [Zbl 0899.42002](#).
- [6] L. Grafakos and A. Stefanov, *L^p bounds for singular integrals and maximal singular integrals with rough kernels*, Indiana Univ. Math. J. **47** (1998), no. 2, 455–469. [MR 99i:42019](#). [Zbl 913.42014](#).
- [7] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, New Jersey, 1993. [MR 95c:42002](#). [Zbl 821.42001](#).

HUSSAIN AL-QASSEM: DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN
E-mail address: husseink@yu.edu.jo

AHMAD AL-SALMAN: DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN
E-mail address: alsalman@yu.edu.jo

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk