

ON A THEOREM OF SCHUR

PETER HILTON

(Received 9 January 2001)

To the memory of a dear friend and colleague, Paul Olum

ABSTRACT. We study the ramifications of Schur's theorem that, if G is a group such that G/ZG is finite, then G' is finite, if we restrict attention to nilpotent group. Here ZG is the center of G , and G' is the commutator subgroup. We use localization methods and obtain relativized versions of the main theorems.

2000 Mathematics Subject Classification. 20B07, 20D15.

1. Introduction. The theorem to which we refer is that which asserts that if G is a group and ZG is its center, then

$$G/ZG \text{ finite} \implies G' \text{ finite}, \quad (1.1)$$

where G' is the commutator subgroup of G . This theorem has a nice homological proof, using the 5-term exact sequence

$$H_2 G \xrightarrow{\alpha_4} H_2(G/ZG) \xrightarrow{\alpha_3} ZG \xrightarrow{\alpha_2} G_{ab} \xrightarrow{\alpha_1} (G/ZG)_{ab} \quad (1.2)$$

derived from the short exact sequence $ZG \rightarrow G \rightarrow G/ZG$. For if G/ZG is finite then $H_2(G/ZG)$ is finite. Thus $G' \cap ZG = \ker \alpha_2 = \text{im } \alpha_3$ is finite. But $G'/G' \cap ZG \subseteq G/ZG$ is also finite, so, finally, G' is finite.

We remark that Schur's theorem has a converse which is valid if G is finitely generated (*fg*). We include a proof for completeness.

THEOREM 1.1. *Let G be an *fg* group such that G' is finite. Then G/ZG is finite.*

PROOF. Let $G = \langle x_1, x_2, \dots, x_k \rangle$. Now, for any $x \in G$, there can only be *finitely* many distinct conjugates of x . For there is a one-one correspondence

$$y^{-1}xy \longleftrightarrow x^{-1}y^{-1}xy \quad (1.3)$$

between the set of conjugates of x and a subset of G' ; and G' is finite. Thus $[G : C_G x]$ is finite for all $x \in G$, where $C_G S$ is the centralizer in G of the subset S of G . But if each $[G : C_G x_i]$, $1 \leq i \leq k$, is finite, so is $[G : \cap_i C_G x_i]$. On the other hand, $\cap_i C_G x_i = ZG$, so $[G : ZG]$ is finite. Thus, as claimed, G/ZG is a finite group. \square

Schur's theorem, and its converse, take on a particular significance in the localization theory of nilpotent groups [1]. For it is one of the main problems in that theory to

calculate the *Mislin genus* $\mathcal{G}(G)$ of an $f\mathcal{G}$ nilpotent group G and to identify its members. Here $\mathcal{G}(G)$ is the set of isomorphism classes of $f\mathcal{G}$ nilpotent groups H such that G and H localize at every prime p to isomorphic groups, $G_p \cong H_p$ for all primes p . It is shown in [2, 3] that if G' is finite then $\mathcal{G}(G)$ may itself be given the structure of a (finite) abelian group, a fact which very much facilitates the study of $\mathcal{G}(G)$.

In the category of nilpotent groups (not necessarily $f\mathcal{G}$) it makes sense to consider P -torsion groups, where P is a family of primes, and to study such groups by the techniques of localization. In this way we are able to prove a P -torsion variant of Schur's theorem, namely,

THEOREM 1.2. *Let G be a nilpotent group such that G/ZG is a P -group. Then G' is a P -group.*

We may also prove a converse of [Theorem 1.2](#); as with Schur's theorem itself, it is necessary to impose a supplementary finiteness condition.

THEOREM 1.3. *Let G be a nilpotent group such that G' is a P -group of exponent m . Then G/ZG is a P -group of exponent dividing m^{c-1} , where $\text{nil } G = c$.*

Actually we regard [Theorems 1.2](#) and [1.3](#) as the *absolute* forms of our results and emphasize the *relative* forms which appear to be quite new. In our relativization we replace the group G by a pair (G, N) consisting of a nilpotent group G and a normal subgroup N of G . Then the absolute case is given by $N = G$; moreover, in our relativization, ZG is replaced by $C_G(N)$, which is easily seen to be a normal subgroup of G ; and G' is replaced by the commutator group $[G, N]$.

We remark that [Theorem 1.2](#) also has a variant in which a finiteness condition is imposed just as in [Theorem 1.3](#). Precisely, we have the following theorem.

THEOREM 1.4. *Let G be a nilpotent group such that G/ZG is a P -group of exponent m . Then G' is a P -group of exponent dividing m^{c-1} , where $\text{nil } G = c$.*

We will prove the relativizations of [Theorems 1.2](#), [1.3](#), and [1.4](#) in [Section 2](#). Proofs of the absolute forms, that is, of [Theorems 1.3](#) and [1.4](#) are to be found in [4]. For Warfield proves (the case $n = 1$ is the critical case).

- (a) If Γ_{n+1} has exponent m , then $G/Z_n G$ has exponent dividing m^{c-n} (see [4, Corollary 2.6]); and
- (b) if $G/Z_n G$ has exponent m , then Γ_{n+1} has exponent dividing m^{c-n} (see [4, Corollary 3.16]).

Here we adopt Warfield's convention that $\Gamma_2 = G'$ and $Z_1 = ZG$.

We do not have available to us a homological proof of a relative version of Schur's theorem. However we do show in the appendix how we may use homological arguments to obtain [Theorem 1.4](#) with a small loss of sharpness in our bound on the exponent of G' .

A key tool in our proof of the *relative* version of [Theorem 1.3](#) is a theorem on the localization of nilpotent groups due to Karl Lorenzen ([Theorem 2.6](#)). This theorem is of considerable interest in its own right. It is a pleasure to acknowledge the crucial help the author received from his friend (and erstwhile student) Karl Lorenzen, not only in the provision of [Theorem 2.6](#).

2. Localization methods. Let P be a family of primes and let Q be the complementary family. We first state and prove the relativization of [Theorem 1.2](#).

THEOREM 2.1. *Let G be a nilpotent group and N a normal subgroup such that $G/C_G(N)$ is a P -group. Then $[G, N]$ is a P -group.*

PROOF. Let $e: G \rightarrow G_Q$ localize at the family Q . Now e maps $C_G(N)$ into $C_{G_Q}(N_Q)$; moreover, $C_{G_Q}(N_Q)$ is Q -local. Thus, in fact, the Q -localization $C_G(N)_Q$ of $C_G(N)$ must be a subgroup of $C_{G_Q}(N_Q)$, that is,

$$C_G(N)_Q \subseteq C_{G_Q}(N_Q) \subseteq G_Q. \quad (2.1)$$

Now since $G/C_G(N)$ is a P -group, $(G/C_G(N))_Q = 1$, so that $G_Q = C_G(N)_Q$. Hence, by (2.1) $G_Q = C_{G_Q}(N_Q)$. Thus every element of G_Q commutes with every element of N_Q , so that $[G_Q, N_Q] = 1$. But $[G_Q, N_Q] = [G, N]_Q$, so $[G, N]$ is a P -group. \square

It is clear from this line of proof that, if we want a result in the opposite direction to that of [Theorem 2.1](#), we will have to establish conditions under which

$$C_{G_Q}(N_Q) = C_G(N)_Q. \quad (2.2)$$

Put another way, we ask when the restriction $e_0: C_G(N) \rightarrow C_{G_Q}(N_Q)$ of the Q -localization $e: G \rightarrow G_Q$ itself Q -localizes. Now certainly e_0 is Q -injective and $C_{G_Q}(N_Q)$ is Q -local. Thus e_0 Q -localizes if and only if it is Q -surjective.

In seeking conditions under which e_0 is Q -surjective—and again in proving Lorenzen's theorem ([Theorem 2.6](#)), we need to apply a basic result in [1], namely, Theorem 6.1. We quote that result here as [Lemma 2.2](#).

LEMMA 2.2 (see [1, Theorem 6.1]). *Let G be a nilpotent group with $\text{nil } G = c$ and let $a, b \in G$ with $b^m = 1$. Then $(ab)^{mc} = a^{mc}$.*

However, we can, in fact, refine this result and it will be valuable to do so. Thus we may enunciate

LEMMA 2.3. *If, in addition, $b \in \Gamma^i G$, then $(ab)^{mc-i+1} = a^{mc-i+1}$.*

(Recall that we adopt Warfield's convention for enumerating the terms of the lower central series of G , so that $\Gamma^1 G = G$, $\Gamma^2 G = G'$.)

PROOF OF LEMMA 2.3. We apply [Lemma 2.2](#), but replace G by $\langle a, b \rangle$. However, if $b \in \Gamma^i G$ then $\text{nil} \langle a, b \rangle \leq c - i + 1$. \square

We now apply [Lemma 2.2](#) (we will need the more refined [Lemma 2.3](#) later) to prove the following theorem.

THEOREM 2.4. *Let G, H be nilpotent groups with subgroups $\tilde{G} \subseteq G$, $\tilde{H} \subseteq H$. Let φ be a Q -bijective homomorphism from G to H sending \tilde{G} into \tilde{H} , and let $\tilde{\varphi}: \tilde{G} \rightarrow \tilde{H}$ be obtained by restricting φ . Then $\tilde{\varphi}$ is Q -surjective (and hence Q -bijective) if and only if, for all $x \in G$ such that $\varphi x \in \tilde{H}$, there exists a P -number m such that $x^m \in \tilde{G}$.*

PROOF. We for brevity, describe the property that, for all $x \in G$ such that $\varphi x \in \tilde{H}$, there exists a P -number m such that $x^m \in \tilde{G}$ as *property S*. Suppose

then that $\bar{\varphi}$ is Q -surjective, and let $x \in G$ satisfy $\varphi x \in \bar{H}$. Since $\bar{\varphi}$ is Q -surjective, there exists a P -number n and an element $\bar{x} \in \bar{G}$ such that $\bar{\varphi} \bar{x} = \varphi x^n$. But then $x^n = \bar{x}z$, $z \in G$ with $z^k = 1$ for some P -number k , since φ is Q -injective. Let $\text{nil } G = c$. Then, by [Lemma 2.2](#), $x^{nk^c} = \bar{x}^{k^c} \in \bar{G}$ and nk^c is a P -number, establishing property S .

Suppose, conversely, that property S holds and let $y \in \bar{H}$. Since φ is Q -surjective, there exists a P -number n and $x \in G$ such that $\varphi x = y^n$. Thus, by property S , there exists a P -number m such that $x^m \in \bar{G}$. Then $\bar{\varphi}(x^m) = y^{mn}$ and mn is a P -number, so $\bar{\varphi}$ is Q -surjective. \square

COROLLARY 2.5. *The restriction $e_0 : C_G(N) \rightarrow C_{G_Q}(N_Q)$ Q -localizes if and only if, for all $x \in G$ such that $ex \in C_{G_Q}(N_Q)$, there exists a P -number n such that $x^n \in C_G(N)$.*

This result enables us to exploit the following theorem due to Karl Lorenzen. With G a nilpotent group, N a normal subgroup of G , and $x \in G$, we write $T_P \Gamma_{(x)}^2 N$ for the P -primary component of the torsion subgroup of $\Gamma_{(x)}^2 N$, which is a subgroup of N generated by commutators $[x^r, a]$, $a \in N$. We then prove the following theorem.

THEOREM 2.6 (Lorenzen). *Let $e_0 : C_G(N) \rightarrow C_{G_Q}(N_Q)$ be obtained by restricting the Q -localization $e : G \rightarrow G_Q$. Then e_0 Q -localizes provided that, for all $x \in G$, $T_P \Gamma_{(x)}^2 N$ has finite exponent.*

PROOF. (This is a small but significant modification of Lorenzen's proof, since it exploits [Lemma 2.3](#).) We will apply [Corollary 2.5](#). Thus we must show that, for all $x \in G$ such that $ex \in C_{G_Q}(N_Q)$, there exists a P -number n such that $x^n \in C_G(N)$. Now let $m = \exp T_P \Gamma_{(x)}^2 N$, and let $y \in N$. Then m is a P -number and $e[x, y] = [ex, ey] = 1$, since $ex \in C_{G_Q}(N_Q)$. Hence $[x, y] \in T_P \Gamma_{(x)}^2 N$, so $[x, y]^m = 1$.

Now $x[x, y] = y^{-1}xy$. Hence, by [Lemma 2.3](#), noting that $[x, y] \in \Gamma^2 G$, we conclude that $x^{m^{c-1}} = (y^{-1}xy)^{m^{c-1}} = y^{-1}x^{m^{c-1}}y$, where $\text{nil } G = c$. Since y is an arbitrary element of N , it follows that $x^{m^{c-1}} \in C_G(N)$ and [Theorem 2.6](#) is proved. \square

REMARK 2.7. Notice that it would have sufficed to assume that $T_P \Gamma_{(x)}^2 N$ has finite exponent for all $x \in G$ such that $ex \in C_{G_Q}(N_Q)$.

Lorenzen's theorem is the key to our relativization of [Theorem 1.3](#), which we now state.

THEOREM 2.8. *Let G be a nilpotent group and N a normal subgroup of G . Then if $[G, N]$ is a P -group of exponent m , $G/C_G(N)$ is a P -group of exponent dividing m^{c-1} , where $\text{nil } G = c$.*

PROOF. Since $\Gamma_{(x)}^2 N \subseteq [G, N]$, and $[G, N]$ is a P -group of exponent m , it follows that we have the conditions for applying Lorenzen's theorem, so that $e_0 : C_G(N) \rightarrow C_{G_Q}(N_Q)$ Q -localizes. Now since $[G, N]$ is a P -group, its Q -localization vanishes, that is, $[G_Q, N_Q] = 1$. This means that $G_Q = C_{G_Q}(N_Q)$, so that every $x \in G$ has the property that $ex \in C_{G_Q}(N_Q)$. Moreover, $\exp \Gamma_{(x)}^2 N$ divides m . Thus, following the proof of [Theorem 2.6](#), we see that $x^{m^{c-1}} \in C_G(N)$ for all $x \in G$, so that $\exp(G/C_G(N)) \mid m^{c-1}$. This, of course, implies that $G/C_G(N)$ is a P -group. \square

REMARK 2.9. This last implication follows immediately from $G_Q = C_{G_Q}(N_Q) = C_G(N)_Q$.

It remains to provide the relativization of [Theorem 1.4](#). In fact, we may simply relativize each step in Warfield's argument in [4, Corollary 3.16], thus obtaining the following theorem.

THEOREM 2.10. *Let G be a nilpotent group and N a normal subgroup such that $G/C_G(N)$ is a P -group of exponent m . Then $[G, N]$ is a P -group of exponent dividing m^{c-1} , where $\text{nil } G = c$.*

Appendix

Homological methods. We show in this appendix how homological arguments may be used to obtain [Theorem 1.2](#), although the numerical estimate is marginally inferior to that given by [Theorem 1.4](#). We emphasize that we have only succeeded in developing a homological method in the absolute case.

We begin with a crucial homological lemma.

LEMMA A.11. *Let G be a nilpotent group with $\text{nil } G = c$ and let $n \geq 1$. If G is a torsion group with $\exp G = m$, then $m^{n(c-1)+1} H_n G = 0$.*

PROOF. We argue by induction on c . If $c = 1$, then G is commutative. If K is an arbitrary $f\mathfrak{g}$ subgroup of G , then K is a direct product of (finitely many) finite cyclic groups whose orders divide m , hence $mH_n K = 0$. Now $H_n G = \lim_{\leftarrow} H_n K$, so that $mH_n G = 0$.

Now we assume $c \geq 2$, and assume the lemma proved for nilpotent groups of class $< c$. We consider the central extension

$$\Gamma \longrightarrow G \longrightarrow G/\Gamma, \quad (\text{A.3})$$

where $\Gamma = \Gamma^c G$, and we exploit the Lyndon-Hochschild-Serre spectral sequence associated with (A.3). In this spectral sequence

$$E_{pq}^2 = H_p(G/\Gamma; H_q \Gamma). \quad (\text{A.4})$$

Since the universal coefficient formula in homology splits, and since $\text{nil } \Gamma = 1$, $\text{nil } G/\Gamma = c - 1$, and $\exp \Gamma \mid m$, $\exp(G/\Gamma) \mid m$, it follows from the inductive hypothesis that, if $p + q > 0$,

$$mE_{pq}^2 = 0, \quad q > 0, \quad m^{p(c-2)+1} E_{p0}^2 = 0. \quad (\text{A.5})$$

(The form of writing in (A.5) and in what follows is acceptable since homology groups and E_{pq}^r are additive abelian groups.)

We may then pass to the limit of the spectral sequence, obtaining

$$mE_{pq}^\infty = 0, \quad q > 0, \quad m^{p(c-2)+1} E_{p0}^\infty = 0. \quad (\text{A.6})$$

Now $H_n G$ admits a finite filtration

$$0 = F^{-1} \subseteq F^0 \subseteq \dots \subseteq F^{p-1} \subseteq F^p \subseteq \dots \subseteq F^{n-1} \subseteq F^n = H_n G, \quad (\text{A.7})$$

such that

$$F^p / F^{p-1} = E_\infty^{pq}, \quad p + q = n, \quad 0 \leq p \leq n. \quad (\text{A.8})$$

From (A.6) and (A.8) an easy finite induction shows that

$$m^{p+1}F^p = 0, \quad 0 \leq p \leq n-1. \quad (\text{A.9})$$

Finally, we exploit the short exact sequence

$$F^{n-1} \longrightarrow H_n G \longrightarrow E_{n0}^\infty \quad (\text{A.10})$$

to infer that $m^{n+n(c-2)+1}H_n G = 0$, or $m^{n(c-1)+1}H_n G = 0$, completing the inductive step. \square

Armed with this lemma, we may prove the following theorem.

THEOREM A.12. *Let G a nilpotent group with $\text{nil } G = c$. Then if G/ZG is a torsion group of exponent m , G' is a torsion group of exponent dividing m^{2c-2} .*

PROOF. We exploit the exact sequence (1.2) and the argument used to prove Schur's theorem. Since $\text{nil } G/ZG = c-1$, we know from Lemma A.11 that

$$m^{2(c-2)+1}H_2(G/ZG) = 0. \quad (\text{A.11})$$

Thus

$$m^{2(c-2)+1}(G' \cap ZG) = 0. \quad (\text{A.12})$$

Now $G'/G' \cap ZG \subseteq G/ZG$, so $\exp(G'/G' \cap ZG) \mid m$. Putting this together with (A.12), we deduce finally that G' is a torsion group and $\exp G' \mid m^{2c-2}$. \square

We remark (again) that our estimate of $\exp G'$ is not best possible.

REFERENCES

- [1] P. Hilton, *Localization and cohomology of nilpotent groups*, Math. Z. **132** (1973), 263–286. [MR 48#438](#). [Zbl 0264.20037](#).
- [2] P. Hilton and G. Mislin, *On the genus of a nilpotent group with finite commutator subgroup*, Math. Z. **146** (1976), no. 3, 201–211. [MR 53#615](#). [Zbl 0311.20013](#).
- [3] G. Mislin, *Nilpotent groups with finite commutator subgroups*, Localization in Group Theory and Homotopy Theory, and Related Topics (Sympos., Battelle Seattle Res. Center, Seattle, Wash., 1974), Lecture Notes in Math., vol. 418, Springer, Berlin, 1974, pp. 103–120. [MR 50#10080](#). [Zbl 0302.20029](#).
- [4] R. B. Warfield Jr., *Nilpotent Groups*, Lecture Notes in Mathematics, vol. 513, Springer-Verlag, New York, 1976. [MR 53#13413](#). [Zbl 0347.20018](#).

PETER HILTON: DEPARTMENT OF MATHEMATICAL SCIENCES, SUNY BINGHAMTON, BINGHAMTON, NY 13902-6000, USA

Current address: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CENTRAL FLORIDA, ORLANDO, FL 32816-1364, USA

E-mail address: marge@math.binghamton.edu

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br