

RESEARCH NOTES

ALTERNATIVE INTEGRATION PROCEDURE FOR SCALE-INVARIANT ORDINARY DIFFERENTIAL EQUATIONS

GERALD ROSEN

Department of Physics
Drexel University
Philadelphia, Pennsylvania 19104
U.S.A.

(Received December 28, 1978)

ABSTRACT. For an ordinary differential equation invariant under a one-parameter group of scale transformations $x \rightarrow \lambda x$, $y \rightarrow \lambda^\alpha y$, $y' \rightarrow \lambda^{\alpha-1} y'$, $y'' \rightarrow \lambda^{\alpha-2} y''$, etc., it is shown by example that an explicit analytical general solution may be obtainable in parametric form in terms of the scale-invariant variable

$\xi \equiv \int^x y^{-1/\alpha} dx$. This alternative integration may go through, as it does for the example equation $y'' = kxy^{-2}y'$, in cases for which the customary dependent and independent variables ($x^{-\alpha}y$) and ($\ln x$) do not yield an analytically integrable transformed equation.

KEY WORDS AND PHRASES. Ordinary differential equations, Integration method, Scale-invariance.

AMS (MOS) SUBJECT CLASSIFICATION (1970) CODES. 34A05.

1. INTRODUCTION.

Consider an ordinary differential equation

$$F(x, y, y', y'', \dots) = 0 \quad (y' \equiv dy/dx, y'' \equiv d^2y/dx^2, \dots) \quad (1.1)$$

that is invariant under a one-parameter group of scale transformations with $F(\lambda x, \lambda^\alpha y, \lambda^{\alpha-1} y', \lambda^{\alpha-2} y'', \dots) \equiv \lambda^\beta F(x, y, y', y'', \dots)$ for certain prescribed real constants α, β and all real $\lambda > 0$. The customary procedure [1] for reducing the order of such an equation, and thus commonly facilitating its solution, involves the introduction of new dependent and independent variables $\eta \equiv x^{-\alpha} y$ and $\xi \equiv \ln x$. For example, in the case of the scale-invariant ($\alpha = 1, \beta = 1$) nonlinear second-order equation.

$$y^2 y'' - kxy' = 0 \quad (k = \text{const}) \quad (1.2)$$

or, equivalently, its inverted form $y^2 (d^2x/dy^2) = -kx(dx/dy)^2$ with $x = x(y)$, introduction of the variables $\xi \equiv \ln x, \eta \equiv x^{-1}y = \eta(\xi)$ yields [1]

$$\eta^2 \frac{d^2\eta}{d\xi^2} + (\eta^2 - k) \frac{d\eta}{d\xi} - k\eta = 0, \quad (1.3)$$

which is equivalent to a first-order equation in $p \equiv d\eta/d\xi = p(\eta)$; however, there is no analytical method for obtaining an integrating factor for the resulting first-order equation, and thus the customary procedure is ineffectual for solving (1.2) by analytical means.

In such cases for which the scale-invariant equation is not amenable to solution by the customary method, an analytical general solution may possibly be obtainable in parametric form in terms of the scale-invariant variable

$$\zeta \equiv \int_{\text{const}}^x y^{-1/\alpha} dx. \quad (1.4)$$

To see how this works, consider the $\alpha = 1$ equation (1.2) with $x = x(\zeta)$ and

$y = dx/d\zeta \equiv x$, according to (1.4). Then $y' = (dx/d\zeta)^{-1}(dy/d\zeta) \equiv y^{-1}\dot{y}$ and (1.2) becomes

$$y \dot{y}^{-1} \ddot{y} - \dot{y} - kx = 0. \quad (1.5)$$

Differentiating (1.5) with respect to ζ and dividing out a common factor of y yields

$$\dot{y}^{-1} \ddot{\ddot{y}} - \dot{y}^{-2} (\ddot{y})^2 - k = 0 \quad (1.6)$$

with the immediate integral

$$\dot{y}^{-1} \ddot{y} = k\zeta. \quad (1.7)$$

A trivial constant of integration, associated with the arbitrariness in the constant lower limit of integration in (1.4), is omitted from the right side of (1.7). Integrating the latter equation twice yields

$$y = A \int_0^{\zeta} (\exp \frac{1}{2} kv^2) dv + B \quad (A, B = \text{const}). \quad (1.8)$$

Finally, by putting (1.8) into the integral formula $x = \int y d\zeta$, one obtains

$$x = A \left[\zeta \int_0^{\zeta} (\exp \frac{1}{2} kv^2) dv - k^{-1} (\exp \frac{1}{2} k\zeta^2) \right] + B\zeta. \quad (1.9)$$

Since there is no way of getting $y = y(x)$ explicitly by eliminating ζ from the general solution to (1.2) given parametrically by (1.8) and (1.9), it is clear why the customary solutional procedure based on (1.3) is ineffectual. The potency of this alternative integration procedure stems in part from the greater generality of parametric representations.

ACKNOWLEDGEMENT. This work was supported by NASA grant NSG 3090.

REFERENCE

1. Kamke, E., Differentialgleichungen Lösungsmethoden und Lösungen, Akademische-Verlag., Leipzig, 1967 (also Chelsea Publishing Co., N.Y., 1959) 68, No. 6.58 on p. 555, No. 6.87 on p. 565.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br