

## RESEARCH NOTES

### ON SLOW OSCILLATION AND NONOSCILLATION IN RETARDED EQUATIONS

BHAGAT SINGH

Department of Mathematics  
University of Wisconsin Center  
Manitowoc, Wisconsin 54220  
U.S.A.

(Received September 11, 1978 and in revised form June 10, 1979)

ABSTRACT. Sufficient conditions have been found to ensure that all oscillatory solutions of

$$(r(t)y'(t))' + a(t)y(t-\zeta(t)) = f(t)$$

are slowly oscillating. This behaviour is further linked to nonoscillation.

KEY WORDS AND PHRASES. Oscillation, Nonoscillation, Moderately Oscillating, Slowly Oscillating.

AMS (MOS) SUBJECT CLASSIFICATION (1970) CODES. 34C10.

#### 1. INTRODUCTION.

In studying the asymptotic nature of oscillatory solutions of the equation

$$(r(t)y'(t))' + a(t)y(t-\zeta(t)) = f(t), \quad (1)$$

this author in [9, Theo. 2] showed that a nontrivial oscillatory solution  $y(t)$  of (1) satisfies  $\lim_{t \rightarrow \infty} y(t) = 0$  if  $\int^{\infty} |a(t)| dt < \infty$ ,  $\int^{\infty} |f(t)| dt < \infty$  and  $\int^{\infty} \frac{1}{r} dt < \infty$ .

In theorem 4 in [9], it was observed that  $\int_1^\infty \frac{1}{r} dt = \infty$  lead to slowly oscillating solutions which do not approach zero as  $t \rightarrow \infty$ .

In this work, we give sufficient conditions which cause all oscillatory solutions of the equation

$$(r(t)y'(t))' + a(t)h(y(g(t))) = f(t) \quad (2)$$

to oscillate slowly.

Even though voluminous literature exists about various types of oscillatory and nonoscillatory criteria for such equations, the asymptotic nature of oscillatory solutions of these equations has not been so extensively studied. For a good literature study of related results Graef [1] and Graef and Spike [2] have included an exhaustive reference list. The work of Travis [11] (c.f[8]) shows that common techniques found for ordinary differential equations fail on retarded differential equations even when the retardations are small. For oscillation criteria see T. Kusano and H. Onose [4] and this author [7].

## 2. ASSUMPTIONS AND DEFINITIONS

The entire study in this work is subject to the following assumptions:

- (i)  $a(t)$ ,  $r(t)$ ,  $f(t)$ ,  $h(t)$  and  $g(t)$  are  $C^0(\mathbb{R})$ , where  $\mathbb{R}$  is the real line,
- (ii)  $r(t) > 0$ ,  $g(t) > 0$ ,  $g'(t) > 0$ ;
- (iii)  $g(t) \leq t$  and  $g(t) \rightarrow \infty$  as  $t \rightarrow \infty$ ;
- (iv)  $h$  is odd,  $\text{sign}(h(t)) = \text{sign}(t)$  and there exists positive constants  $\alpha, m$  such that  $0 < \alpha \leq \frac{h(t)}{t} \leq m$  on some positive half line.

In order to reckon the half line we shall assume that it would mean for  $t \geq t_0$  for some positive  $t_0$ .  $t_0$  will be referred to without further mention. All functions considered are real valued.

We call a function  $H(t) \in C^0[t_0, \infty)$  oscillatory if  $H(t)$  has arbitrarily large zeros in  $[t_0, \infty)$ ; otherwise we call it nonoscillatory. In order to be more precise we shall use the term "solution" only for those nontrivial solutions (of equations under consideration) which can be extended continuously for  $t \geq t_0$ .

We define a function  $H_1(t) \in C^0[t_0, \infty)$  to be slowly oscillating if  $H_1(t)$  is oscillatory and the set:  $Z_{H_1} = \{y_0 - x_0 : y_0 > x_0 ; y_0 \text{ and } x_0 \text{ are consecutive zeros of } H_1(t), x_0 > t_0\}$  is unbounded. If  $Z_{H_1}$  is bounded,  $H_1$  is called "moderately oscillating".

### 3. ON SLOW OSCILLATION

Theorem 4 in [9] states that moderately oscillatory solutions of (1) approach zero if  $\int^{\infty} |a(t)| dt < \infty$  and  $\int^{\infty} |f(t)| dt < \infty$ . Our next Theorem gives sufficient conditions for all oscillatory solutions of (2) to be slowly oscillating.

THEOREM (1). Suppose  $a(t) = a_1(t) + a_2(t)$  where  $a_1(t) > 0$  and  $\frac{a_2(t)}{a_1(t)}$  is bounded for large  $t$ . Further suppose that  $\int^{\infty} |f(t)| dt < \infty$  and  $\liminf_{t \rightarrow \infty} \frac{|f(t)|}{a_1(t)} > 0$ . Then all oscillatory solutions  $y(t)$  of equation (2) satisfy

$$\limsup_{t \rightarrow \infty} |y(t)| > 0 \quad (3)$$

and

$$y(t) \text{ is slowly oscillating.} \quad (4)$$

PROOF. Let  $y(t)$  be an oscillatory solution of (2). Rewriting (2) we have

$$\frac{(ry')'}{a_1(t)} + h(y(g(t))) + \frac{a_2(t)}{a_1(t)} h(y(g(t))) = \frac{f(t)}{a_1(t)} \quad . \quad (5)$$

Suppose to the contrary that  $\lim_{t \rightarrow \infty} y(t) = 0$ . Since  $\frac{a_2(t)}{a_1(t)}$  is bounded and

$h(x) \rightarrow 0$  as  $x \rightarrow 0$  we get that

$(1 + a_2(t)/a_1(t)) h(y(g(t))) \rightarrow 0$  as  $t \rightarrow \infty$ . Since  $\liminf_{t \rightarrow \infty} (|f(t)| / a_1(t)) > 0$ , (5) reveals that  $y'(t)$  assumes a constant

sign making  $y(t)$  nonoscillatory. This contradiction shows that

$$\limsup_{t \rightarrow \infty} |y(t)| > 0.$$

Now

$$\begin{aligned} \int_T^\infty |a(t)| dt &\leq \int_T^\infty a_1(t) dt + \int_T^\infty |a_2(t)| dt \\ &= \int_T^\infty a_1(t) dt + \int_T^\infty \frac{|a_2(t)|}{a_1(t)} a_1(t) dt \\ &< \infty, \end{aligned}$$

since  $\frac{a_2(t)}{a_1(t)}$  is bounded as  $t \rightarrow \infty$  and  $\int_T^\infty |f(t)| dt < \infty$ . By Theorem

4 of this author [9],  $y(t)$  is slowly oscillating. The proof is now complete.

EXAMPLE (1). Consider the equation.

$$y''(t) + \frac{5}{4t^2} y(t) = \frac{1}{t^{3/2}}, \quad t > 0. \quad (6)$$

Here  $a_1(t) \equiv a(t) = \frac{5}{4t^2}$ . All conditions of Theorem 3 are satisfied. Hence

all oscillatory solutions of (6) are slowly oscillating and satisfy

$\limsup_{t \rightarrow \infty} |y(t)| > 0$ . In fact  $y(t) = \sqrt{t} (1 + 2 \sin(\lambda \ln t))$  is one such

solution.

EXAMPLE (2). Let  $y(t)$  be a solution of

$$y''(t) + \frac{1 + 2 \sin t}{t^3} y(t - \pi) = \frac{2}{t^3} + \frac{\cos t}{t^4}, \quad t > \pi. \quad (7)$$

Taking  $a_1(t) = 1/t^3$ ,  $a_2(t) = 2 \sin t / t^3$  we find that all conditions of this theorem are satisfied. Hence either  $y(t)$  is nonoscillatory or slowly oscillating with no limit at  $\infty$ .

Example 1 suggests the following theorem:

THEOREM (2). Suppose  $a(t) = a_1(t) + a_2(t)$  where  $a_1(t) > 0$  and  $a_2(t) / a_1(t)$  is bounded for large  $t$ . Further suppose that

$\int_{-\infty}^{\infty} |f(t)| dt < \infty$  and  $\frac{|f(t)|}{a_1(t)} \rightarrow \infty$  as  $t \rightarrow \infty$ . Let  $y(t)$  be an oscillatory solution of (2). Then  $y(t)$  is slowly oscillating and  $\limsup_{t \rightarrow \infty} |y(t)| = \infty$ .

PROOF. We only need to show that  $\limsup_{t \rightarrow \infty} |y(t)| = \infty$ . Suppose to the

contrary that  $y(t)$  is bounded. Then (5) immediately reveals that  $(ry')'/a_1(t) \rightarrow \infty$  as  $t \rightarrow \infty$ .

Since  $a_1(t) > 0$ ,  $|y'(t)| > 0$  eventually. This forces  $y(t)$  to be nonoscillatory, a contradiction. This proves the theorem.

EXAMPLE (3). The equation

$$y''(t) + \left( \frac{5}{4t^2} + \frac{6 \sin(\ln t)}{t^4} \right) y(t) = \frac{1}{t^{3/2}} + \frac{6}{t^{7/2}} \left( \sin(\ln t) + 2 \sin^2(\ln t) \right), \quad (8)$$

$t > 0$ ,

satisfies all conditions of Theorem 2 by choosing  $a_1$  and  $a_2$  as

$$a_1(t) = \frac{5}{4t^2} \text{ and } a_2(t) = \frac{6 \sin(\ln t)}{t^4}. \text{ This equation has}$$

$$y(t) = \sqrt{t} (1 + 2 \sin(\ln t))$$

as an oscillatory solution satisfying the conclusion of this theorem.

Our next theorem gives conditions for boundedness of all moderately oscillating solutions of equation 2. It is well known that if  $a(t) = a_1(t) + a_2(t)$  where  $a_1(t) \rightarrow L > 0$  as  $t \rightarrow \infty$ ;  $a_1(t)$  is of bounded variation and

$$\int_{-\infty}^{\infty} |a(t)| dt < \infty \text{ then all solutions of}$$

$$y''(t) + a(t)y(t) = 0 \quad (9)$$

are bounded with bounded derivatives. See Cesari [3, p.85]. It is not true for retarded equations

$$y''(t) + a(t)h(y(g(t))) = 0 \quad (10)$$

as the following example shows.

EXAMPLE (4). The equation

$$y'' + 2e^{\pi/2}y(t - \pi/2) = 0 \quad (11)$$

has  $y(t) = e^t \sin t$  as a solution. However our next theorem gives the partial result. We prove it in more generality for equation 2.

THEOREM (3). Suppose  $a(t) = a_1(t) + a_2(t)$  where  $a_1(t) \rightarrow L > 0$  as  $t \rightarrow \infty$ ,  $\int_1^\infty |a_2(t)| dt < \infty$  and  $\int_1^\infty |f(t)| dt < \infty$ . Further suppose that there exists a  $\lambda > 0$  such that  $L + \lambda < (4 - \lambda)/\alpha^2$  where for any oscillatory solution  $y$  of (31),  $\alpha = \sup \{x_2 - x_1 : y(x_2) = y(x_1) = 0, y(x) \neq 0, x \in (x_1, x_2)\} < \infty$  then  $y(t)$  is bounded.

PROOF. Let  $T$  be large enough so that for  $t > T$ ,  $a_1(t) \leq L + \lambda$ . Let  $T_1 > T$  be large enough so that  $g(T_1) > T$  and  $y(T_1) = 0$ ,  $\int_T^\infty |a_2(t)| dt < \lambda/2$  and  $\int_T^\infty |f(t)| dt < \lambda/2$ .

Suppose to the contrary that  $\limsup_{t \rightarrow \infty} |y(t)| = \infty$ . Then there is a  $T_1' > T_2 > T_1$  such that  $y(T_2) = 0$  and  $P_1 = \max \{ |y(t)| : T \leq t \leq T_1' \} = |y(T_1')| > \lambda$ .  $y(x_1) = y(x_2) = 0$ . Let  $M_1 = \max \{ |y(t)| : x_1 \leq t \leq x_2 \}$ , where  $[x_1, x_2]$  is the smallest closed interval containing  $T_1'$ . It is clear that  $T_2 \leq x_1$ ,

$$|y(t)| \leq M_1 \quad (12)$$

and

$$|y(g(t))| \leq M_1 \quad (13)$$

for  $t \in [x_1, x_2]$ . Let  $x_0 \in [x_1, x_2]$  such that  $M_1 = |y(x_0)|$ . Since

$$M_1 = \int_{x_1}^{x_0} y'(t) dt = - \int_{x_2}^{x_0} y'(t) dt$$

we have

$$2M_1 \leq \int_{x_1}^{x_2} |y'(t)| dt$$

$$= \int_{x_1}^{x_2} |y'(t)|^{\frac{1}{2}} |y''(t)|^{\frac{1}{2}} dt$$

$$4M_1^2 \leq \left( \int_{x_1}^{x_2} dt \right) \left( \int_{x_1}^{x_2} y'(t) \cdot y''(t) dt \right) .$$

by Schwarz's inequality. Integrating by parts and using (2) we get

$$4M_1^2 \leq (x_2 - x_1) \left[ \int_{x_1}^{x_2} y(t) a(t) h(y(g(t))) dt - \int_{x_1}^{x_2} y(t) f(t) dt \right] . \quad (14)$$

From (14) we have

$$4M_1^2 \leq \alpha \int_{x_1}^{x_2} a(t) \frac{h(y(g(t)))}{y(g(t))} y(g(t)) dt + \alpha \int_{x_1}^{x_2} |f(t)| dt$$

since  $x_2 - x_1 \leq \alpha$ . This yields in view of (12) and (13)

$$4 \leq \alpha m \int_{x_1}^{x_2} (a_1(t) + |a_2(t)|) dt + \frac{2}{M_1} \int_{x_1}^{x_2} |f(t)| dt \quad (15)$$

which gives

$$4 \leq \alpha^2 m (L + \lambda) + \alpha m \int_{x_1}^{x_2} |a_2(t)| dt + \frac{\alpha}{M_1} \int_{x_1}^{x_2} |f(t)| dt \quad (16)$$

or

$$4 \leq \alpha^2 m (L + \lambda) + \lambda , \quad (17)$$

$$\text{since } \frac{\alpha}{M_1} < 1 , \alpha m \int_{x_1}^{x_2} |a_2(t)| dt < \lambda/2 \text{ and } \int_{x_1}^{x_2} |f(t)| dt \leq \lambda/2 .$$

This contradiction, in (17), completes the proof.

REMARK. Coming back to example (4) we see that for the solution  $y = e^t \sin t$ ,  $\alpha = \pi$ ,  $L = 2e^{\pi/2}$ ,  $m = 1$ ,  $f(t) \equiv 0$  and  $a_2(t) \equiv 0$ . Thus far any  $\lambda > 0$ ,  $\alpha^2 m (L + \lambda) = \pi^2 (2 e^{\pi/2} + \lambda) > 4 - \lambda$ , satisfying the conclusion of the theorem.

EXAMPLE (5). The equation

$$y''(t) + 2 e^{-3\pi/2} y(t - \frac{3\pi}{2}) = 0 \quad (18)$$

has  $y = e^{-t} \sin t$  as a solution. Here  $\alpha = \pi$   
 $L = e^{-\frac{3\pi}{2}} \approx e^{-4.7}$ ,  $m = 1$ .

Therefore for  $\lambda = .01$

$$m\alpha^2(L + \lambda) = \pi^2(e^{-4.7} + .01) < 4, \text{ once again, satisfying this theorem.}$$

#### REFERENCES

1. Graef, John R. Oscillation, Nonoscillation and Growth of Solutions of Nonlinear Functional Differential Equations of Arbitrary Order, J. Math. Anal. Appl. 60 (1977) 398-409.
2. Graef, John R. and Paul W. Spikes. Asymptotic Properties of Solutions of Functional Differential Equations of Arbitrary Order, J. Math. Anal. Appl. 60 (1977) 339-348.
3. Cesari, L. "Asymptotic Behavior and Stability Problems in Ordinary Differential Equations," Springer-Verlag, Berlin, 1959.
4. Kusano, T. and H. Onose. Oscillations of Functional Differential Equations with Retarded Arguments, J. Differential Equations, 15 (1974) 269-277.
5. Londen, S. Some Nonoscillation Theorems for a Second Order Nonlinear Differential Equation, SIAM J. Math. Anal. 4 (1973) 460-465.
6. Norkin, S.B. "Differential Equations of Second Order with Retarded Arguments," American Mathematical Society, Providence, Rhode Island (1972) 1-8.
7. Singh, B. Oscillation and Nonoscillation of Even-order Nonlinear Delay-differential Equations, Quart. Appl. Math. (1973) 343-349.
8. Singh, B. Asymptotic Nature of Nonoscillatory Solutions of nth Order Retarded Differential Equations, SIAM J. Math. Anal. 6 (1975) 784-795.
9. Singh, B. Asymptotically Vanishing Oscillatory Trajectories in Second Order Regarded Equations, SIAM J. Math. Anal. 7 (1976) 37-44.
10. Singh, B. Forced Oscillations in Second Order Functional Equations, Hiroshima Math. J. 7 (1977) to appear.
11. Travis, C.C. Oscillation Theorems for Second Order Differential Equations with Functional Arguments, Proc. Amer. Math. Soc. 30 (1972) 199-201.
12. Wallgren, T. Oscillation of Solutions of the Differential Equation  $y'' + p(x)y = f(x)$ , SIAM J. Math. Anal. 7 (1976) 848-857.

# Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

### Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

### Lead Guest Editor

**Juan J. Nieto**, Departamento de Análisis Matemático,  
Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;  
[juanjose.nieto.roig@usc.es](mailto:juanjose.nieto.roig@usc.es)

### Guest Editor

**Donal O'Regan**, Department of Mathematics, National  
University of Ireland, Galway, Ireland;  
[donal.oregan@nuigalway.ie](mailto:donal.oregan@nuigalway.ie)