

THE DE RHAM THEOREM FOR THE NONCOMMUTATIVE COMPLEX OF CENKL AND PORTER

LUIS FERNANDO MEJIAS

Received 22 May 2001

We use noncommutative differential forms (which were first introduced by Connes) to construct a noncommutative version of the complex of Cenkl and Porter $\Omega^{*,*}(X)$ for a simplicial set X . The algebra $\Omega^{*,*}(X)$ is a differential graded algebra with a filtration $\Omega^{*,q}(X) \subset \Omega^{*,q+1}(X)$, such that $\Omega^{*,q}(X)$ is a \mathbb{Q}_q -module, where $\mathbb{Q}_0 = \mathbb{Q}_1 = \mathbb{Z}$ and $\mathbb{Q}_q = \mathbb{Z}[1/2, \dots, 1/q]$ for $q > 1$. Then we use noncommutative versions of the Poincaré lemma and Stokes' theorem to prove the noncommutative tame de Rham theorem: if X is a simplicial set of finite type, then for each $q \geq 1$ and any \mathbb{Q}_q -module M , integration of forms induces a natural isomorphism of \mathbb{Q}_q -modules $I : H^i(\Omega^{*,q}(X), M) \rightarrow H^i(X; M)$ for all $i \geq 0$. Next, we introduce a complex of noncommutative tame de Rham currents $\Omega_{*,*}(X)$ and we prove the noncommutative tame de Rham theorem for homology: if X is a simplicial set of finite type, then for each $q \geq 1$ and any \mathbb{Q}_q -module M , there is a natural isomorphism of \mathbb{Q}_q -modules $I : H_i(X; M) \rightarrow H_i(\Omega_{*,q}(X), M)$ for all $i \geq 0$.

2000 Mathematics Subject Classification: 55N35.

1. Introduction. Sullivan [22] used the de Rham complex $\Omega^*(X, \mathbb{Q})$ for a simplicial space X of Whitney [23], to construct a free algebra model $M(X)$ of Ω^* such that the rational homotopy of X could be computed. In an effort to use the idea of a “model” to compute homotopy group Miller [18] and Cartan [2] constructed a filtration $\mathcal{A}^*(X, \mathbb{Z})$ of the de Rham complex of polynomial forms $\mathcal{A}^*(X, \mathbb{Q})$ such that the cohomology $H^i(\mathcal{A}^{*,q}(X; \mathbb{Z}))$ is isomorphic to the singular cohomology $H^i(X; \mathbb{Z})$ for $i \leq q$. It turned out that a free model for \mathcal{A}^* could not be constructed.

Cenkl and Porter [6, 7] constructed the so-called tame de Rham complex of polynomial forms $T^*(X, \mathbb{Z})$ with filtration $T^{*,q}(X) \subset T^{*,q+1}(X)$, depending on the degree of the polynomials and the forms, such that $T^{*,q}(X)$ is a \mathbb{Q}_q -module ($\mathbb{Q}_0 = \mathbb{Q}_1 = \mathbb{Z}$ and $\mathbb{Q}_q = \mathbb{Z}[1/2, \dots, 1/q]$ for $q > 1$). They proved that there exists an isomorphism $I : H^i(T^{*,q}(X; \mathbb{Q}_q)) \rightarrow H^i(X; \mathbb{Q}_q)$ which is induced by integration of forms for all $q \geq 1$ and for all i . They also constructed a free model $TM(X)$ for $T(X, \mathbb{Z})$ which computes most of the torsion in the homotopy groups of X . However, the Steenrod operations cannot be introduced on $TM(X)$ (with proper localization).

The proof of Cenkl and Porter of the de Rham theorem for $T^*(X, \mathbb{Z})$ was done for a finite simplicial complex. This was later extended for a simplicial set by Boullay et al. [1]. They also dualized the situation and proved a homology version of the tame de Rham theorem [20]. Several attempts were made to build free models for spaces with \mathbb{Z}_p -coefficients (see [10, 13, 19]).

Trying to construct a model for a space (along the lines of Sullivan, etc.) that would have cohomology operations, Karoubi [11] proposed enlarging the de Rham complex of commutative forms by considering noncommutative algebras (along the lines of Connes). Karoubi defined a noncommutative de Rham complex $\Omega(X)$ and proved the noncommutative de Rham theorem for a simplicial space X . A slightly more general version of the noncommutative de Rham theorem was proved by Cenkl in [4, 6]. Both proofs are functorial and in principle are based on the idea of Cartan. Some Steenrod operations are induced.

In this paper, we present another proof of the noncommutative de Rham theorem for a simplicial set of finite type. This proof is in the spirit of the classical de Rham theorem, that is, using integration. The possibility of such a proof (over a ring containing \mathbb{Q}) was suggested by Karoubi in [12]. However, we give a stronger result by defining a noncommutative tame de Rham complex $\Omega^{*,*}(X)$ (a noncommutative version of the de Rham complex of Cenkl and Porter).

THEOREM 1.1. *Let X be a simplicial set of finite type. Then for each $q \geq 1$ and any \mathbb{Q}_q -module M , there is a natural isomorphism of \mathbb{Q}_q -modules*

$$H^i(\Omega^{*,q}(X), M) \xrightarrow{\cong} H^i(X; M) \quad \forall i \geq 0. \quad (1.1)$$

The isomorphism is induced by integration.

Motivated by the work of Scheerer et al. [20], we also introduce a complex of noncommutative tame currents $\Omega_{*,*}(X)$ (dual of noncommutative tame forms) and prove the noncommutative tame de Rham theorem for homology.

THEOREM 1.2. *Let X be a simplicial set of finite type. Then for each $q \geq 1$ and any \mathbb{Q}_q -module M , there is a natural isomorphism of \mathbb{Q}_q -modules*

$$H_i(X; M) \xrightarrow{\cong} H_i(\Omega_{*,q}(X), M) \quad \forall i \geq 0. \quad (1.2)$$

The isomorphism is induced by integration.

Finally, we introduce a noncommutative version of the complex of tame currents presented in [20] and compare it with the complex $\Omega_{*,*}(X)$.

2. Simplicial objects. In this section, we present a brief introduction to the concept of simplicial objects as well as some examples. For more details see [15, 16, 17].

A *simplicial set* X is a graded set indexed on the nonnegative integers together with the *face operators* $d_i : X_k \rightarrow X_{k-1}$ and the *degeneracy operators* $s_i : X_k \rightarrow X_{k+1}$, $0 \leq i \leq k$, which satisfy the following identities:

- (i) $d_i d_{j+1} = d_j d_i$ if $i \leq j$,
- (ii) $s_i s_j = s_{j+1} s_i$ if $i \leq j$,
- (iii) $d_i s_j = s_{j-1} d_i$ if $i < j$,
- (iv) $d_j s_j = \text{identity} = d_{j+1} s_j$,
- (v) $d_i s_j = s_j d_{i-1}$ if $i > j+1$.

The elements of X_k are called *k-simplices*. Let X and Y be two simplicial sets. A *simplicial map* $f : X \rightarrow Y$ is a map of graded sets of degree zero which commutes with the face and degeneracy operators.

If X and Y are two simplicial sets, the *Cartesian product* $X \times Y$ is the simplicial set with $(X \times Y)_k = X_k \times Y_k$ and

$$d_i(x, y) = (d_i x, d_i y), \quad s_i(x, y) = (s_i x, s_i y), \quad \forall x \in X_k, y \in Y_k, 0 \leq i \leq k. \quad (2.1)$$

EXAMPLE 2.1. Let V be any partially ordered set. Let X_k be the set of all finite sequences (x_0, \dots, x_k) , with $x_0 \leq \dots \leq x_k$, $x_0, \dots, x_k \in V$. Define $d_i : X_k \rightarrow X_{k-1}$ and $s_i : X_k \rightarrow X_{k+1}$, $0 \leq i \leq k$, by

$$\begin{aligned} d_i(x_0, \dots, x_k) &= (x_0, \dots, x_{i-1}, \hat{x}_i, x_{i+1}, \dots, x_k) \quad (\text{omit } x_i), \\ s_i(x_0, \dots, x_k) &= (x_0, \dots, x_{i-1}, x_i, x_i, x_{i+1}, \dots, x_k) \quad (\text{double } x_i). \end{aligned} \quad (2.2)$$

Then $X = \{X_k\}$ is a simplicial set.

EXAMPLE 2.2. Let Δ denote the category whose objects are all finite sequences of integers $\Delta(n) = \{0, 1, \dots, n\}$ and the morphisms are all the increasing functions $f : \Delta(n) \rightarrow \Delta(m)$ (for all $0 \leq i \leq j \leq n$, we have $f(i) \leq f(j)$).

Define the morphisms $\delta_i : \Delta(n-1) \rightarrow \Delta(n)$ and $\sigma_i : \Delta(n+1) \rightarrow \Delta(n)$, for $0 \leq i \leq n$, by

$$\delta_i(j) = \begin{cases} j & \text{if } j < i, \\ j+1 & \text{if } j \geq i, \end{cases} \quad \sigma_i(j) = \begin{cases} j & \text{if } j \leq i, \\ j-1 & \text{if } j > i. \end{cases} \quad (2.3)$$

Then every $f \in \text{Hom}(\Delta(n), \Delta(m))$ can be written as the product of finitely many δ 's and σ 's.

A *simplicial object in a category* \mathcal{C} is a contravariant functor $F : \Delta \rightarrow \mathcal{C}$. A simplicial set X can be identified with a simplicial object X in the category of sets **Set**, $F : \Delta \rightarrow \mathbf{Set}$, $X = F(\Delta_n) = X$ (see [15, page 233] or [17, page 4]).

A *simplicial Λ -module* is a simplicial object in the category of Λ -modules **Mod**. If M and N are simplicial Λ -modules, then the tensor product $M \otimes N$ is a simplicial Λ -module. The face and degeneracy operators d_i and s_i on $(M \otimes N)_k = M_k \otimes N_k$ are given by

$$d_i(x \otimes y) = d_i x \otimes d_i y, \quad s_i(x \otimes y) = s_i x \otimes s_i y, \quad \forall x \in X_k, y \in Y_k, 0 \leq i \leq k. \quad (2.4)$$



A *simplicial graded algebra* $\mathcal{A}^* = \oplus_{n \geq 0} \mathcal{A}^n$ is a family of graded algebras $\mathcal{A}_k^* = \oplus_{n \geq 0} \mathcal{A}_k^n$, $k = 0, 1, 2, \dots$, over a commutative ring Λ which is a simplicial set and the face and degeneracy operators d_i and s_i are morphisms of graded algebras.

EXAMPLE 2.3. Let $\Delta_n = \{(a_0, \dots, a_n) \in \mathbb{R}^{n+1} \mid 0 \leq a_i \leq 1, \sum a_i = 1\}$ be the *standard n -simplex* (Figure 2.1). The maps $\delta_i : \Delta_{n-1} \rightarrow \Delta_n$ and $\sigma_i : \Delta_{n+1} \rightarrow \Delta_n$ are defined by

$$\begin{aligned} \delta_i(x_0, \dots, x_{n-1}) &= (x_0, \dots, x_{i-1}, 0, x_i, \dots, x_{n-1}), \\ \sigma_i(x_0, \dots, x_{n+1}) &= (x_0, \dots, x_{i-1}, x_i + x_{i+1}, x_{i+2}, \dots, x_{n+1}). \end{aligned} \quad (2.5)$$

Let \mathcal{P}_n be the collection of the polynomials $f : \Delta_n \rightarrow \mathbb{R}$ with \mathbb{Z} -coefficients and let $\mathcal{P} = \{\mathcal{P}_n\}_{n \geq 0}$. Then \mathcal{P} is a simplicial set. The face and degeneracy maps are the maps $\delta_i : \mathcal{P}_n \rightarrow \mathcal{P}_{n-1}$ and $s_i : \mathcal{P}_n \rightarrow \mathcal{P}_{n+1}$ defined, for each $f \in \mathcal{P}_n$, by

$$\delta_i(f) = f \circ \delta_i, \quad s_i(f) = f \circ \sigma_i, \quad (2.6)$$

FIGURE 2.1. The standard simplexes Δ_1 and Δ_2 .FIGURE 2.2. The 2-simplexes Δ_2 .

(δ_i and s_i are the pullbacks of δ_i and σ_i). Multiplication of polynomials induces an algebra structure on \mathcal{P} . Then \mathcal{P} is a simplicial algebra.

EXAMPLE 2.4. Instead of the standard n -simplex Δ_n as in [Example 2.3](#), we consider Δ_n to be the subset of the boundary of I^{n+1} (the standard $(n+1)$ -cube in \mathbb{R}^{n+1}) given by


$$\left\{ (x_0, \dots, x_n) \in \mathbb{R}^{n+1} : 0 \leq x_i \leq 1, \prod_i x_i = 0 \right\}, \quad (2.7)$$

that is, Δ_n is identified with the backfaces of I^{n+1} ([Figure 2.2](#)).

Define the maps $\delta_i : \Delta_{n-1} \rightarrow \Delta_n$ and $\sigma_i : \Delta_{n+1} \rightarrow \Delta_n$ by

$$\begin{aligned} \delta_i(x_0, \dots, x_{n-1}) &= (x_0, \dots, x_{i-1}, 1, x_i, x_{i+1}, \dots, x_{n-1}), \\ \sigma_i(x_0, \dots, x_{n+1}) &= (x_0, \dots, x_{i-1}, x_i \cdot x_{i+1}, x_{i+2}, \dots, x_{n+1}) \end{aligned} \quad (2.8)$$

(see [\[5, 7\]](#)). A k -face F of Δ_n is determined for two disjoint sets $A = \{a_1, a_2, \dots, a_{k+1}\}$ and $B = \{b_1, b_2, \dots, b_{n-k}\}$ such that $0 \leq a_1 < a_2 < \dots < a_{k+1} \leq n$, $0 \leq b_1 < b_2 < \dots < b_{n-k} \leq n$, $A \cup B = \{0, 1, \dots, n\}$, $0 \leq x_i \leq 1$, $\prod_{i \in A} x_i = 0$, and $x_j \equiv 1$ for all $j \in B$. Sometimes we use the notation $F = F(A, B)$. [Figure 2.3](#) shows the 1-face $F(A, B)$ of the 2-simplex Δ_2 for $A = \{1, 2\}$ and $B = \{0\}$.

FIGURE 2.3. A 1-face of Δ_2 .

Let \mathcal{T}_n be the collection of the polynomials $f : \Delta_n \rightarrow \mathbb{R}$ with \mathbb{Z} -coefficients and let $\mathcal{T} = \{\mathcal{T}_n\}_{n \geq 0}$. Then \mathcal{T} is a simplicial algebra. The face and degeneracy maps are the maps $\partial_i : \mathcal{P}_n \rightarrow \mathcal{P}_{n-1}$ and $s_i : \mathcal{P}_n \rightarrow \mathcal{P}_{n+1}$ defined, for each $f \in \mathcal{P}_n$, by

$$\partial_i(f) = f \circ \delta_i, \quad s_i(f) = f \circ \sigma_i, \quad (2.9)$$

(∂_i and s_i are the pullbacks of δ_i and σ_i).

EXAMPLE 2.5. Let \mathcal{I}_n be the ideal generated by the polynomial $\prod_{j=0}^n x_i$. Then \mathcal{P}_n can be identified with the quotient $\mathcal{T}_n = \mathbb{Z}[x_0, \dots, x_n]/\mathcal{I}_n$. Let $\mathcal{T} = \{\mathcal{T}_n\}_{n \geq 0}$. Multiplication on $\mathbb{Z}[x_0, \dots, x_n]$ induces structure of \mathbb{Z} -algebra on \mathcal{T}_n . Then \mathcal{T} is a simplicial algebra.

Let X be a simplicial set and let $C_n(X)$ be the free group on X_n . Denote by $C_*(X)$ the chain complex $(C_n(X), \partial)$ with the *boundary operator* $\partial = \sum_{i=0}^n (-1)^i d_i$. Elements of $C_n(X)$ are called *n-chains* in X . If X is a simplicial set and G is an abelian group, then the *homology of X with coefficients in G* is defined by

$$H_*(X; G) = H(C_*(X) \otimes G). \quad (2.10)$$

Denote by $C^*(X)$ the complex $(C^n(X), \delta)$ of *cochains in X with coefficients in G* where $C^n(X; G) = \text{Hom}(C_n(X), G)$ and the *coboundary operator* δ is the dual of ∂ . The *cohomology* of X with coefficients in G is defined by

$$H^*(X; G) = H(C^*(X), G). \quad (2.11)$$

3. The complex of Cenkl-Porter. Cenkl and Porter [7] first proved the de Rham theorem for the complex of cubical differential forms for a space of finite type using integration. Later Boullay, Kiefer, Majewski, Stelzer, Scheerer, Unsöld, and Vogt [1] introduced the *complex of Cenkl and Porter* or the *de Rham complex of tame differential forms* and proved the de Rham theorem for a simplicial set following Cartan's ideas.

In this section, we present the complex of Cenkl and Porter which is the complex of compatible differential forms on the backfaces of the standard cube and state the tame de Rham theorem.

Let $\Delta_n \subset \mathbb{R}^{n+1}$ denote the standard simplex (Example 2.4). A basic form of weight q on Δ_n in the coordinates x_0, x_1, \dots, x_n is a differential form

$$x_{i_1}^{a_1} \cdots x_{i_j}^{a_j} x_{k_1}^{b_1} dx_{k_1} \wedge \cdots \wedge x_{k_p}^{b_p} dx_{k_p}, \quad (3.1)$$

where $\{i_1, \dots, i_j\}$ and $\{k_1, \dots, k_p\}$ are disjoint subsets of $\{0, 1, \dots, n\}$, the a 's and b 's are nonnegative integers, and $q = \max\{a_1, \dots, a_j, b_1+1, \dots, b_p+1\}$. Let $\mathbb{Q}_q = \mathbb{Z}[1/2, \dots, 1/q]$ be the smallest subring of the rationals such that Q_q contains $1/p$ if $0 < p \leq q$ for $q > 1$, and $\mathbb{Q}_0 = \mathbb{Q}_1 = \mathbb{Z}$. Denote by $T^{p,q}(\Delta_n)$ the module of \mathbb{Q}_q -linear combinations of basic p -forms of weight less than or equal to q . The wedge product \wedge extends to a map

$$\wedge : T_n^{p_1, q_1}(\mathbb{Z}) \otimes T_n^{p_2, q_2}(\mathbb{Z}) \longrightarrow T_n^{p_1 + p_2, q_1 + q_2}(\mathbb{Z}) \quad (3.2)$$

and the usual differential d extends to a morphism of \mathbb{Z} -modules $d : T_n^{p,q}(\mathbb{Z}) \rightarrow T_n^{p+1,q}(\mathbb{Z})$. We also have the inclusion map $T_n^{p,q}\mathbb{Z} \hookrightarrow T_n^{p,q+1}(\mathbb{Z})$. Then for every $n \geq 0$, $T^{*,*} = \{T^{p,q}(\Delta_n)\}_{n \geq 0}$ is a simplicial differential graded algebra (DGA) with filtration. For the proofs of the next two results we refer to [1, 7].

PROPOSITION 3.1. *If Δ_p is a p -simplex contained in Δ_n , and $\omega \in T^{p,q}(\Delta_n)$, then*

$$\int_{\Delta_p} \omega \in \mathbb{Q}_q. \quad (3.3)$$

Let $X = \{X_n\}$ be a simplicial set and let $T(X) = \text{Mor}(T^{*,*}, X)$ (morphisms of simplicial sets). The Stokes' theorem implies that for any $q \geq 0$, integration of tame forms induces a map of cochain complexes $I : T^{*,q}(X) \rightarrow C^*(X; \mathbb{Q}_q)$. Then we have the following theorem.

THEOREM 3.2 (the tame de Rham theorem). *Let X be a simplicial set of finite type. Then for each $q \geq 1$ and any \mathbb{Q}_q -module M there is a natural isomorphism of \mathbb{Q}_q -modules*

$$H^i(T^{*,q}(X), M) \xrightarrow{\cong} H^i(X; M) \quad \forall i \geq 0. \quad (3.4)$$

The isomorphism is induced by integration.

4. Noncommutative differential forms. In this section, we present the complex of noncommutative differential forms or the noncommutative de Rham complex, which is a generalization of the standard de Rham complex on a manifold M , but the algebra of smooth functions on M is replaced by an arbitrary associative algebra with unit. Noncommutative forms were introduced by Connes [8, 9]. Karoubi [11] used noncommutative forms to define the noncommutative de Rham complex $\Omega(X)$ and proved a noncommutative version of the de Rham theorem for a simplicial space X [12]. Here we present the basic properties of the noncommutative de Rham complex of an algebra \mathcal{A} over a commutative ring Λ .

Let \mathcal{A} be an algebra over a commutative ring Λ (with unit). Let $\mu : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$ denote the multiplication operation on \mathcal{A} (all rings are considered to be commutative and

unitary, and all algebras are with unit). The *differential forms of degree n* are the elements of the tensor product of Λ -algebras

$$\mathcal{T}^n(\mathcal{A}) = \underbrace{\mathcal{A} \otimes_{\Lambda} \mathcal{A} \otimes_{\Lambda} \cdots \otimes_{\Lambda} \mathcal{A}}_{n+1 \text{ times}}. \quad (4.1)$$

The algebra $\mathcal{T}^*(\mathcal{A}) = \otimes_{n \geq 0} \mathcal{T}^n(\mathcal{A})$ is a Λ -algebra with the multiplication $\cdot : \mathcal{T}^n(\mathcal{A}) \otimes \mathcal{T}^m(\mathcal{A}) \rightarrow \mathcal{T}^{n+m}(\mathcal{A})$ defined by the formula

$$(a_0 \otimes a_1 \otimes \cdots \otimes a_n) \cdot (b_0 \otimes b_1 \otimes \cdots \otimes b_m) = a_0 \otimes a_1 \otimes \cdots \otimes (a_n \cdot b_0) \otimes b_1 \otimes \cdots \otimes b_m. \quad (4.2)$$

The *differential* operator $D : \mathcal{T}^n(\mathcal{A}) \rightarrow \mathcal{T}^{n+1}(\mathcal{A})$ is defined by the formula

$$\begin{aligned} D(a_0 \otimes a_1 \otimes \cdots \otimes a_n) &= 1 \otimes a_0 \otimes a_1 \otimes \cdots \otimes a_n \\ &+ \sum_{j=1}^n (-1)^j a_0 \otimes a_1 \otimes \cdots \otimes a_{j-1} \otimes 1 \otimes a_j \cdots a_n \\ &+ (-1)^{n+1} a_0 \otimes a_1 \otimes \cdots \otimes a_n \otimes 1. \end{aligned} \quad (4.3)$$

THEOREM 4.1. *If $\omega \in \mathcal{T}^n(\mathcal{A})$ and $\theta \in \mathcal{T}^m(\mathcal{A})$, then*

- (1) $D^2(\omega) = 0$;
- (2) $D(\omega \cdot \theta) = D(\omega) \cdot \theta + (-1)^n \omega \cdot D(\theta)$ (Leibniz identity).

Then $\mathcal{T}^*(\mathcal{A})$ is a DGA and the cohomology of the complex $(\mathcal{T}^*(\mathcal{A}), D)$ is trivial. Suppose that \mathcal{A} is an augmented Λ -algebra with an augmentation $\lambda : \mathcal{A} \rightarrow \Lambda$ (morphism of rings) such that $\lambda(1) = 1$. Consider the map of modules $\iota_\lambda : \mathcal{T}^n(\mathcal{A}) \rightarrow \mathcal{T}^{n-1}(\mathcal{A})$ defined by

$$\iota_\lambda(a_0 \otimes \cdots \otimes a_n) = \lambda(a_0)(a_1 \otimes \cdots \otimes a_n). \quad (4.4)$$

THEOREM 4.2. *Let \mathcal{A} be an augmented Λ -algebra with an augmentation $\lambda : \mathcal{A} \rightarrow \Lambda$ such that $\lambda(1) = 1$. The map $\iota_\lambda : \mathcal{T}^n(\mathcal{A}) \rightarrow \mathcal{T}^{n-1}(\mathcal{A})$ is a contracting homotopy,*

$$D\iota_\lambda + \iota_\lambda D = 1. \quad (4.5)$$

Define $\Omega^0(\mathcal{A}) = \mathcal{A}$ and $\Omega^1(\mathcal{A}) = \ker \mu$, the Λ -module $\Omega^1(\mathcal{A})$ is an \mathcal{A} -bimodule. The *noncommutative differential forms of degree n* are the elements of the tensor product of \mathcal{A} -modules

$$\Omega^n(\mathcal{A}) = \underbrace{\Omega^1(\mathcal{A}) \otimes_{\mathcal{A}} \Omega^1(\mathcal{A}) \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} \Omega^1(\mathcal{A})}_{n \text{ times}}. \quad (4.6)$$

The product of differential forms is defined by juxtaposition of tensor products. Then the direct sum

$$\Omega^*(\mathcal{A}) = \bigoplus_{n \geq 0} \Omega^n(\mathcal{A}) \quad (4.7)$$

is a graded algebra. The *differential* $d : \Omega^0(\mathcal{A}) \rightarrow \Omega^1(\mathcal{A})$ is defined by the formula

$$d(a) = 1 \otimes a - a \otimes 1. \quad (4.8)$$

Thus we have the isomorphism of Λ -modules $\mathcal{A} \otimes \mathcal{A}/\Lambda \rightarrow \Omega^1(\mathcal{A})$ such that $a \otimes \bar{b} \mapsto adb$. Then $\Omega^n(\mathcal{A})$ can be identified with the tensor product of Λ -modules

$$\mathcal{A} \otimes \underbrace{\mathcal{A}/\Lambda \otimes \mathcal{A}/\Lambda \otimes \cdots \otimes \mathcal{A}/\Lambda}_{n \text{ times}}. \quad (4.9)$$

A noncommutative differential form of degree n can be written as a linear combination of terms of the form $a_0 da_1 da_2 \cdots da_n$ and the morphism d extends to forms of degree n of $\Omega^n(\mathcal{A})$ by the formula

$$d(a_0 da_1 \cdots da_n) = da_0 da_1 \cdots da_n = 1 da_0 da_1 \cdots da_n. \quad (4.10)$$

THEOREM 4.3. *If $\omega \in \Omega^n(\mathcal{A})$ and $\theta \in \Omega^m(\mathcal{A})$, then*

- (1) $d^2(\omega) = 0$;
- (2) $d(\omega \cdot \theta) = d(\omega) \cdot \theta + (-1)^n \omega \cdot d(\theta)$ (Leibniz identity).

REMARK 4.4. The DGA $\Omega^*(\mathcal{A})$ is called the *differential enveloping* of \mathcal{A} , and it is the solution of a universal problem: for a DGA \mathcal{B}^* and an algebra morphism $f: \mathcal{A} \rightarrow \mathcal{B}^0$, there exists a unique morphism of DGA's $f^*: \Omega^*(\mathcal{A}) \rightarrow \mathcal{B}^*$ which agrees with f at degree 0. The complex $(\Omega^*(\mathcal{A}), d)$ is known as the *universal differential calculus* of \mathcal{A} or as the *noncommutative de Rham complex* of \mathcal{A} .

There is an inclusion of DGA's sending $\Omega^*(\mathcal{A})$ to $\mathcal{T}^*(\mathcal{A})$. On the other hand, for any $n \geq 0$ there is a *projection operator* $J: \mathcal{T}^n(\mathcal{A}) \rightarrow \Omega^n(\mathcal{A})$ defined by $J(a_0 \otimes a_1 \otimes \cdots \otimes a_n) = a_0 da_1 \cdots da_n$.

A noncommutative differential form ω is said to be *closed* if $d\omega = 0$. We say that $\omega \in \Omega^n(\mathcal{A})$ is *exact* if there exists $\eta \in \Omega^{n-1}(\mathcal{A})$ such that $\omega = d\eta$. The fact that the complex $(\Omega^*(\mathcal{A}), d)$ has trivial cohomology is known as the noncommutative Poincaré lemma.

LEMMA 4.5. *Let \mathcal{A} be an augmented Λ -algebra, then every closed form $\omega \in \Omega^*(\mathcal{A})$ is exact.*

PROOF. As in [Theorem 4.2](#), let $\lambda: \mathcal{A} \rightarrow \Lambda$ be a Λ -linear form with $\lambda(1) = 1$. We prove that there exists a homotopy contraction $J_\lambda: \Omega^n(\mathcal{A}) \rightarrow \Omega^{n-1}(\mathcal{A})$. To define J_λ we express elements of $\Omega^n(\mathcal{A})$ as elements of $\mathcal{T}^{n+1}(\mathcal{A})$ using inclusion, next we apply ι_λ , and then we apply the projection J . Thus for $\omega = a_0 da_1 \cdots da_n$, we get $J_\lambda(\omega) = \lambda(a_0) a_1 da_2 \cdots da_n - \lambda(a_0 a_1) da_2 \cdots da_n$. First we show that ι_λ is well defined. Obviously it is enough to prove that $\iota_\lambda: \Omega^1(\mathcal{A}) \rightarrow \Omega^0(\mathcal{A})$ is well defined. Let $\omega = a_0 da_1 \in \Omega^1(\mathcal{A})$, $a \in \Lambda$, and $a'_1 = a_1 + a \cdot 1 \in \mathcal{A}$. Then $J_\lambda(a_0 da'_1) = \lambda(a_0) a'_1 - \lambda(a_0 a'_1) \cdot 1 = \lambda(a_0) a_1 - \lambda(a_0 a_1) \cdot 1 = J_\lambda(a_0 da_1)$. Now for $\omega = a_0 da_1 \cdots da_n \in \Omega^n(\mathcal{A})$, we have

$$\begin{aligned} dJ_\lambda(\omega) + J_\lambda d(\omega) &= \lambda(a_0) da_1 da_2 \cdots da_n - \lambda(a_0 a_1) d1 da_2 \cdots da_n \\ &\quad + \lambda(1) a_0 da_1 da_2 \cdots da_n - \lambda(1 \cdot a_0) da_1 \cdots da_n \\ &= \omega. \end{aligned} \quad (4.11)$$

But $d\omega = 0$. Therefore $dJ_\lambda(\omega) = \omega$. □

If $\mathcal{A} = \{\mathcal{A}_n\}_{n \geq 0}$ is a simplicial algebra, then $\Omega^*(\mathcal{A}) = \{\Omega^*(\mathcal{A}_n)\}_{n \geq 0}$ is a simplicial DGA. Next we define the face and degeneracy operators for $\Omega^*(\mathcal{A})$.

Let $\mathcal{A} = \{\mathcal{A}_n^*\}_{n \geq 0}$ be a simplicial graded algebra. For each n consider the simplicial tensor algebra

$$\mathcal{T}^*(\mathcal{A}_n) = \bigoplus_{p \geq 0} \mathcal{A}_n^{\otimes p}, \quad (4.12)$$

where the face and degeneracy operators $\partial_i : \mathcal{A}_n^{\otimes p} \rightarrow \mathcal{A}_{n-1}^{\otimes p}$ and $s_i : \mathcal{A}_n^{\otimes p} \rightarrow \mathcal{A}_{n+1}^{\otimes p}$ are defined by

$$\begin{aligned} \partial_i(a_0 \otimes a_1 \otimes \cdots \otimes a_p) &= \partial_i a_0 \otimes \partial_i a_1 \otimes \cdots \otimes \partial_i a_p, \\ s_i(a_0 \otimes a_1 \otimes \cdots \otimes a_p) &= s_i a_0 \otimes s_i a_1 \otimes \cdots \otimes s_i a_p. \end{aligned} \quad (4.13)$$

PROPOSITION 4.6. *Let $\mathcal{A} = \{\mathcal{A}_n^*\}_{n \geq 0}$ be a simplicial algebra. If D is the differential on $\mathcal{T}^*(\mathcal{A}_n)$, then $\mathcal{T}^*(\mathcal{A})$ is a simplicial DGA.*

Observe that the restriction of ∂_i to $\Omega^1(\mathcal{A}_n)$ applied to $adb \in \Omega^1(\mathcal{A}_n)$ is

$$\begin{aligned} \partial_i(adb) &= \partial_i(a \otimes b - ab \otimes 1) = \partial_i(a) \otimes \partial_i(b) - \partial_i(ab) \otimes 1 \\ &= \partial_i(a) \otimes \partial_i(b) - \partial_i(a)\partial_i(b) \otimes 1 \in \mathcal{A}_{n-1} \otimes \mathcal{A}_{n-1}. \end{aligned} \quad (4.14)$$

If μ_{n-1} denotes multiplication on \mathcal{A}_{n-1} , then

$$\begin{aligned} \mu_{n-1}(\partial_i(adb)) &= \mu_{n-1}(\partial_i(a) \otimes \partial_i(b) - \partial_i(a)\partial_i(b) \otimes 1) \\ &= \partial_i(a) \cdot \partial_i(b) - \partial_i(a)\partial_i(b) \cdot 1 = 0. \end{aligned} \quad (4.15)$$

Therefore $adb \in \ker \mu_{n-1} = \Omega^1(\mathcal{A}_{n-1})$. In particular, if we take elements of the form $da \in \Omega^1(\mathcal{A}_{n-1})$, then we get

$$\partial_i(da) = \partial_i(1 \otimes a - a \otimes 1) = 1 \otimes \partial_i a - \partial_i a \otimes 1 = d(\partial_i a). \quad (4.16)$$

Then extend ∂_i to $\Omega^P(\mathcal{A})$ by setting

$$\partial_i(a_0 da_1 \cdots da_p) = \partial_i(a_0) d(\partial_i a_1) \cdots d(\partial_i a_p) \in \Omega^P(\mathcal{A}_{n-1}). \quad (4.17)$$

Similarly s_i can be extended to $\Omega^P(\mathcal{A})$. Then we have the following proposition.

PROPOSITION 4.7. *If $\mathcal{A} = \{\mathcal{A}_n\}_{n \geq 0}$ is a simplicial graded algebra, then $\Omega^*(\mathcal{A}) = \{\Omega^*(\mathcal{A}_n)\}_{n \geq 0}$ is a simplicial DGA.*

A noncommutative version of the de Rham theorem was proved by Karoubi in [12]. Karoubi considered \mathcal{A}_n to be the quotient Λ -algebra $\Lambda[x_0, x_1, \dots, x_n]/(x_0 + x_1 + \cdots + x_n - 1)$. Let $\Omega^*(\mathcal{A}_n)$ be the algebra of noncommutative forms on \mathcal{A}_n and let $\mathcal{A} = \{\mathcal{A}_n\}_{n \geq 0}$. The algebra $\Omega^*(\mathcal{A}_n)$ is the noncommutative algebra generated by the symbols x_i and $dx_i, 0 \leq i \leq n$ and the following relations:

$$\sum_{i=0}^n x_i = 1, \quad \sum_{i=0}^n dx_i = 0, \quad x_i x_j = x_j x_i. \quad (4.18)$$

Then we have the following theorem.

THEOREM 4.8 (the noncommutative de Rham theorem). *Let X be a simplicial set and let $\Omega^*(X) = \text{Mor}(X, \Omega^*(\mathcal{A}))$. Then there exists a natural isomorphism of Λ -modules*

$$H^i(\Omega^*(X)) \cong H^i(X; \Lambda) \quad \forall i \geq 0. \quad (4.19)$$

A slightly more general version of [Theorem 4.8](#) was proved by Cenkl in [\[3, 4\]](#).

5. The noncommutative complex of Cenkl-Porter. In [\[12\]](#), Karoubi conjectured that the noncommutative de Rham theorem could be proved using integration of noncommutative differential forms assuming that Λ is a ring containing the ring of the rational numbers \mathbb{Q} . We present a solution of a more general problem by considering a noncommutative version of the tame de Rham complex of Cenkl-Porter. This complex is constructed by defining a filtration on $\Omega^*(\mathcal{T})$, the algebra of noncommutative differential forms on $\mathcal{T} = \bigoplus_{n \geq 0} \mathcal{T}_n$, where \mathcal{T}_n are the polynomials restricted to n -simplex Δ_n (see [Example 2.4](#)). Then we prove some basic properties of that complex. In particular we prove the noncommutative tame Poincaré lemma.

REMARK 5.1. [Propositions 4.6](#) and [4.7](#) imply that $\Omega^*(\mathcal{T})$ is a simplicial DGA.

We establish some conventions of notation. Let \mathbb{Z}_+ be the set of nonnegative integers. Let \mathbb{Z}_+^{n+1} be the set of multi-indexes $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_n)$ with $\alpha_i \in \mathbb{Z}_+$, and let $|\alpha| = \sum_i \alpha_i$. For $x = (x_0, x_1, \dots, x_n) \in \Delta_n$ and $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_n)$, $\varepsilon = (\varepsilon_0, \varepsilon_1, \dots, \varepsilon_n) \in \mathbb{Z}_+^{n+1}$, let

$$\begin{aligned} x^\alpha &= x_0^{\alpha_0} x_1^{\alpha_1} \cdots x_n^{\alpha_n}, \\ dx^\varepsilon &= dx_0^{\varepsilon_0} dx_1^{\varepsilon_1} \cdots dx_n^{\varepsilon_n} = (dx_0)^{\varepsilon_0} (dx_1)^{\varepsilon_1} \cdots (dx_n)^{\varepsilon_n}. \end{aligned} \quad (5.1)$$

If $A = \{a_1, a_2, \dots, a_p\} \subset \{0, 1, \dots, n\}$, we write

$$\begin{aligned} x_A^\alpha &= x_{a_1}^{\alpha_{a_1}} x_{a_2}^{\alpha_{a_2}} \cdots x_{a_p}^{\alpha_{a_p}}, \\ (1 - x_A)^\alpha &= (1 - x_{a_1})^{\alpha_{a_1}} (1 - x_{a_2})^{\alpha_{a_2}} \cdots (1 - x_{a_p})^{\alpha_{a_p}}, \\ dx_A^\varepsilon &= dx_{a_1}^{\varepsilon_{a_1}} dx_{a_2}^{\varepsilon_{a_2}} \cdots dx_{a_p}^{\varepsilon_{a_p}}. \end{aligned} \quad (5.2)$$

Let $\Omega_n(\mathbb{Z})$ be the algebra of all \mathbb{Z} -linear combinations of basic *tame noncommutative differential forms*

$$\omega = x^{\alpha_1} dx^{\varepsilon_1} x^{\alpha_2} dx^{\varepsilon_2} \cdots x^{\alpha_r} dx^{\varepsilon_r}, \quad \alpha_i, \varepsilon_i \in \mathbb{Z}_+^{n+1}, \quad i = 1, 2, \dots, r, \quad (5.3)$$

with $0 \leq x_j \leq 1$ and $\prod_{j=0}^n x_j = 0$. These are the compatible noncommutative differential forms on the backfaces of the cube I^{n+1} (see [Example 2.4](#)). If $\sum_i |\varepsilon_i| = p$ we say that ω is a p -form (note that 0-forms are polynomials).

Let $\|\omega\|_j = \sum_i (\alpha_{ij} + \varepsilon_{ij})$. The *weight* of ω is defined by $\|\omega\| = \max\{\|\omega\|_j : j = 0, 1, \dots, n\}$.

Let $\Omega_n^{p,q}(\mathbb{Z})$ be the set of all the p -forms of $\Omega_n(\mathbb{Z})$ of weight $\|\omega\| \leq q$ and let $\Omega_n^{*,*}(\mathbb{Z}) = \{\Omega_n^{p,q}(\mathbb{Z})\}_{p,q \geq 0}$.

REMARK 5.2. For all n , p , and q , $\Omega_n^{p,q}(\mathbb{Z})$ is a finitely generated free \mathbb{Z} -module.

PROPOSITION 5.3. *For every $n \geq 0$, $\Omega_n^{*,*}(\mathbb{Z}) = \{\Omega_n^{p,q}(\mathbb{Z})\}_{p,q \geq 0}$ is a differential graded algebra with filtration.*

PROOF. We have to prove that \cdot and d on $\Omega^*(\mathcal{T})$ induce maps \cdot and d such that

$$(1) \quad \Omega_n^{p_1,q_1}(\mathbb{Z}) \otimes \Omega_n^{p_2,q_2}(\mathbb{Z}) \xrightarrow{\cdot} \Omega_n^{p_1+p_2,q_1+q_2}(\mathbb{Z}),$$

$$(2) \quad \Omega_n^{p,q}(\mathbb{Z}) \xrightarrow{d} \Omega_n^{p+1,q}(\mathbb{Z}),$$

$$(3) \quad \Omega_n^{p,q}(\mathbb{Z}) \hookrightarrow \Omega_n^{p,q+1}(\mathbb{Z}).$$

PROOF OF (1). Let

$$\begin{aligned} \omega &= \omega_1 \cdot \omega_2 \cdot \dots \cdot \omega_r \in \Omega_n^{p_1,q_1}(\mathbb{Z}), \quad \omega_i = x^{\alpha_i} dx^{\varepsilon_i}, \quad i = 1, 2, \dots, r, \\ \eta &= \eta_1 \cdot \eta_2 \cdot \dots \cdot \eta_s \in \Omega_n^{p_2,q_2}(\mathbb{Z}), \quad \eta_j = x^{\beta_j} dx^{\varepsilon_j}, \quad j = 1, 2, \dots, s. \end{aligned} \quad (5.4)$$

Then $\omega \cdot \eta = \omega_1 \cdot \omega_2 \cdot \dots \cdot \omega_r \cdot \eta_1 \cdot \eta_2 \cdot \dots \cdot \eta_s = \omega_1 \cdot \omega_2 \cdot \dots \cdot \omega_r \cdot \omega_{r+1} \cdot \dots \cdot \omega_{r+s}$ where $\omega_{r+j} = \eta_j (\|\omega_{r+j}\|_k = \|\eta_j\|_k \text{ for all } j \text{ and } k)$;

$$\begin{aligned} \|\omega \cdot \eta\|_k &= \sum_{l=1}^{r+s} \|\omega_l\|_k = \sum_{l=1}^r \|\omega_l\|_k + \sum_{l=r+1}^{r+s} \|\omega_l\|_k \\ &= \sum_{l=1}^r \|\omega_l\|_k + \sum_{l=1}^s \|\eta_l\|_k \leq \|\omega_l\|_k + \|\eta_l\|_k, \end{aligned} \quad (5.5)$$

therefore

$$\|\omega \cdot \eta\| \leq \|\omega_l\| + \|\eta_l\|. \quad (5.6)$$

On the other hand, we have $\sum_i |\varepsilon_i| + \sum_j |\varepsilon_j| = p_1 + p_2$. Therefore,

$$\|\omega \cdot \eta\| \in \Omega_n^{p_1+p_2,q_1+q_2}(\mathbb{Z}). \quad (5.7)$$

PROOF OF (2). Let $\omega = \omega_1 \cdot \omega_2 \cdot \dots \cdot \omega_r \in \Omega_n^{p,q}(\mathbb{Z})$ with $\omega_i = x^{\alpha_i} dx^{\varepsilon_i} \in \Omega_n^{p_i,q_i}(\mathbb{Z})$, $i = 1, 2, \dots, r$. Then

$$d\omega = d(x^{\alpha_i}) \cdot dx^{\varepsilon_i} + x^{\alpha_i} \cdot d(dx^{\varepsilon_i}) = d(x^{\alpha_i}) \cdot dx^{\varepsilon_i}. \quad (5.8)$$

Note that

$$\begin{aligned} d(x^{\alpha_i}) &= d(x_0^{\alpha_{i0}} \cdots x_n^{\alpha_{in}}) \\ &= \sum_{j=0}^n x_0^{\alpha_{i0}} \cdots d(x_j^{\alpha_{ij}}) \cdots x_n^{\alpha_{in}} \\ &= \sum_{j=0}^n x_0^{\alpha_{i0}} \cdots x_{j-1}^{\alpha_{i,j-1}} \left(\sum_{k=1}^n x_j^{k-1} dx_j x_j^{\alpha_{i,j-k}} \right) \cdots x_n^{\alpha_{in}} \\ &= \sum_{j=0}^n \sum_{k=1}^n x_0^{\alpha_{i0}} \cdots x_{j-1}^{\alpha_{i,j-1}} x_j^{k-1} dx_j x_j^{\alpha_{i,j-k}} \cdots x_n^{\alpha_{in}}. \end{aligned} \quad (5.9)$$

Then $\|d(x^{\alpha_i})\|_j = \|x^{\alpha_i}\|_j$ for $j = 0, 1, \dots, n$, therefore $\|d(x^{\alpha_i})\| = \|x^{\alpha_i}\|$. Hence $\|d(x^{\alpha_i}) dx^{\varepsilon_i}\| = \|x^{\alpha_i} dx^{\varepsilon_i}\|_j$ and $\|d\omega\| = \|\omega\|$. On the other hand, x^{α_i} is a 0-form, therefore $d(x^{\alpha_i})$ is a 1-form and $d\omega$ is a $(p+1)$ -form, then $d\omega \in \Omega_n^{p+1,q}(\mathbb{Z})$.

PROOF OF (3). The proof is obvious. \square

Now we show that $\Omega^{*,*}(\mathbb{Z}) = \{\Omega_n^{*,*}(\mathbb{Z})\}_{n \geq 0}$ is a simplicial algebra (n denotes the simplicial index). If $p \geq 0$ and $q \geq 1$ are fixed we consider a form $\omega \in \Omega_n^{p,q}(\mathbb{Z})$ as an element of $\Omega^p(\mathcal{T}_n)$ for all n . But $\Omega^*(\mathcal{T})$ is a simplicial algebra so we may restrict the face and degeneracy operators $\partial_i : \Omega^p(\mathcal{T}_n) \rightarrow \Omega^p(\mathcal{T}_{n+1})$ and $s_i : \Omega^p(\mathcal{T}_n) \rightarrow \Omega^p(\mathcal{T}_{n+1})$ to $\Omega_n^{p,q}(\mathbb{Z})$. Then we must verify that $\text{im } \partial_i \subset \Omega_{n-1}^{p,q}(\mathbb{Z})$ and $\text{im } s_i \subset \Omega_{n+1}^{p,q}(\mathbb{Z})$ for all $p \geq 0$, $q \geq 1$. Suppose that

$$\omega = x^{\alpha_1} dx^{\varepsilon_1} x^{\alpha_2} dx^{\varepsilon_2} \cdots x^{\alpha_r} dx^{\varepsilon_r}. \quad (5.10)$$

Then

$$\partial_i(\omega) = \partial_i(x^{\alpha_1}) \partial_i(dx^{\varepsilon_1}) \partial_i(x^{\alpha_2}) \partial_i(dx^{\varepsilon_2}) \cdots \partial_i(x^{\alpha_r}) \partial_i(dx^{\varepsilon_r}), \quad (5.11)$$

where $\partial_i(x^{\alpha_j}) = (\partial_i x_0)^{\alpha_{j0}} \cdots (\partial_i x_n)^{\alpha_{jn}}$ and $\partial_i(dx^{\varepsilon_j}) = (d\partial_i x_0)^{\varepsilon_{j0}} \cdots (d\partial_i x_n)^{\varepsilon_{jn}}$.

If $\varepsilon_{ji} \neq 0$, for some j , then $(d\partial_i(x_i))^{\varepsilon_{ji}} = (d1)^{\varepsilon_{ji}} = 0$. Then $\partial_i \omega = 0$ and $\|\partial_i \omega\| \leq \|\omega\|$. If $\varepsilon_{ji} = 0$ for all j , it is enough to consider one block,

$$\omega = x^\alpha dx^\varepsilon = x_0^{\alpha_0} \cdots x_n^{\alpha_n} dx_0^{\varepsilon_0} \cdots dx_n^{\varepsilon_n} \quad \text{with } \alpha_i, \varepsilon_i \in \mathbb{Z}_+, \quad i = 0, 1, \dots, n. \quad (5.12)$$

Then

$$\begin{aligned} \partial_i(\omega) &= (\partial_i x_0)^{\alpha_0} \cdots (\partial_i x_n)^{\alpha_n} (\partial_i dx_0)^{\varepsilon_0} \cdots (\partial_i dx_n)^{\varepsilon_n} \\ &= x_0^{\alpha_0} \cdots x_{i-1}^{\alpha_{i-1}} 1^{\alpha_i} x_i^{\alpha_{i+1}} \cdots x_{n-1}^{\alpha_n} dx_0^{\varepsilon_0} \cdots dx_{i-1}^{\varepsilon_{i-1}} 1^0 dx_i^{\varepsilon_{i+1}} \cdots dx_{n-1}^{\varepsilon_n}. \end{aligned} \quad (5.13)$$

Then we have

$$\|\partial_i(\omega)\|_k = \begin{cases} \|\omega\|_k, & \text{for } 0 \leq k < i, \\ \|\omega\|_{k+1}, & \text{for } 0 \leq i \leq k < n, \\ 0, & \text{for } 0 \leq i < k = n. \end{cases} \quad (5.14)$$

Therefore, $\|\partial_i(\omega)\| \leq \|\omega\|$. Then

$$\partial_i(\omega) \in \Omega_{n-1}^{p,q}(\mathbb{Z}). \quad (5.15)$$

Note that if $\|\omega\|_k < \|\omega\|_i$, for all $k \neq i$, then we have a sharp inequality $\|\partial_i(\omega)\| < \|\omega\|$.

Similarly, for $\omega = x^{\alpha_1} dx^{\varepsilon_1} x^{\alpha_2} dx^{\varepsilon_2} \cdots x^{\alpha_r} dx^{\varepsilon_r}$, we have

$$s_i(\omega) = s_i(x^{\alpha_1}) s_i(dx^{\varepsilon_1}) s_i(x^{\alpha_2}) s_i(dx^{\varepsilon_2}) \cdots s_i(x^{\alpha_r}) s_i(dx^{\varepsilon_r}), \quad (5.16)$$

where

$$s_i(x^{\alpha_j}) = (s_i x_0)^{\alpha_{j0}} \cdots (s_i x_n)^{\alpha_{jn}}, \quad s_i(dx^{\varepsilon_j}) = (ds_i x_0)^{\varepsilon_{j0}} \cdots (ds_i x_n)^{\varepsilon_{jn}}. \quad (5.17)$$

Then

$$\begin{aligned} s_i(\omega) &= (s_i x_0)^{\alpha_0} \cdots (s_i x_n)^{\alpha_n} (s_i dx_0)^{\varepsilon_0} \cdots (s_i dx_n)^{\varepsilon_n} \\ &= x_0^{\alpha_0} \cdots x_{i-1}^{\alpha_{i-1}} (x_i \cdot x_{i+1})^{\alpha_i} x_{i+2}^{\alpha_{i+1}} \cdots x_{n+1}^{\alpha_{n+1}} \\ &\quad \cdot dx_0^{\varepsilon_0} \cdots d(x_i \cdot x_{i+1})^{\varepsilon_i} dx_{i+2}^{\varepsilon_{i+1}} \cdots dx_{n+1}^{\varepsilon_{n+1}} \\ &= x_0^{\alpha_0} \cdots x_{i-1}^{\alpha_{i-1}} x_i^{\alpha_i} x_{i+1}^{\alpha_{i+1}} x_{i+2}^{\alpha_{i+2}} \cdots x_{n+1}^{\alpha_{n+1}} \\ &\quad \cdot dx_0^{\varepsilon_0} \cdots (dx_i x_{i+1} + x_i dx_{i+1})^{\varepsilon_i} dx_{i+2}^{\varepsilon_{i+1}} \cdots dx_{n+1}^{\varepsilon_{n+1}}. \end{aligned} \quad (5.18)$$

Thus $s_i(\omega)$ is given by the expression

$$\begin{aligned} \sum_{j=0}^{\varepsilon_i} \binom{\varepsilon_i}{j} x_0^{\alpha_0} \cdots x_{i-1}^{\alpha_{i-1}} x_i^{\alpha_i} x_{i+1}^{\alpha_i} x_{i+2}^{\alpha_{i+1}} \cdots x_{n+1}^{\alpha_{n+1}} \\ \cdot dx_0^{\varepsilon_0} \cdots dx_i^{\varepsilon_{i-j}} x_{i+1}^{\varepsilon_{i-j}} x_i^j dx_{i+1}^j dx_{i+2}^{\varepsilon_{i+1}} \cdots dx_{n+1}^{\varepsilon_n}. \end{aligned} \quad (5.19)$$

Note that $\|s_i(\omega)\|_k = \alpha_i + \varepsilon_i - j + j = \|\omega\|_i$ for $k = i$, $i+1$, and $\|s_i(\omega)\|_k = \alpha_{k-1} + \varepsilon_{k-1} = \|\omega\|_{k-1}$ for $k > i+1$. Then

$$\|s_i(\omega)\|_k = \begin{cases} \|\omega\|_k, & \text{for } k < i, \\ \|\omega\|_i, & \text{for } k = i, i+1, \\ \|\omega\|_{k-1}, & \text{for } k > i+1. \end{cases} \quad (5.20)$$

Therefore, $\|s_i(\omega)\| \leq \|\omega\|$ and $s_i(\omega) \in \Omega_{n+1}^{p,q}(\mathbb{Z})$.

From the definition and [Proposition 4.7](#) it follows that ∂_i and s_i are morphisms of DGA's. Then we have the following proposition.

PROPOSITION 5.4. *The algebra $\Omega^{*,*}(\mathbb{Z}) = \{\Omega_n^{*,*}(\mathbb{Z})\}_{n \geq 0}$ is a simplicial DGA.*

Now consider $0 \leq x_j \leq 1$ for $j = 0, 1, \dots, n$ with $\prod_j x_j = 0$. Define

$$\Omega^{p,q}(\Delta_n) = \Omega_n^{p,q}(\mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q}_q, \quad (5.21)$$

where $\mathbb{Q}_q = \mathbb{Z}[1/2, \dots, 1/q]$, for $q > 1$, and $\mathbb{Q}_0 = \mathbb{Q}_1 = \mathbb{Z}$.

Let Δ_n^k , $0 \leq k \leq n$, be the k -skeleton of Δ_n , and let

$$\Omega^{p,q}(\Delta_n, \Delta_n^k) = \left\{ \omega \in \Omega^{p,q}(\Delta_n) : \omega|_{\Delta_n^k} \equiv 0 \right\}. \quad (5.22)$$

Let $\Omega^{p,q}(\Delta_n^k)$ be the set of all \mathbb{Q}_q -linear combinations of forms which are nonzero on exactly one k -face of Δ_n^k .

PROPOSITION 5.5. *The sequence*

$$\Omega^{p,q}(\Delta_n, \Delta_n^{k-1}) \xrightarrow{\text{r restrictions}} \Omega^{p,q}(\Delta_n^k, \Delta_n^{k-1}) \longrightarrow 0 \quad (5.23)$$

is an exact sequence of \mathbb{Q}_q -modules for all $p \geq 0$, $q \geq 1$.

PROOF. Let $\omega \in \Omega^{p,q}(\Delta_n, \Delta_n^{k-1})$ with $q \geq 1$. Then the form ω is a linear combination $\omega = \sum_i \omega_i$, where each ω_i is nonzero on exactly one face of Δ_n^k . Let F be such a face. Then

$$\omega|_G \equiv 0 \quad \text{if } G \text{ is any } k\text{-face of } \Delta_n^k \text{ different from } F. \quad (5.24)$$

We write $F = F(A, B)$ (using the notation of [Example 2.4](#)) where A and B are two disjoint sets $A = \{a_1, a_2, \dots, a_{k+1}\}$ and $B = \{b_1, b_2, \dots, b_{n-k}\}$ such that $0 \leq a_1 < a_2 < \dots <$

$a_{k+1} \leq n$, $0 \leq b_1 < b_2 < \dots < b_{n-k} \leq n$, $A \cup B = \{0, 1, \dots, n\}$, $0 \leq x_i \leq 1$, for all $i \in A$, $\prod_{i \in A} x_i = 0$, and $x_j \equiv 1$ for all $j \in B$.

We have $\omega|_{\Delta_n^{k-1}} \equiv 0$. Therefore ω is a linear combination of forms of the type

$$f^1(x)dx^{\varepsilon_1} \cdots f^s(x)dx^{\varepsilon_s}, \quad (5.25)$$

where $f^j(x) = x_A^{\alpha_1}(1-x_A)^{\alpha_2}x_B^{\beta_1}(1-x_B)^{\beta_2}f_j(x)$ ($f_j(x)$ is a polynomial in x). For F we may assume that $|\beta_2| = 0$. Thus $f^j(x) = x_A^{\alpha_1}(1-x_A)^{\alpha_2}x_B^{\beta_1}f_j(x)$. Note that if $|\alpha_i| = 0$ for $i = 1, 2$ (and for all j), then $\varepsilon_{it} \neq 0$ for some t .

On F we have $f^j(x) = x_A^{\alpha_1}(1-x_A)^{\alpha_2}f_j(x_A)$.

Let $G = G(A', B')$ be another k -face of Δ_n^k . If $A = A'$, then there exists $i \in B$ with $x_i \equiv 0$, then $f^j(x) = 0$ and $\omega|_G \equiv 0$.

If $A \neq A'$ then there exists $i \in A$ such that $x_i \equiv 1$. Then either $f^j(x) \equiv 0$ or $dx_i^{\varepsilon_{it}} = 0$ for some t , in both cases we have $\omega|_G \equiv 0$.

Let $\phi(x) = x_B$, then $\phi \in \Omega^{0,1}(\Delta_n)$. Define $\omega_k = \phi \cdot \omega$. Note that there exists $i \in A$ such that $\|\omega_k\|_j = 1 \leq \|\omega_k\|_i = q$ for all $j \in B$. Therefore $\omega_k \in \Omega^{p,q}(\Delta_n)$. Moreover,

$$\omega_k|_{F(A,B)} = \phi|_{F(A,B)} \cdot \omega|_{F(A,B)} \equiv 1 \cdot \omega = \omega. \quad (5.26)$$

If $F(A', B')$ is another k -face of Δ_n^k , then we have

$$\omega_k|_{F(A',B')} = \phi|_{F(A',B')} \cdot \omega|_{F(A',B')} \equiv \phi|_{F(A',B')} \cdot 0 = 0. \quad (5.27)$$

If $F(E, H)$ is a $(k-1)$ -face of Δ_n^k , then there exists at least one $i \in A$ such that $i \notin E$. Then x_i is either 0 or 1 on $F(E, H)$. Therefore $\omega|_{F(E, H)} = 0$ and then $\omega_k|_{F(E, H)} \equiv 0$. We deduce that $\omega_k \in \Omega^{p,q}(\Delta_n, \Delta_n^{k-1})$ and $\omega_k|_{\Delta_n^k} \equiv \omega$. \square

PROPOSITION 5.6. *The sequence*

$$\Omega^{p,q}(\Delta_n) \xrightarrow{r \text{ restrictions}} \Omega^{p,q}(\partial\Delta_n) \longrightarrow 0 \quad (5.28)$$

is an exact sequence of \mathbb{Q}_q -modules for all $p \geq 0, q \geq 1$.

PROOF. For $k = 0$, the sequence

$$\Omega^{p,q}(\Delta_n) \longrightarrow \Omega^{p,q}(\Delta_n^0) \longrightarrow 0 \quad (5.29)$$

is exact, thus any element $a \in \mathbb{Q}_q$ can be pulled back to the form $\omega(x_0, \dots, x_n) = a$, that is, it is a constant \mathbb{Q}_q -polynomial.

Assume by induction that

$$\Omega^{p,q}(\Delta_n) \longrightarrow \Omega^{p,q}(\partial\Delta_n^{k-1}) \longrightarrow 0 \quad (5.30)$$

is exact. Consider the following commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \Omega^{p,q}(\Delta_n, \Delta_n^{k-1}) & \xrightarrow{r_1} & \Omega^{p,q}(\Delta_n^k, \Delta_n^{k-1}) & \longrightarrow & 0 \\
 & & i_1 \downarrow & & i_2 \downarrow & & \\
 & & \Omega^{p,q}(\Delta_n) & \xrightarrow{r_2} & \Omega^{p,q}(\Delta_n^k) & \longrightarrow & 0 \\
 & & p_1 \downarrow & & p_2 \downarrow & & \\
 & & \Omega^{p,q}(\Delta_n^{k-1}) & \xlongequal{\quad} & \Omega^{p,q}(\Delta_n^{k-1}) & \longrightarrow & 0 \\
 & & \downarrow & & & & \\
 & & 0. & & & &
 \end{array} \tag{5.31}$$

The left column is exact by induction hypothesis (and by the definition of $\Omega^{p,q}(\Delta_n^k, \Delta_n^{k-1})$). The right column is exact by definition. The first row is exact by [Proposition 5.5](#). We show that the second row is exact. Let $\omega \in \Omega^{p,q}(\Delta_n^k)$.

CASE 1. If $\omega \in \ker p_2$, then there exists $\omega' \in \Omega^{p,q}(\Delta_n^k, \Delta_n^{k-1})$ with $i_2(\omega') = \omega$. But the first row is exact, then there exists $\omega'' \in \Omega^{p,q}(\Delta_n, \Delta_n^{k-1})$ such that $\omega'' = r_1(\omega')$. Hence $\omega = i_2(r_1(\omega')) = r_2(i_1(\omega'))$ and $\omega \in \text{im } r_2$.

CASE 2. If $p_2(\omega) \neq 0$, then there exists $\omega' \in \Omega^{p,q}(\Delta_n)$ such that $p_1(\omega') = p_2(\omega)$. Then

$$\begin{aligned}
 p_2(\omega - r_2(\omega')) &= p_1(\omega)p_2(r_2(\omega')) \\
 &= p_1(\omega') - p_2(r_2(\omega')) \\
 &= 0,
 \end{aligned} \tag{5.32}$$

thus $\omega' - r_2(\omega') \in \ker p_2$, then $\omega' - r_2(\omega') \in \text{im } r_2$ (by [Case 1](#)), then $\omega \in \text{im } r_2$.

Finally for $k = n - 1$, we have $\Delta_n^{n-1} = \partial \Delta_n$. Thus the sequence

$$\Omega^{p,q}(\Delta_n) \longrightarrow \Omega^{p,q}(\partial \Delta_n) \longrightarrow 0 \tag{5.33}$$

is exact. \square

Now we prove that, for any $q \geq 0$ the complex $\Omega^{*,q}(\Delta_n)$ has trivial cohomology. If $\lambda : \mathcal{T} \rightarrow \mathbb{Z}$ is any linear form with $\lambda(1) = 1$. For each $p \geq 0$, let $j_\lambda : \Omega^p(\mathcal{T}) \rightarrow \Omega^{p-1}(\mathcal{T})$ be the map defined at $\omega = f^0 df^1 \cdots df^p \in \Omega^p(\mathcal{T})$ by

$$j_\lambda(\omega) = \lambda(f^0) f^1 df^2 \cdots df^p - \lambda(f^0 f^1) df^2 \cdots df^p. \tag{5.34}$$

Then j_λ is a contracting homotopy ([Lemma 4.5](#)). Consider $\Omega^{p,q}(\Delta_n)$ as a submodule of $\Omega^p(\mathcal{T}_n)$ and j_λ restricted to $\Omega^{p,q}(\Delta_n)$. Suppose that

$$\omega = x^{\alpha_1} dx^{\epsilon_1} \cdots x^{\alpha_r} dx^{\epsilon_r} = \sum f^0 df^1 \cdots df^p. \tag{5.35}$$

Note that $\|J_\lambda(\omega)\|_j \leq \|\omega\|_j$ for all j . Therefore $\|J_\lambda(\omega)\| \leq \|\omega\| \leq q$ and $J_\lambda(\omega) \in \Omega^{p-1,q}(\Delta_n)$. Thus $J_\lambda : \Omega^{p,q}(\Delta_n) \rightarrow \Omega^{p-1,q}(\Delta_n)$ is a contracting homotopy (by [Lemma 4.5](#)).

If $\omega \in \Omega^{p,q}(\Delta_n)$ and $d\omega = 0$ we say that ω is *closed*. We say that ω is *exact* if there exists $\eta \in \Omega^{p-1,q}(\Delta_n)$ such that $\omega = d\eta$. Then we have the following lemma.

LEMMA 5.7 (the noncommutative tame Poincaré lemma). *If $\omega \in \Omega^{p,q}(\Delta_n)$ is a closed form, then ω is exact.*

6. Integration of noncommutative tame forms. In this section, we introduce integration of noncommutative tame forms, prove the tame noncommutative Stokes' theorem, and use this result to define a morphism of \mathbb{Q}_q -modules $I : \Omega^{*,q}(\Delta_n) \rightarrow C^*(\Delta_n; \mathbb{Q}_q)$ which plays an important role in the proof of the de Rham theorem. The definition of the integral of noncommutative tame forms is motivated by the ideas presented in [\[12\]](#).

Let $\Omega^{*,*}(\Delta_n)$ denote the algebra of noncommutative tame differential forms in the variables x_0, x_1, \dots, x_n .

Let $T^{*,*}(\Delta_n)$ be the algebra of differential forms of Cenkl-Porter with \mathbb{Q}_q -coefficients on the standard cube $I^{n+1} \subset \mathbb{R}^{n+1}$. Define $F : \Omega^{p,q}(\Delta_n) \rightarrow T^{p,q}(\Delta_n)$ as follows: if $\omega \in \Omega^{0,q}(\Delta_n)$ or $\omega = dx_j \in \Omega^{1,1}(\Delta_n)$ then $F(\omega) = \omega$; if $\omega = f^0 df^1 \cdots df^p \in \Omega^{p,q}(\Delta_n)$ for $p > 1$, then $F(f^0 df^1 \cdots df^p) = f^0 df^1 \wedge \cdots \wedge df^p$. Then $F : \Omega^{p,q}(\Delta_n) \rightarrow T^{p,q}(\Delta_n)$ defines a morphism of \mathbb{Q}_q -modules.

Note that for all $p \geq 0$, we have

$$F(f^0 df^1 \cdots df^p) = F(f^0) F(df^1) \wedge \cdots \wedge F(df^p). \quad (6.1)$$

In particular, if $\omega \in \Omega^{p_1,q_1}(\Delta_n)$ and $\eta \in \Omega^{p_2,q_2}(\Delta_n)$ then

$$F(\omega \cdot \eta) = F(\omega) \wedge F(\eta). \quad (6.2)$$

In other words F is a morphism of algebras. To prove this identity it is enough to consider $\omega = f^0 df^1$, $\eta = g^0 dg^1 \in \Omega^1(\mathcal{T}_n)$. Then

$$\begin{aligned} F(\omega \cdot \eta) &= F(f^0 df^1 \cdot g^0 dg^1) \\ &= F(f^0 d(f^1 g^0) dg^1 - f^0 f^1 dg^0 dg^1) \\ &= f^0 d(f^1 g^0) \wedge dg^1 - f^0 f^1 dg^0 dg^1 \\ &= f^0 df^1 g^0 \wedge dg^1 + f^0 f^1 dg^0 \wedge dg^1 - f^0 f^1 dg^0 \wedge dg^1 \\ &= f^0 g^0 df^1 \wedge dg^1 \\ &= F(\omega) \wedge F(\eta). \end{aligned} \quad (6.3)$$

Similarly, the following propositions can be proved by direct computations.

PROPOSITION 6.1. *The diagram*

$$\begin{array}{ccc} \Omega^{p,q}(\Delta_n) & \xrightarrow{d} & \Omega^{p,q}(\Delta_n) \\ F_p \downarrow & & \downarrow F_{p+1} \\ T^{p,q}(\Delta_n) & \xrightarrow{d} & T^{p,q}(\Delta_n) \end{array} \quad (6.4)$$

commutes for all $p \geq 0$.

PROPOSITION 6.2. *Let $\omega = f^0 df^1 \cdots df^p \in \Omega^{p,q}(\Delta_n)$. If $p > n$ then $F(\omega) = 0$. If $0 < p \leq n$ and \mathcal{G}_p denotes the permutation group of the set $\{1, 2, \dots, p\}$, then*

$$F(\omega) = \sum_{0 \leq j_1 < \cdots < j_p \leq n} \sum_{\tau \in \mathcal{G}_p} \operatorname{sgn} \tau f^0 \frac{\partial f^1}{\partial x_{j_{\tau(1)}}} \cdots \frac{\partial f^p}{\partial x_{j_{\tau(p)}}} dx_{j_1} \wedge \cdots \wedge dx_{j_p}. \quad (6.5)$$

PROPOSITION 6.3. *Suppose that $\omega = x^{\alpha_1} dx^{\varepsilon_1} \cdots x^{\alpha_r} dx^{\varepsilon_r} \in \Omega_n^{p,q}(\Delta_n)$ and $0 < p \leq n$, then*

- (1) *if $\sum_i \varepsilon_{ij} \geq 2$ for some j , then $F(\omega) = 0$;*
- (2) *if $0 \leq i_1 < \cdots < i_p \leq n$ and $\tau \in \mathcal{G}_p$ such that $\tau(1) < \cdots < \tau(p)$ and $\sum_i \varepsilon_{i\tau(j)} \leq 1$ for all j , then*

$$F(\omega) = \operatorname{sgn} \tau x^{\alpha_1} \cdots x^{\alpha_r} dx_{i_{\tau(1)}} \wedge \cdots \wedge dx_{i_{\tau(p)}}. \quad (6.6)$$

REMARK 6.4. *Proposition 6.5 implies that F is a simplicial map.*

From [Proposition 3.1](#), we obtain the following result.

PROPOSITION 6.5. *Let $p \geq 0, q \geq 1$. If $\omega \in \Omega_n^{p,q}(\mathbb{Z})$ and if G is a p -face of Δ_n , then*

$$\int_G F(\omega) \in \mathbb{Q}_q. \quad (6.7)$$

If $\omega = \Omega^{p,q}(\Delta_n)$ and $\sigma : \Delta_p \rightarrow \Delta_n$ is a p -simplex, we define *the integral of ω on σ* by

$$\int_{\sigma} \omega = \int_{\sigma} F(\omega) = \int_{\Delta_p} \sigma^* F(\omega). \quad (6.8)$$

If $\sigma = \sum_i \sigma_i \otimes a_i \in C_p(\Delta_n; \mathbb{Q}_q)$, then *the integral of ω on σ* is defined by

$$\int_{\sigma} \omega = \sum_i a_i \int_{\sigma_i} \omega. \quad (6.9)$$

PROPOSITION 6.6 (noncommutative Stokes' theorem). *Let σ be a p -chain on Δ_n and let $\omega \in \Omega^{p,q}(\Delta_n)$. Then*

$$\int_{\sigma} d\omega = \int_{\partial\sigma} \omega. \quad (6.10)$$

PROOF. Let $\omega \in \Omega^{p,q}(\Delta_n)$ and let $\sigma : \Delta_p \rightarrow \Delta_n$ be a p -simplex. By [Proposition 6.1](#) and by the classical Stokes' theorem, we get

$$\int_{\sigma} d\omega = \int_{\sigma} F(d\omega) = \int_{\sigma} d(F(\omega)) = \int_{\partial\sigma} F(\omega) = \int_{\partial\sigma} \omega. \quad (6.11)$$

□

Let $(C^*(\Delta_n; \mathbb{Q}_q), \delta)$ denote the standard complex of cochains on Δ_n with coefficients in \mathbb{Q}_q . Let

$$I : \Omega^{*,q}(\Delta_n) \longrightarrow C^*(\Delta_n; \mathbb{Q}_q) \quad (6.12)$$

be the morphism of \mathbb{Q}_q -modules defined as follows: given $\sigma \in C_p(\Delta_n; \mathbb{Q}_q)$ and $\omega \in \Omega^{p,q}(\Delta_n)$,

$$I_p(\omega)(\sigma) = \int_{\sigma} \omega. \quad (6.13)$$

The Stokes theorem implies that I is a map of cochain complexes. We also have that the diagram

$$\begin{array}{ccccc} \Omega^{p,q}(\Delta_{n-1}) & \xleftarrow{\delta_i} & \Omega^{p,q}(\Delta_n) & \xrightarrow{s_i} & \Omega^{p,q}(\Delta_{n+1}) \\ I \downarrow & & I \downarrow & & I \downarrow \\ C^p(\Delta_{n-1}; \mathbb{Q}_q) & \xleftarrow{\tilde{\delta}_i} & C^p(\Delta_n; \mathbb{Q}_q) & \xrightarrow{\tilde{s}_i} & C^p(\Delta_{n+1}; \mathbb{Q}_q) \end{array} \quad (6.14)$$

commutes for $0 \leq i \leq n$. Then I is a simplicial map.

PROPOSITION 6.7. *The diagram*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega^{p,q}(\Delta_K, \partial\Delta_K) & \xrightarrow{i} & \Omega^{p,q}(\Delta_K) & \xrightarrow{r} & \Omega^{p,q}(\partial\Delta_{K-1}) \longrightarrow 0 \\ & & I \downarrow & & I \downarrow & & I \downarrow \\ 0 & \longrightarrow & C^p(\Delta_K, \partial\Delta_K; \mathbb{Q}_q) & \xrightarrow{i} & C^p(\Delta_K; \mathbb{Q}_q) & \xrightarrow{r} & C^p(\partial\Delta_{K-1}; \mathbb{Q}_q) \longrightarrow 0 \end{array} \quad (6.15)$$

commutes for all $p \geq 0$, $q \geq 1$. (The i 's and r 's denote the inclusions and restrictions, respectively.)

PROOF. Let $\sigma \in C_p(\Delta_K; \mathbb{Q}_q)$ and $\omega \in \Omega^{p,q}(\Delta_K, \partial\Delta_K)$, then $r(\omega) = r(i(\omega))$. Therefore,

$$\begin{aligned} I(i(\omega))(\sigma) &= \int_{\sigma} i(\omega) = \int_{\sigma \cap \partial\Delta_K} \omega + \int_{\sigma - (\sigma \cap \partial\Delta_K)} \omega \\ &= \int_{\partial - (\sigma \cap \partial\Delta_K)} \omega = i(I(\omega))(\sigma). \end{aligned} \quad (6.16)$$

On the other hand, we have

$$\begin{aligned} I(r(\omega))(\sigma) &= I(\sigma)(r(\omega)) = \int_{\sigma} r(\omega) = \int_{\sigma} \omega|_{\partial\Delta_K} \\ &= \int_{\sigma \cap \partial\Delta_K} \omega = \int_{r(\sigma)} \omega = I(r(\sigma))(\omega). \end{aligned} \quad (6.17)$$

□

7. The noncommutative tame de Rham theorem for cohomology. In this section, we introduce the noncommutative de Rham complex of Cenkl and Porter for a simplicial set of finite type X . Then we use the noncommutative versions of the Poincaré lemma and the Stokes' theorem to prove the noncommutative tame de Rham theorem.

Let X be a simplicial complex of finite type. Let X_n be the collection of nondegenerated n -simplices in X . A *noncommutative differential form of type (p, q) on X* is a simplicial map $\omega : X_n \rightarrow \Omega^{p,q}(\Delta_n)$ (in other words, ω is a map such that for $G \in X_n$ and any face F of G , $\omega(F)$ is the restriction of $\omega(G)$ to F). The collection of all such forms is denoted by $\Omega^{p,q}(X)$.

For a p -chain $\sigma = \sum_i \sigma_i \otimes a_i \in C_p(X; \mathbb{Q}_q)$, $\sigma_i : \Delta_p \rightarrow X$ and $\omega \in \Omega^{p,q}(X)$, we define

$$\int_{\sigma} \omega = \sum_i a_i \int_{\Delta_p} \omega|_{\sigma_i}, \quad (7.1)$$

so we may define the map $I : \Omega^{p,q}(X) \rightarrow C^p(X; \mathbb{Q}_q)$ by

$$I_p(\omega)(\sigma) = \int_{\sigma} \omega. \quad (7.2)$$

Then

$$\delta I_p(\omega)(\sigma) = I_p(\omega)(\partial \sigma) = \int_{\partial \Delta_p} \omega|_{\partial \sigma}. \quad (7.3)$$

On the other hand,

$$I_{p+1}(d\omega)(\sigma) = I_p(d\omega)(\sigma) = \int_{\Delta_p} d\omega|_{\sigma}. \quad (7.4)$$

Thus integration induces a map of cochain complexes. Then we have the following theorem.

THEOREM 7.1. *Let X be a simplicial set of finite type. Then for $q \geq 1$ the map*

$$I : H^i(\Omega^{*,q}(X)) \xrightarrow{\cong} H^i(X; \mathbb{Q}_q), \quad (7.5)$$

induced by integration, is an isomorphism of \mathbb{Q}_q -modules for all $i \geq 0$.

PROOF. Induction on the skeleta of X . For $k = 0$ the statement is true because

$$H^i(\Omega^{*,q}(X)) = \begin{cases} \mathbb{Q}_q, & \text{if } i = 0, \\ 0, & \text{if } i > 0, \end{cases} \quad H^i(X; \mathbb{Q}_q) = \begin{cases} \mathbb{Q}_q, & \text{if } i = 0, \\ 0, & \text{if } i > 0. \end{cases} \quad (7.6)$$

Suppose that the statement is true on the ℓ -skeleton, X_ℓ of X for $\ell < k$.

From [Proposition 6.7](#) it follows that the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega^{*,q}(X_k, X_{k-1}) & \longrightarrow & \Omega^{*,q}(X_k) & \longrightarrow & \Omega^{*,q}(X_{k-1}) \longrightarrow 0 \\ & & I \downarrow & & I \downarrow & & I \downarrow \\ 0 & \longrightarrow & C^*(X_k, X_{k-1}; \mathbb{Q}_q) & \longrightarrow & C^*(X_k; \mathbb{Q}_q) & \longrightarrow & C^*(X_{k-1}; \mathbb{Q}_q) \longrightarrow 0 \end{array} \quad (7.7)$$

is commutative. Then the following diagram commutes:

$$\begin{array}{ccccccc}
\cdots & \longrightarrow & H^{i-1}(\Omega^{*,q}(X_{k-1})) & \xrightarrow{\delta} & H^i(\Omega^{*,q}(X_k, X_{k-1})) & \xrightarrow{\delta} & H^i(\Omega^{*,q}(X_k)) \longrightarrow H^i(\Omega^{*,q}(X_{k-1})) \longrightarrow \cdots \\
& & \downarrow \iota & & \downarrow \kappa & & \downarrow I \\
\cdots & \longrightarrow & H^{i-1}(X_{k-1}; \mathbb{Q}_q) & \xrightarrow{\delta} & H^i(X_k, X_{k-1}; \mathbb{Q}_q) & \xrightarrow{\delta} & H^i(X_k; \mathbb{Q}_q) \longrightarrow H^i(X_{k-1}; \mathbb{Q}_q) \longrightarrow \cdots
\end{array} \tag{7.8}$$

The rows are exact, an ι is an isomorphism by assumption. We prove that κ is an isomorphism. Let $\{\Delta_{k,j} : j \in J\}$ be the set of k -simplices of X_k . Then

$$\begin{aligned}
\Omega^{*,q}(X_k, X_{k-1}) &\cong \bigoplus_j \Omega^{*,q}(\Delta_{k,j}, \partial\Delta_{k,j}), \\
C^*(X_k, X_{k-1}; \mathbb{Q}_q) &\cong \bigoplus_j C^*(\Delta_{k,j}, \partial\Delta_{k,j}; \mathbb{Q}_q)
\end{aligned} \tag{7.9}$$

are isomorphisms of \mathbb{Q}_q -modules. Then it is enough to prove that integration induces an isomorphism

$$I : H^i(\Omega^{*,q}(\Delta_k, \partial\Delta_k)) \longrightarrow H^i(C^*(\Delta_k, \partial\Delta_k; \mathbb{Q}_q)). \tag{7.10}$$

Consider the following commutative diagram (Proposition 6.7):

$$\begin{array}{ccccccc}
0 & \longrightarrow & \Omega^{*,q}(\Delta_k, \partial\Delta_k) & \longrightarrow & \Omega^{*,q}(\Delta_k) & \longrightarrow & \Omega^{*,q}(\partial\Delta_k) \longrightarrow 0 \\
& & \downarrow I & & \downarrow I & & \downarrow I \\
0 & \longrightarrow & C^*(\Delta_k, \partial\Delta_k; \mathbb{Q}_q) & \longrightarrow & C^*(\Delta_k; \mathbb{Q}_q) & \longrightarrow & C^*(\partial\Delta_k; \mathbb{Q}_q) \longrightarrow 0.
\end{array} \tag{7.11}$$


The first row is exact by Proposition 5.6.

$$\begin{array}{ccccccc}
\cdots & \longrightarrow & H^{i-1}(\Omega^{*,q}(\partial\Delta_k)) & \xrightarrow{\delta} & H^i(\Omega^{*,q}(\Delta_k, \partial\Delta_k)) & \xrightarrow{\delta} & H^i(\Omega^{*,q}(\Delta_k)) \longrightarrow H^i(\Omega^{*,q}(\partial\Delta_k)) \longrightarrow \cdots \\
& & \downarrow \iota & & \downarrow \kappa & & \downarrow I \\
\cdots & \longrightarrow & H^{i-1}(\partial\Delta_k; \mathbb{Q}_q) & \xrightarrow{\delta} & H^i(\Delta_k, \partial\Delta_k; \mathbb{Q}_q) & \xrightarrow{\delta} & H^i(\Delta_k; \mathbb{Q}_q) \longrightarrow H^i(\partial\Delta_k; \mathbb{Q}_q) \longrightarrow \cdots
\end{array} \tag{7.12}$$

Therefore I is an isomorphism by lemma five (“so named because of the five-term exact sequence involved in its formulation,” Spanier [21, page 185]). \square

In the following examples, we consider X to be the circle S^1 and we verify the isomorphism $H^*(\Omega^{*,q}(X)) \cong H^*(X; \mathbb{Q}_q)$ (for $q = 1, 2$) by computing directly $H^i(\Omega^{*,q}(X))$.

EXAMPLE 7.2. In this example, we consider the complex of noncommutative tame differential forms of weight ≤ 1 on the circle $\Omega^{*,2}(S^1)$, where $\Omega^{*,2}(S^1)$ denotes the complex of noncommutative differential forms of weight ≤ 2 on the circle. In this case the ground ring is $\mathbb{Q}_2 = \mathbb{Z}[1/2]$.

FIGURE 7.1. Differential forms on S^1 .

Consider the triangulation of S^1 with 0-simplexes v_0, v_1, v_2 , and 1-simplexes e_0, e_1, e_2 oriented as indicated in Figure 7.1. We can consider each 1-simplex e_i is the image of a continuous injective function $\psi_i : \Delta_1 \rightarrow S^1$. Let $\phi_i = \psi_i^{-1} : e_i \rightarrow \Delta_1$, $i = 0, 1, 2$. Recall that the 1-simplex Δ_1 is considered to be the union of the two backfaces of the 2-cube I^2 in \mathbb{R}^2 , as in Example 2.4.

A form $\eta \in \Omega^{0,2}(S^1)$ is a map such that $\eta(e_i) = \eta_i \in \Omega^{0,2}(\Delta_1) = \Omega_1^{0,2}(\mathbb{Z}) \otimes \mathbb{Q}_2$, satisfying

$$\begin{aligned} \eta_0(\phi_0(v_0)) &= \eta_1(\phi_1(v_0)), \\ \eta_1(\phi_1(v_1)) &= \eta_2(\phi_2(v_1)), \\ \eta_2(\phi_2(v_2)) &= \eta_0(\phi_0(v_2)). \end{aligned} \tag{7.13}$$

Note that $x_0 x_1 = y_0 y_1 = z_0 z_1 = 0$, therefore we have

$$\begin{aligned} \eta_0(x_0, x_1) &= k_0 + a_0 x_0 + a_1 x_1 + a_2 x_0^2 + a_3 x_1^2, \\ \eta_1(y_0, y_1) &= k_1 + b_0 y_0 + b_1 y_1 + b_2 y_0^2 + b_3 y_1^2, \\ \eta_2(z_0, z_1) &= k_2 + c_0 z_0 + c_1 z_1 + c_2 z_0^2 + c_3 z_1^2, \end{aligned} \tag{7.14}$$

with $k_i, a_j, b_j, c_j \in \mathbb{Q}_2$ for $i = 0, 1, 2$ and $j = 0, 1, 2, 3$, such that

$$\begin{aligned} k_0 + a_0 + a_2 &= k_1 + b_1 + b_3, \\ k_1 + b_0 + b_2 &= k_2 + c_1 + c_3, \\ k_2 + c_0 + c_2 &= k_0 + a_1 + a_3. \end{aligned} \tag{7.15}$$

If η is closed then $d\eta_i = 0$ for $i = 0, 1, 2$, then $a_j = b_j = c_j = 0$. Then (7.15) implies $k_0 = k_1 = k_2 = 0$ and η is constant. Then $H^0(\Omega^{*,2}(S^1)) \cong \mathbb{Q}_2$.

Now let $\omega \in \Omega^{1,2}(S^1)$. Then $\omega(e_i) = \omega_i \in \Omega^{1,2}(\Delta_1)$. We have

$$\begin{aligned} \omega_0(x_0, x_1) &= a_0 dx_0 + a_1 dx_1 + a_2 x_0 dx_0 + a_3 dx_0 x_0 + a_4 x_1 dx_1 + a_5 dx_1 x_1, \\ \omega_1(y_0, y_1) &= b_0 dy_0 + b_1 dy_1 + b_2 y_0 dy_0 + b_3 dy_0 y_0 + b_4 y_1 dy_1 + b_5 dy_1 y_1, \\ \omega_2(z_0, z_1) &= c_0 dz_0 + c_1 dz_1 + c_2 z_0 dz_0 + c_3 dz_0 z_0 + c_4 z_1 dz_1 + c_5 dz_1 z_1. \end{aligned} \tag{7.16}$$

Then

$$\begin{aligned} d\omega_0 &= a_2 dx_0^2 - a_3 dx_0^2 + a_4 dx_1^2 - a_5 dx_1^2, \\ d\omega_1 &= b_2 dy_0^2 - b_3 dy_0^2 + b_4 dy_1^2 - b_5 dy_1^2, \\ d\omega_2 &= c_2 dz_0^2 - c_3 dz_0^2 + c_4 dz_1^2 - c_5 dz_1^2. \end{aligned} \quad (7.17)$$

If ω is closed we have $a_i = a_{i+1}$, $b_i = b_{i+1}$, and $c_i = c_{i+1}$, for $i = 2, 4$. Then we have $\dim \ker(d : \Omega^{1,2}(S^1) \rightarrow \Omega^{2,2}(S^1)) = 12$.

Note that the linear system (7.15) is equivalent to

$$\begin{aligned} k_0 - k_1 &= b_1 + b_3 - a_0 - a_2, \\ k_1 - k_2 &= c_1 + c_3 - b_0 b_2, \\ 0 &= a_0 - a_1 + a_2 - a_3 + b_0 - b_1 + b_2 - b_3 + c_0 - c_1 + c_2 - c_3. \end{aligned} \quad (7.18)$$

Then $\dim \text{im}(d : \Omega^{0,2}(S^1) \rightarrow \Omega^{1,2}(S^1)) = 11$ and $H^1(\Omega^{*,2}(S^1)) \cong \mathbb{Q}_2$.

Now let $\theta \in \Omega^{2,2}(S^1)$, then θ is a map such that $\theta(e_i) = \theta_i \in \Omega^{2,2}(\Delta_1)$, that is,

$$\begin{aligned} \theta_0(x_0, x_1) &= a_0 dx_0^2 + a_1 dx_1^2, \\ \theta_1(y_0, y_1) &= b_0 dy_0^2 + b_1 dy_1^2, \\ \theta_2(z_0, z_1) &= c_0 dz_0^2 + c_1 dz_1^2. \end{aligned} \quad (7.19)$$

Any such form θ is closed. We also have that θ is exact, in fact $\theta = d\omega$ where $\omega \in \Omega^{2,2}(S^1)$ is given by (note that 2 is invertible in \mathbb{Q}_2)

$$\begin{aligned} \omega_0(x_0, x_1) &= \frac{a_0}{2} x_0 dx_0 - \frac{a_0}{2} dx_0 x_0 + \frac{a_1}{2} x_1 dx_1 - \frac{a_1}{2} dx_1 x_1, \\ \omega_1(y_0, y_1) &= \frac{b_0}{2} y_0 dy_0 - \frac{b_0}{2} dy_0 y_0 + \frac{b_1}{2} y_1 dy_1 - \frac{b_1}{2} dy_1 y_1, \\ \omega_2(z_0, z_1) &= \frac{c_0}{2} z_0 dz_0 - \frac{c_0}{2} dz_0 z_0 + \frac{c_1}{2} z_1 dz_1 - \frac{c_1}{2} dz_1 z_1. \end{aligned} \quad (7.20)$$

Then $H^2(\Omega^{*,2}(S^1)) = 0$. Finally, $H^i(\Omega^{*,2}(S^1)) = 0$ for $i > 2$ because $\Omega^{p,2}(S^1) = 0$ for $p > 2$.

We conclude this section with the presentation of a more general version of the noncommutative tame de Rham theorem (Theorem 7.1). Let M be a \mathbb{Q}_q -module and let $\omega = \eta \otimes a \in \Omega^{p,q}(X) \otimes M$, and $\sigma = \theta \otimes b \in C_p(X; \mathbb{Q}_q) \otimes M$. Then the *integral of ω on σ* is defined by

$$\int_{\sigma} \omega = I(\eta)(\theta) \cdot a \otimes b = \int_{\theta} \eta \cdot a \otimes b. \quad (7.21)$$

Thus integration defines a morphism of modules \mathbb{Q}_q -modules $I : \Omega^{p,q}(X) \otimes M \rightarrow C^p(X; M)$. Finally, we apply Künneth's theorem and Theorem 7.1 to obtain the noncommutative tame de Rham theorem for cohomology.

THEOREM 7.3. *Let X be a simplicial set of finite type. Then for $q \geq 1$ and any \mathbb{Q}_q -module M there is a natural isomorphism of \mathbb{Q}_q -modules*

$$H^i(\Omega^{*,q}(X), M) \xrightarrow{\cong} H^i(X; M) \quad (7.22)$$

for all $i \geq 0$. The isomorphism is induced by integration.

Let $f : M_1 \rightarrow M_2$ be a morphism of \mathbb{Q}_q -modules. Then for each $p \geq 0$, f induces two morphisms of \mathbb{Q}_q -modules $f^* : H^i(\Omega^{p,q}(X), M_1) \rightarrow H^i(\Omega^{p,q}(X), M_2)$ and $f^* : H^i(X; M_1) \rightarrow H^i(X; M_2)$. The word “natural” in [Theorem 7.3](#) means that if $f : M_1 \rightarrow M_2$ is a morphism of \mathbb{Q}_q -modules then the diagram

$$\begin{array}{ccc} H^i(\Omega^{p,q}(X), M_1) & \xrightarrow{I} & H^i(X; M_2) \\ f^* \downarrow & & \downarrow f^* \\ H^i(\Omega^{p,q}(X), M_1) & \xrightarrow{I} & H^i(X; M_2) \end{array} \quad (7.23)$$

commutes for all $p \geq 0$, $i \geq 0$.

The existence of an isomorphism $H^i(\Omega^*(X)) \rightarrow H^i(X; \Lambda)$, for any commutative ring with a unit Λ , can be obtained following Cartan’s ideas. Such a proof is found in [\[3, 4\]](#). Cenkl’s (1998) noncommutative de Rham theorem is a generalization of a result of Karoubi proved in [\[12\]](#). In that same paper Karoubi conjectured the noncommutative de Rham theorem using integration being Λ a commutative ring containing the rationals.

8. The dual noncommutative complex of Cenkl-Porter. In [\[20\]](#), Scheerer et al. presented a dual version of the tame de Rham theorem. They introduced the chain complex of tame de Rham currents and proved the tame de Rham theorem for homology. In this section, we study the dual of the complex of noncommutative tame forms $\Omega_{*,*}(X)$ on a simplicial set of finite type X . We use the facts that X is a simplicial set of finite type and that $\Omega_n^{p,q}$ is a finitely generated free \mathbb{Z} -module for $p \geq 0$, $q \geq 0$, $n > 0$, and some classical results to prove some basic properties of the complex $\Omega_{*,*}(X)$. Then we prove that for $q \geq 1$ there exists an isomorphism of \mathbb{Q}_q -modules $H_i(X; \mathbb{Q}_q) \cong H_i(\Omega_{*,q}(X))$ for all $i \geq 0$.

Let $\Omega^{p,q}(\Delta_n) = \Omega_n^{p,q} \otimes_{\mathbb{Z}} \mathbb{Q}_q$ be the collection of noncommutative tame p -forms of weight $\leq q$ on the n -simplex Δ_n . Define

$$\Omega_{p,q}(\Delta_n) := \text{Hom}(\Omega^{p,q}(\Delta_n), \mathbb{Q}_q). \quad (8.1)$$

Elements of $\Omega_{p,q}(\Delta_n)$ are called *noncommutative tame de Rham currents*.

Consider the maps

$$\begin{aligned} \Omega_{p,q}(\Delta_n) &\xrightarrow{\partial} \Omega_{p-1,q}(\Delta_n), & \Omega_{p,q}(\Delta_{n-1}) &\xrightarrow{\delta^i} \Omega_{p,q}(\Delta_n), \\ & & \Omega_{p,q}(\Delta_{n+1}) &\xrightarrow{s^i} \Omega_{p,q}(\Delta_n), \end{aligned} \quad (8.2)$$

where $\partial = \tilde{d}$, $\delta^i = \tilde{s}_i$, and $s^i = \tilde{s}_i$ denote the dual of the maps

$$\begin{aligned} \Omega^{p,q}(\Delta_n) &\xrightarrow{d} \Omega^{p+1,q}(\Delta_n) \quad (\text{the differential}), \\ \Omega^{p,q}(\Delta_n) &\xrightarrow{\delta_i} \Omega^{p,q}(\Delta_{n-1}) \quad (\text{the face operators}), \\ \Omega^{p,q}(\Delta_n) &\xrightarrow{s_i} \Omega^{p,q}(\Delta_{n+1}) \quad (\text{the degeneracy operators}). \end{aligned} \quad (8.3)$$

Next we use the dual of the contracting homotopy $J_\lambda : \Omega^{p+1,q}(\Delta_n) \rightarrow \Omega^{p,q}(\Delta_n)$ (see [Lemma 5.7](#)) to prove the Poincaré lemma for noncommutative de Rham currents.

LEMMA 8.1. *For every $q \geq 1$,*

$$H_i(\Omega_{*,q}(\Delta_n)) = \begin{cases} \mathbb{Q}_q, & \text{for } i = 0, \\ 0, & \text{for } i > 0. \end{cases} \quad (8.4)$$

PROOF. The map $J_\lambda : \Omega^{p+1,q}(\Delta_n) \rightarrow \Omega^{p,q}(\Delta_n)$ is a contracting homotopy, that is, $dJ_\lambda + J_\lambda d = 1$ (see the proof of [Lemma 5.7](#)). Now consider the dual of $J_\lambda \tilde{J}_\lambda : \Omega_{p,q}(\Delta_n) \rightarrow \Omega_{p+1,q}(\Delta_n)$. For all $w \in \Omega_{p,q}(\Delta_n)$ and for any form $\omega \in \Omega^{p,q}(\Delta_n)$, we have

$$\begin{aligned} \partial \tilde{J}_\lambda(w)(\omega) + \tilde{J}_\lambda \partial(w)(\omega) &= (w)(J_\lambda d\omega) + (w)(dJ_\lambda \omega) \\ &= (w)(J_\lambda d(w) + dJ_\lambda(w)) = w(\omega), \end{aligned} \quad (8.5)$$

in other words, $\partial \tilde{J}_\lambda + \tilde{J}_\lambda \partial = 1$. Therefore, if $\partial(w) = 0$ there exists $v \in \Omega_{p+1,q}(\Delta_n)$ such that $\partial v = w$. Namely, $v = \tilde{J}(w)$. \square

Using [Proposition 5.6](#) and [[14](#), Theorem III.6.3] we deduce the following proposition.

PROPOSITION 8.2. *The sequence*

$$0 \rightarrow \Omega_{p,q}(\partial \Delta_k) \rightarrow \Omega_{p,q}(\Delta_k) \rightarrow \Omega_{p,q}(\Delta_k, \partial \Delta_k) \rightarrow 0 \quad (8.6)$$

is an exact sequence of \mathbb{Q}_q -modules for all $p \geq 0$ and $q \geq 1$.

Let X be a simplicial set of finite type. Let X_k denote the k -skeleton of X . Let $C_*(X)$ be the complex of unreduced chains of X (as an abelian group) and let

$$C_*(X; \mathbb{Q}_q) := C_*(X) \otimes_{\mathbb{Z}} \mathbb{Q}_q. \quad (8.7)$$

The *noncommutative tame de Rham currents of type (p, q) (of first kind)* on X are the elements of

$$\Omega_{p,q}(X) := \text{Hom}(\Omega^{p,q}(X), \mathbb{Q}_q). \quad (8.8)$$

For $q \geq 1$ define $I : C_p(X; \mathbb{Q}_q) \rightarrow \Omega_{p,q}(X)$ as follows: given $\sigma = \sum_\ell \sigma_\ell \otimes a_\ell \in C_p(X; \mathbb{Q}_q)$ and $\omega \in \Omega^{p,q}(X)$ then

$$I(\sigma)(\omega) = \int_\sigma \omega = \sum_\ell a_\ell \int_{\sigma_\ell} \omega. \quad (8.9)$$

Let $\{\Delta_{k,j} : j \in J\}$ be the set of simplexes of X_k . Because X is of finite type, we have

$$\Omega_{*,q}(X_k, X_{k-1}) = \text{Hom}(\Omega^{*,q}(X_k, X_{k-1}), \mathbb{Q}_q) = \bigoplus_j \Omega_{*,q}(\Delta_{k,j}, \partial\Delta_{k,j}). \quad (8.10)$$

Then as a consequence of [Proposition 8.2](#), we have the following result.

PROPOSITION 8.3. *The sequence*

$$0 \longrightarrow \Omega_{*,q}(X_{k-1}) \longrightarrow \Omega_{*,q}(X_k) \longrightarrow \Omega_{*,q}(X_k, X_{k-1}) \longrightarrow 0 \quad (8.11)$$

is an exact sequence of \mathbb{Q}_q -modules for all $q \geq 1$.

If X is a simplicial set of finite type, then there exists a natural isomorphism of \mathbb{Q}_q -modules (by [\[16, Theorem V.4.1\]](#))

$$C_p(X_k; \mathbb{Q}_q) \cong \text{Hom}(\text{Hom}(C_p(X_k), \mathbb{Q}_q), \mathbb{Q}_q) = \text{Hom}(C^p(X_k; \mathbb{Q}_q), \mathbb{Q}_q) \quad (8.12)$$

for all $k \geq 0, p \geq 0$. Let $\tilde{I} : C_p(\Delta_k; \mathbb{Q}_q) \rightarrow \Omega_{p,q}(\Delta_k)$ be dual to the morphism $I : \Omega^{p,q}(\Delta_k) \rightarrow C^p(\Delta_k; \mathbb{Q}_q)$. Then [Proposition 6.7](#) implies the following result.

PROPOSITION 8.4. *The diagram*

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_*(\partial\Delta_k; \mathbb{Q}_q) & \xrightarrow{\iota} & C_*(\Delta_k; \mathbb{Q}_q) & \xrightarrow{\rho} & C_p(\Delta_k, \partial\Delta_k; \mathbb{Q}_q) \longrightarrow 0 \\ & & \tilde{I} \downarrow & & \tilde{I} \downarrow & & \tilde{I} \downarrow \\ 0 & \longrightarrow & \Omega_{*,q}(\partial\Delta_k) & \xrightarrow{\tilde{\rho}} & \Omega_{*,q}(\Delta_k) & \xrightarrow{\tilde{\iota}} & \Omega_{*,q}(\Delta_k, \partial\Delta_k) \longrightarrow 0 \end{array} \quad (8.13)$$

commutes for all $q \geq 1, p \geq 0$ (ι and ρ denote inclusion and projection, respectively).

9. The noncommutative tame de Rham theorem for homology. In this section, we prove that the dual map I induces an isomorphism of \mathbb{Q}_q -modules between the homology of a simplicial set of finite type and the homology of the complex of non-commutative tame de Rham currents.

THEOREM 9.1. *Let X be a simplicial set of finite type. Then for $q \geq 1$, the map*

$$\tilde{I} : H_i(X; \mathbb{Q}_q) \xrightarrow{\cong} H_i(\Omega_{*,q}(X)), \quad (9.1)$$

induced by integration, is an isomorphism of \mathbb{Q}_q -modules for all $i \geq 0$.

PROOF. Induction on the skeleta of X . For $k = 0$, the statement is true because

$$H_i(X; \mathbb{Q}_q) = \begin{cases} \mathbb{Q}_q, & \text{if } i = 0, \\ 0, & \text{if } i > 0, \end{cases} \quad H_i(\Omega_{*,q}(X)) = \begin{cases} \mathbb{Q}_q, & \text{if } i = 0, \\ 0, & \text{if } i > 0. \end{cases} \quad (9.2)$$

Suppose that the proposition is true for the ℓ -skeleta X_ℓ , for $\ell < k$. Consider the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_p(X_{k-1}; \mathbb{Q}_q) & \longrightarrow & C_p(X_k; \mathbb{Q}_q) & \longrightarrow & C_p(X_k, X_{k-1}; \mathbb{Q}_q) \longrightarrow 0 \\ & & \tilde{I} \downarrow & & \tilde{I} \downarrow & & \tilde{I} \downarrow \\ 0 & \longrightarrow & \Omega_{p,q}(X_{k-1}) & \longrightarrow & \Omega_{p,q}(X_k) & \longrightarrow & \Omega_{p,q}(X_k, X_{k-1}) \longrightarrow 0. \end{array} \quad (9.3)$$

The first row is exact. The second row is exact by [Proposition 8.3](#). Then we have the following commutative diagram, where the rows are exact

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_i(X_{k-1}; \mathbb{Q}_q) & \xrightarrow{\delta} & H_i(X_k; \mathbb{Q}_q) & \xrightarrow{\delta} & H_i(X_k, X_{k-1}; \mathbb{Q}_q) \longrightarrow H_{i-1}(X_{k-1}; \mathbb{Q}_q) \longrightarrow \cdots \\ & & \downarrow \iota & & \downarrow \kappa & & \downarrow \bar{I} \\ \cdots & \longrightarrow & H_i(\Omega_{*,q}(X_{k-1})) & \xrightarrow{\delta} & H_i(\Omega_{*,q}(X_k)) & \xrightarrow{\delta} & H_i(\Omega_{*,q}(x_k, X_{k-1})) \longrightarrow H_i(\Omega_{*,q}(X_{k-1})) \longrightarrow \cdots \end{array} \quad (9.4)$$

The ι 's are isomorphisms by induction hypothesis. We prove that κ is an isomorphism. Let $\{\Delta_{k,j} : j \in J\}$ be the k -simplices of X_k . Then the morphisms of \mathbb{Q}_q -modules

$$\begin{aligned} \Omega_{*,q}(X_k, X_{k-1}) &\cong \bigoplus_j \Omega_{*,q}(\Delta_{k,j}, \partial\Delta_{k,j}), \\ C_*(X_k, X_{k-1}; \mathbb{Q}_q) &\cong \bigoplus_j C_*(\Delta_{k,j}, \partial\Delta_{k,j}; \mathbb{Q}_q) \end{aligned} \quad (9.5)$$

are isomorphism of \mathbb{Q}_q -modules.

Then it is enough to prove that integration induces an isomorphism

$$\bar{I} : H_i(\Delta_k, \partial\Delta_k; \mathbb{Q}_q) \longrightarrow H_i(\Omega_{*,q}(\Delta_k, \partial\Delta_k)). \quad (9.6)$$

By [Proposition 8.4](#) the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_*(\partial\Delta_k; \mathbb{Q}_q) & \xrightarrow{\iota} & C_*(\Delta_k; \mathbb{Q}_q) & \xrightarrow{\rho} & C_*(\Delta_k, \partial\Delta_k; \mathbb{Q}_q) \longrightarrow 0 \\ & & \downarrow \bar{I} & & \downarrow \bar{I} & & \downarrow \bar{I} \\ 0 & \longrightarrow & \Omega_{*,q}(\partial\Delta_k) & \xrightarrow{\tilde{\rho}} & \Omega_{*,q}(\Delta_k) & \xrightarrow{\tilde{\iota}} & \Omega_{*,q}(\Delta_k, \partial\Delta_k) \longrightarrow 0 \end{array} \quad (9.7)$$

commutes. Then we have the following commutative diagram, with exact rows

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_i(\partial\Delta_k; \mathbb{Q}_q) & \xrightarrow{\delta} & H_i(\Delta_k; \mathbb{Q}_q) & \xrightarrow{\delta} & H_i(\Delta_k, \partial\Delta_k; \mathbb{Q}_q) \longrightarrow H_{i-1}(\partial\Delta_k; \mathbb{Q}_q) \longrightarrow \cdots \\ & & \downarrow \iota & & \downarrow \kappa & & \downarrow \bar{I} \\ \cdots & \longrightarrow & H_i(\Omega_{*,q}(\partial\Delta_k)) & \xrightarrow{\delta} & H_i(\Omega_{*,q}(\Delta_k)) & \xrightarrow{\delta} & H_i(\Omega_{*,q}(\Delta_k, \partial\Delta_k)) \longrightarrow H_i(\Omega_{*,q}(\partial\Delta_k)) \longrightarrow \cdots \end{array} \quad (9.8)$$

Then we apply five lemma and conclude that \bar{I} is an isomorphism. \square

Finally, we apply Künneth's theorem and obtain the noncommutative tame de Rham theorem for homology.

THEOREM 9.2. *Let X be a simplicial set of finite type. Then for any $q \geq 1$ and any \mathbb{Q}_q -module M , there is a natural isomorphism of \mathbb{Q}_q -modules*

$$H_i(X; M) \xrightarrow{\cong} H_i(\Omega_{*,q}(X), M) \quad (9.9)$$

for all $i \geq 0$. The isomorphism is induced by integration.

The word “natural” in the previous theorem means that for any \mathbb{Q}_q -modules M_1 and M_2 and for any morphism of \mathbb{Q}_q -modules $f : M_1 \rightarrow M_2$, then the diagram

$$\begin{array}{ccc} H_i(X; M_1) & \xrightarrow{\tilde{I}} & H_i(\Omega_{p,q}(X), M_1) \\ f_* \downarrow & & \downarrow f_* \\ H_i(X; M_2) & \xrightarrow{\tilde{I}} & H_i(\Omega_{p,q}(X), M_2) \end{array} \quad (9.10)$$

commutes for all $p \geq 0$, $i \geq 0$, where $f_* : H_i(X; M_1) \rightarrow H_i(X; M_2)$ and $f_* : H_i(\Omega_{p,q}(X), M_1) \rightarrow H_i(\Omega_{p,q}(X), M_2)$ are the morphisms of \mathbb{Q}_q -modules induced by f .

10. Cofiltered chain complexes and noncommutative tame de Rham currents.

Our study of noncommutative de Rham currents and the noncommutative tame de Rham theorem for homology was originally inspired by the investigations of Scheerer et al. in [20], but our approach differs from theirs. In this section, we introduce the *complex of noncommutative tame de Rham currents of second kind on a simplicial set X , $\mathcal{T}_{*,*}(X)$* which is a noncommutative version of the tame de Rham currents of Scheere et al. Then we show that, for all $q \geq 0$ there exists an isomorphism of \mathbb{Q}_q -modules $\Omega_{*,*}(X) \cong \mathcal{T}_{*,*}(X)$.

For the complex $F(X)$, such that for all n ,

$$F_n(X)_{p,q} = \begin{cases} 0, & \text{for } p > 0, \\ C_n(X) \otimes \mathbb{Q}_q, & \text{for } p = 0. \end{cases} \quad (10.1)$$

Note that $F_n(X)_{0,q-1} \otimes \mathbb{Q}_q = C_n(X) \otimes \mathbb{Q}_{q-1} \otimes \mathbb{Q}_q \cong C_n(X) \otimes \mathbb{Q}_q$. Then we can consider the respective identities as restrictions maps $\rho_q : F_n(X)_{0,q} \rightarrow F_n(X)_{0,q-1}$. Therefore, $F(X)$ is a cofiltered chain complex.

Consider $\Omega_{p,q}(\Delta_n) := \text{Hom}_{\mathbb{Z}}(\Omega^{p,q}(\Delta_n), \mathbb{Q}_q)$ and define the restriction maps $\rho_q : \Omega_{p,q}(\Delta_n) \rightarrow \Omega_{p,q-1}(\Delta_n) \otimes \mathbb{Q}_q$ as the dual of the inclusions $i_q : \Omega^{p,q-1}(\Delta_n) \rightarrow \Omega^{p,q}(\Delta_n)$ (Proposition 5.3). Then $\Omega_{*,*}^* = \bigoplus_{p,n \geq 0} (\cup_{q \geq 1} \Omega_{p,q}(\Delta_{n+1}))$ is a cosimplicial cofiltered chain complex. It is also a coalgebra; the coproduct is obtained by dualizing the multiplication of noncommutative differential forms

$$\Omega^{p_1, q_1}(\Delta_n) \otimes \Omega^{p_2, q_2}(\Delta_n) \xrightarrow{\mu} \Omega^{p_1 + p_2, q_1 + q_2}(\Delta_n), \quad (10.2)$$

μ is a noncommutative graded bilinear map of simplicial groups such that for all $0 \leq a_1 \leq q_1$, $0 \leq a_2 \leq q_2$

$$\begin{array}{ccc} \Omega^{p_1, a_1}(\Delta_n) \otimes \Omega^{p_2, a_2}(\Delta_n) & \xrightarrow{\mu} & \Omega^{p_1 + p_2, a_1 + a_2}(\Delta_n) \\ i \otimes i \downarrow & & \downarrow i \\ \Omega^{p_1, q_1}(\Delta_n) \otimes \Omega^{p_2, q_2}(\Delta_n) & \xrightarrow{\mu} & \Omega^{p_1 + p_2, q_1 + q_2}(\Delta_n). \end{array} \quad (10.3)$$

Recall that if $V = \{V_q, \mathbb{Q}_q\}$ and $W = \{W_q, \mathbb{Q}_q\}$ are two cofiltered chain complexes then the tensor product

$$(V \otimes W)_q = \varprojlim_{q_1 + q_2 \leq q} (V_{q_1} \otimes W_{q_2} \otimes \mathbb{Q}_q) \quad (10.4)$$

is a cofiltered chain complex.

Consider the cofiltered chain complex $\mathcal{T}(X)$ of *noncommutative tame de Rham currents of second kind* on X , where $\mathcal{T}_{p,q}(X)$ is defined as the quotient of

$$\bigoplus_{n \geq 0} (F_n(X) \otimes \Omega^n)_{p,q} \quad (10.5)$$

by the subspaces generated by the images of the maps

$$(\alpha_* \otimes 1 - 1 \otimes \alpha^*) : F_m(X) \otimes \Omega^k \rightarrow \bigoplus_{n \geq 0} (F_n(X) \otimes \Omega^n)_{p,q} \quad (10.6)$$

induced by all the morphisms $\alpha : \Delta(m) \rightarrow \Delta(k)$ in the category of simplicial sets Δ . For all n there exists a natural isomorphism

$$(F_n(X) \otimes \Omega^n)_{p,q} \cong C_n(X) \otimes \Omega_{p,q}^n. \quad (10.7)$$

Then $F(X)_{p,q}$ has a structure of simplicial cofiltered coalgebra.

The de Rham theorem for the complex $\mathcal{T}(X)$ follows from [Theorem 9.2](#) and from the following proposition.

PROPOSITION 10.1. *Let X be a simplicial set of finite type. Then for all $q \geq 0$ there is a natural isomorphism of \mathbb{Q}_q -modules*

$$\Omega_{*,q}(X) \cong \mathcal{T}_{*,q}(X) \quad \forall q \geq 1. \quad (10.8)$$

PROOF. First, we prove that the cochain complex $\Omega^{*,q}(X)$ is a naturally isomorphic to the cochain complex $\text{Hom}(\mathcal{T}_{*,q}(X), \mathbb{Q}_q)$. Let $\omega : X_* \rightarrow \Omega^{p,q}(\Delta_*)$ be a simplicial map (ω is a form on X). Let $\sigma \in X_n$ be an n -simplex and $w \in \Omega_{p,q}(\Delta_n) := \text{Hom}(\Omega^{p,q}(\Delta_n), \mathbb{Q}_q)$. Let $[\sigma \otimes w] \in \mathcal{T}_{p,q}(X)$ be the equivalence class containing $\sigma \otimes w$ (a current of second kind). Define a homomorphism $\zeta(\omega) : \mathcal{T}_{p,q}(X) \rightarrow \mathbb{Q}_q$ by

$$\zeta(\omega)([\sigma \otimes w]) = w(\omega(\sigma)). \quad (10.9)$$

We prove that $\zeta(\omega)$ is well defined: suppose that $\theta \otimes v \in [\sigma \otimes w]$. Then for any morphism $\alpha : \Delta(k) \rightarrow \Delta(m)$ in the category Δ there are elements $\eta \in X_m$ and $z \in \Omega^k$ such that

$$\theta \otimes v = \sigma \otimes w + (\alpha_* \otimes 1 - 1 \otimes \alpha^*)(\eta \otimes z) = \sigma \otimes w + \alpha_*(\eta) \otimes z - \eta \otimes \alpha^*(z). \quad (10.10)$$

Then

$$\begin{aligned} \zeta(\omega)([\theta \otimes v]) &= w(\omega(\sigma)) + z(\omega(\alpha^*(\eta))) - \alpha_*(z)(\omega(\eta)) \\ &= w(\omega(\sigma)) + z(\alpha^*(\omega(\eta))) - \alpha_*(z)(\omega(\eta)) \\ &= w(\omega(\sigma)) + \alpha_*(z)(\omega(\eta)) - \alpha_*(z)(\omega(\eta)) \\ &= w(\omega(\sigma)) \\ &= \zeta(\omega)([\sigma \otimes w]). \end{aligned} \quad (10.11)$$

Thus $\zeta : \Omega^{p,q}(X) \rightarrow \text{Hom}(\mathcal{T}_{p,q}(X), \mathbb{Q}_q)$ is a homomorphism. Note that

$$\zeta(d\omega)([\sigma \otimes \omega]) = w(d\omega(\sigma)) = \partial(w(\omega(\sigma))) = \partial\zeta(\omega)([\sigma \otimes \omega]). \quad (10.12)$$

Now we prove that ζ is injective. Suppose that $\omega, \eta \in \Omega^{p,q}(X)$ with $\zeta(\omega) = \zeta(\eta)$. Then for all n -simplex $\sigma \in X_n$ and $w \in \Omega_{p,q}(\Delta_n)$ we have $\zeta(\omega)([\sigma \otimes w]) = \zeta(\eta)([\sigma \otimes w])$. This is equivalent to $w(\omega(\sigma)) = w(\eta(\sigma))$ for all σ . Then $\omega(\sigma) = \eta(\sigma)$ for all σ . Hence $\omega = \eta$.

Now we prove that ζ is surjective. Let $\bar{w} : \mathcal{T}_{*,q}(X) \rightarrow \mathbb{Q}_q$ be a morphism of \mathbb{Q}_q -modules. Let $\sigma \in X_n$ be an n -simplex. Consider the homomorphism $\bar{w}(\sigma) : \Omega_{*,q}(\Delta_n) \rightarrow \mathbb{Q}_q$ defined by

$$\bar{w}(\sigma)(w) = \bar{w}([\sigma \otimes w]). \quad (10.13)$$

Thus $\bar{w}(\sigma) \in \text{Hom}(\Omega_{*,q}(\Delta_n), \mathbb{Q}_q)$. Then there exists a unique form $\omega(\sigma) \in \Omega^{*,q}(\Delta_n)$ such that

$$\bar{w}(\sigma)(w) = w(\omega(\sigma)), \quad (10.14)$$

$\varphi(\omega(\sigma)) = \bar{w}(\sigma)$, where φ is the natural isomorphism between a module and its dual (see [16, Theorem V.4.1]). Let $\alpha : \Delta(n) \rightarrow \Delta(k)$ be a morphism in the category Δ . Then $[\alpha_* \sigma \otimes z] = [\sigma \otimes \alpha^* z]$ for all $z \in \Omega_{p,q}(\Delta_k)$. Then

$$z(\omega(\alpha_* \sigma)) = \alpha^* z(\omega(\sigma)) = z(\alpha^*(\omega(\sigma))). \quad (10.15)$$

Thus the map $\sigma \mapsto \omega(\sigma)$ is a simplicial map $\omega : X_* \rightarrow \Omega^{p,q}(\Delta_*)$. Moreover,

$$\zeta(\omega)([\sigma \otimes w]) = w(\omega(\sigma)) = \bar{w}(\sigma)(w) = \bar{w}([\sigma \otimes w]), \quad (10.16)$$

and $\zeta(\omega) = \bar{w}$.

Finally we apply [16, Theorem V.4.1] to conclude the proof. \square

ACKNOWLEDGMENTS. This paper is essentially a part of the author's thesis written under supervision of Professor Bohumil Cenkl (Northeastern University), whom I gratefully thank for advice and encouragement. The author was supported by University of the Andes (Mérida and Trujillo, Venezuela) and the National Council for Scientific and Technological Researches (Caracas, Venezuela).

REFERENCES

- [1] P. Boullay, F. Kiefer, M. Majewski, M. Stelzer, H. Scheerer, M. Unsöld, and E. Vogt, *Tame Homotopy Theory via Differential Forms*, Freie Universität, Berlin, preprint No. 233, 1986.
- [2] H. Cartan, *Théories cohomologiques*, Invent. Math. 35 (1976), 261–271.
- [3] B. Cenkl, *Noncommutative differential forms and cohomology operations*, preprint, 1998.
- [4] ———, *Noncommutative Geometry*, Northeastern University, Boston, 1998, Course Notes.
- [5] B. Cenkl, G. Hector, and M. Saralegi, *Cohomologie d'intersection modérée. Un théorème de de Rham*, Pacific J. Math. 169 (1995), no. 2, 235–289 (French).
- [6] B. Cenkl and R. Porter, *Differential forms and torsion in the fundamental group*, Adv. in Math. 48 (1983), no. 2, 189–204.

- [7] ———, *de Rham theorem with cubical forms*, Pacific J. Math. **112** (1984), no. 1, 35–48.
- [8] A. Connes, *Noncommutative differential geometry*, Inst. Hautes Études Sci. Publ. Math. (1985), no. 62, 257–360.
- [9] ———, *Noncommutative Geometry*, Academic Press, California, 1994.
- [10] P. G. Goerss, *Simplicial chains over a field and p -local homotopy theory*, Math. Z. **220** (1995), no. 4, 523–544.
- [11] M. Karoubi, *Homologie cyclique et K -théorie*, Astérisque **149** (1987), 1–147 (French).
- [12] ———, *Formes différentielles non commutatives et cohomologie à coefficients arbitraires* [*Noncommutative differential forms and cohomology with arbitrary coefficients*], Trans. Amer. Math. Soc. **347** (1995), no. 11, 4277–4299 (French).
- [13] I. Kris, *p -adic homotopy theory*, Topology and Its Applications **52** (1993), 279–308.
- [14] S. Lang, *Algebra*, Addison-Wesley, Massachusetts, 1993.
- [15] D. Lehmann, *Théorie homotopique des formes différentielles (d'après D. Sullivan)*, Astérisque (1990), no. 45, 1–145 (French).
- [16] S. Mac Lane, *Categories for the Working Mathematician*, 2nd ed., Springer-Verlag, New York, 1998.
- [17] J. Madore, *An Introduction to Noncommutative Differential Geometry and Its Physical Applications*, 2nd ed., London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 1999.
- [18] J. P. Miller, *Simplicial Objects in Algebraic Topology*, Chicago Lectures in Mathematics, vol. 11, D. Van Nostrand, London, 1967.
- [19] C. Mouet, *q -cohomologie non commutative*, C. R. Acad. Sci. Paris Sér. I Math. **323** (1996), no. 8, 849–851.
- [20] H. Scheerer, K. Schuch, and E. Vogt, *Tame Homotopy Theory via de Rham Currents*, Freie Universität Berlin, Berlin, preprint No. A91-20, 1991.
- [21] E. H. Spanier, *Algebraic Topology*, McGraw-Hill Book, New York, 1966.
- [22] M. Sullivan, *A Comprehensive Introduction to Differential Geometry. Vol. I*, 3rd ed., Publish or Perish, Houston, 1999.
- [23] F. W. Whitney, *Foundations of Differentiable Manifolds and Lie Groups*, Springer-Verlag, New York, 1983.

LUIS FERNANDO MEJIAS: DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, 567 LAKE HALL, BOSTON, MA 02115, USA

E-mail address: lmejias@lynx.neu.edu

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.tero@usc.es