

ON THE FIRST POWER MEAN OF L -FUNCTIONS WITH THE WEIGHT OF GENERAL KLOOSTERMAN SUMS

ZHANG WENPENG

Received 20 January 2002

The main purpose of this paper is using the estimates for character sums and the analytic method to study the first power mean of Dirichlet L -functions with the weight of general Kloosterman sums, and give an interesting asymptotic formula.

2000 Mathematics Subject Classification: 11L05, 11M20.

1. Introduction. Let $q \geq 2$ be an integer, χ denotes a Dirichlet character modulo q . For any integers m and n , we define the general Kloosterman sums $S(m, n, \chi, q)$ as follows:

$$S(m, n, \chi, q) = \sum_{a=1}^q \chi(a) e\left(\frac{ma + n\bar{a}}{q}\right), \quad (1.1)$$

where \bar{a} denotes the inverse of a modulo q and $e(y) = e^{2\pi i y}$. This summation is very important, because it is a generalization of the classical Kloosterman sums. Many authors had studied the properties of $S(m, n, \chi, q)$. For instance, Chowla [1] and Malyšev [3] obtained a sharper upper bound estimation for $S(m, n, \chi, q)$. That is,

$$|S(m, n, \chi, p)| \ll (m, n, p)^{1/2} p^{1/2+\epsilon}, \quad (1.2)$$

where p is a prime, ϵ is any fixed positive number, and (m, n, p) denotes the greatest common divisor of m , n , and p . But for an arbitrary composite number q , we do not know how large $|S(m, n, \chi, q)|$ is. In fact the value of $|S(m, n, \chi, q)|$ is quite irregular if q is not a prime. The main purpose of this paper is to obtain some good distribution properties of $|S(m, n, \chi, q)|$ in some weight mean value problems. For convenience, in this paper we always suppose $q \geq 3$ be an integer and $L(s, \chi)$ denotes the Dirichlet L -function corresponding to character $\chi \bmod q$. Then we can use the estimates for character sums and the analytic method to prove the following main result.

THEOREM 1.1. *For any integers m and n with $(mn, q) = 1$, we have the asymptotic formula*

$$\sum_{\chi \neq \chi_0} |S(m, n, \chi, q)|^2 \cdot |L(1, \chi)| = C \cdot \phi^2(q) + O(q^{3/2} \cdot d^2(q) \cdot \ln^2 q), \quad (1.3)$$

where

$$C = \prod_p' \left[1 + \frac{\binom{2}{1}^2}{4^2 \cdot p^2} + \frac{\binom{4}{2}^2}{4^4 \cdot p^4} + \cdots + \frac{\binom{2n}{n}^2}{4^{2n} \cdot p^{2n}} + \cdots \right] \quad (1.4)$$

is a constant, $\sum_{\chi \neq \chi_0}$ denotes the summation over all nonprincipal characters modulo q , \prod'_p denotes the product over all primes p with $(p, q) = 1$, $d(q)$ is the divisor function, $\phi(q)$ is the Euler function, and $\binom{2n}{n} = (2n)!/(n!)^2$.

For general integer $k \geq 2$, whether there exists an asymptotic formula for

$$\sum_{\chi \neq \chi_0} |S(m, n, \chi, q)|^{2k} |L(1, \chi)| \quad (1.5)$$

is an unsolved problem.

2. Some lemmas. In order to complete the proof of [Theorem 1.1](#), we need the following lemmas.

LEMMA 2.1. *For any integer $q \geq 3$, we have the estimate*

$$\sum_{d|q} d^{1/2} \sum_{s=1}^{q/d-1} \left| \sum_{\chi \neq \chi_0} \chi(sd+1) |L(1, \chi)| \right| = O(q \cdot d(q) \cdot \ln^2 q). \quad (2.1)$$

PROOF. Let $N = q^{3/2}$, χ be a nonprincipal character mod q and $A(\chi, y) = \sum_{N < n \leq y} \chi(n)$. Then by Abel identity and Pólya-Vinogradov inequality, we have

$$\begin{aligned} L(1, \chi) &= \sum_{n \leq N} \frac{\chi(n)}{n} + \int_N^{+\infty} \frac{A(\chi, y)}{y^2} dy \\ &= \sum_{n \leq N} \frac{\chi(n)}{n} + O\left(\frac{\ln q}{q}\right). \end{aligned} \quad (2.2)$$

So that

$$|L(1, \chi)| = \left| \sum_{n \leq N} \frac{\chi(n)}{n} \right| + O\left(\frac{\ln q}{q}\right). \quad (2.3)$$

On the other hand, let $r(n)$ be a multiplicative function defining by

$$r(p^\alpha) = \frac{\binom{2\alpha}{\alpha}}{4^\alpha}, \quad r(1) = 1, \quad (2.4)$$

where p is a prime and α is any positive integer. For the function $r(n)$, it is easy to prove that

$$\sum_{d|n} r(d) \cdot r\left(\frac{n}{d}\right) = 1, \quad (2.5)$$

$$\begin{aligned} \left(\sum_{n \leq N} \frac{\chi(n)r(n)}{n} \right)^2 &= \sum_{m \leq N} \sum_{n \leq N} \frac{\chi(nm)r(m)r(n)}{mn} \\ &= \sum_{n \leq N} \frac{\chi(n)}{n} + \sum_{N < n \leq N^2} \frac{\chi(n)r(n, N)}{n}, \end{aligned} \quad (2.6)$$

where

$$r(n, N) = \sum_{\substack{d|n \\ d, n/d \leq N}} r(d) \cdot r\left(\frac{n}{d}\right). \quad (2.7)$$

Note that the triangle inequalities

$$|a| - |b| \leq |a \pm b|, \quad |a \pm b| \leq |a| + |b|, \quad (2.8)$$

from (2.3), (2.6), Cauchy inequality, and the orthogonality relationship for character sums

$$\begin{aligned} \sum_{\chi \bmod q} \chi(n) &= \begin{cases} \phi(q), & \text{if } n \equiv 1 \pmod{q}; \\ 0, & \text{otherwise,} \end{cases} \\ \sum_{a=1}^q \chi(a) &= \begin{cases} \phi(q), & \text{if } \chi = \chi_0; \\ 0, & \text{otherwise,} \end{cases} \end{aligned} \quad (2.9)$$

we have

$$\begin{aligned} & \sum_{d|q} d^{1/2} \sum_{s=1}^{q/d-1} \left| \sum_{\chi \neq \chi_0} \chi(sd+1) |L(1, \chi)| \right| \\ &= \sum_{d|q} d^{1/2} \sum_{s=1}^{q/d-1} \left| \sum_{\chi \neq \chi_0} \chi(sd+1) \left| \sum_{n \leq N} \frac{\chi(n)}{n} \right| \right| + O(\ln q) \\ &= \sum_{d|q} d^{1/2} \sum_{s=1}^{q/d-1} \left| \sum_{\chi} \chi(sd+1) \left| \sum_{n \leq N} \frac{\chi(n)}{n} \right| \right| + O(q \cdot d(q) \cdot \ln q) \\ &\ll \sum_{d|q} d^{1/2} \sum_{s=1}^{q/d-1} \left| \sum_{\chi} \chi(sd+1) \left(\left| \sum_{n \leq N} \frac{\chi(n)}{n} \right| - \left| \sum_{n \leq N} \frac{\chi(n)r(n)}{n} \right|^2 \right) \right| \\ &\quad + \sum_{d|q} d^{1/2} \sum_{s=1}^{q/d-1} \left| \sum_{\chi} \chi(sd+1) \left| \sum_{n \leq N} \frac{\chi(n)r(n)}{n} \right|^2 \right| + q \cdot d(q) \cdot \ln q \\ &\ll \sum_{d|q} d^{1/2} \cdot \frac{q}{\sqrt{d}} \cdot \left[\sum_{\chi} \left(\left| \sum_{n \leq N} \frac{\chi(n)}{n} \right| - \left| \sum_{n \leq N^2} \frac{\chi(n)r(n, N)}{n} \right| \right)^2 \right]^{1/2} \\ &\quad + \phi(q) \sum_{d|q} d^{1/2} \sum_{s=1}^{q/d-1} \sum'_{\substack{m \leq N \\ (sd+1)m \equiv n \pmod{q}}} \sum'_{\substack{n \leq N \\ m \neq n}} \frac{r(m)r(n)}{mn} + q \cdot d(q) \cdot \ln q \\ &\ll q \cdot d(q) \cdot \left[\sum_{\chi} \left| \sum_{N < n \leq N^2} \frac{\chi(n)r(n, N)}{n} \right|^2 \right]^{1/2} + q \cdot d(q) \cdot \ln q \\ &\quad + \phi(q) \sum_{d|q} d^{1/2} \sum'_{\substack{m \leq N \\ m \equiv n \pmod{d}}} \sum'_{\substack{n \leq N \\ m \neq n}} \frac{r(m)r(n)}{mn} \end{aligned}$$

$$\begin{aligned}
&\ll q^{3/2} \cdot d(q) \cdot \left(\sum'_{\substack{N < m \leq N^2 \\ m \equiv n \pmod{q}}} \sum'_{N < n \leq N^2} \frac{r(m, N) \cdot r(n, N)}{mn} \right)^{1/2} + q \cdot d(q) \cdot \ln^2 q \\
&\ll q \cdot d(q) \cdot \ln^2 q.
\end{aligned} \tag{2.10}$$

This proves [Lemma 2.1](#). \square

LEMMA 2.2. *For any integer $q \geq 3$, we have the asymptotic formula*

$$\sum_{\chi \neq \chi_0} |L(1, \chi)| = C \cdot \phi(q) + O(q^{1/2} \cdot \ln q), \tag{2.11}$$

where

$$C = \sum_{n=1}^{\infty} \frac{r^2(n)}{n^2} = \prod_p' \left[1 + \frac{\binom{2}{1}^2}{4^2 \cdot p^2} + \frac{\binom{4}{2}^2}{4^4 \cdot p^4} + \cdots + \frac{\binom{2n}{n}^2}{4^{2n} \cdot p^{2n}} + \cdots \right] \tag{2.12}$$

is an absolute constant.

PROOF. Let $N = q^{3/2}$. Then from (2.3), (2.6), the orthogonality relationship for character sums, and the method of proving [Lemma 2.1](#), we have

$$\begin{aligned}
\sum_{\chi \neq \chi_0} |L(1, \chi)| &= \sum_{\chi \neq \chi_0} \left| \sum_{n \leq N} \frac{\chi(n)}{n} \right| + O(\ln q) \\
&= \sum_{\chi} \left| \sum_{n \leq N} \frac{\chi(n) r(n)}{n} \right|^2 + O(\ln q) \\
&\quad + \sum_{\chi} \left(\left| \sum_{n \leq N} \frac{\chi(n)}{n} \right| - \left| \sum_{n \leq N} \frac{\chi(n) r(n)}{n} \right|^2 \right) \\
&= \phi(q) \cdot \sum'_{\substack{m \leq N \\ m \equiv n \pmod{q}}} \sum_{n \leq N} \frac{r(m) \cdot r(n)}{mn} + O(\ln q) \\
&\quad + O \left(\sum_{\chi} \left| \sum_{N < n \leq N^2} \frac{\chi(n) \cdot r(n, N)}{n} \right| \right) \\
&= \phi(q) \cdot \sum_{\substack{n=1 \\ (n, q)=1}}^{\infty} \frac{r^2(n)}{n^2} + O(\ln^2 q) \\
&\quad + O \left(q^{1/2} \left(\sum_{\chi} \left| \sum_{N < n \leq N^2} \frac{\chi(n) \cdot r(n, N)}{n} \right|^2 \right)^{1/2} \right) \\
&= \phi(q) \cdot \sum_{\substack{n=1 \\ (n, q)=1}}^{\infty} \frac{r^2(n)}{n^2} + O(q^{1/2} \cdot \ln q) \\
&= C \cdot \phi(q) + O(q^{1/2} \cdot \ln q).
\end{aligned} \tag{2.13}$$

This proves [Lemma 2.2](#). \square

LEMMA 2.3. *Let m, n , and q be integers with $q \geq 3$. Then we have the estimates*

$$S(m, n, q) = \sum_{a=1}^q' e\left(\frac{ma+n\bar{a}}{q}\right) \ll (m, n, q)^{1/2} q^{1/2} d(q), \quad (2.14)$$

where \sum_a' denotes the summation over all a such that $(a, q) = 1$.

PROOF. See [2]. □

3. Proof of Theorem 1.1. In this section, we will complete the proof of [Theorem 1.1](#). First we have

$$\begin{aligned} & \sum_{\chi \neq \chi_0} |L(1, \chi)| \cdot |S(m, n, \chi, q)|^2 \\ &= \sum_{r=1}^q' \sum_{s=1}^q' e\left(\frac{(r-s)m + (\bar{r} - \bar{s})n}{q}\right) \sum_{\chi \neq \chi_0} \chi(r\bar{s}) |L(1, \chi)| \\ &= \sum_{r=1}^q' \sum_{s=1}^q' e\left(\frac{r(1-\bar{s})m + \bar{r}(1-s)n}{q}\right) \sum_{\chi \neq \chi_0} \chi(s) |L(1, \chi)| \\ &= \phi(q) \sum_{\chi \neq \chi_0} |L(1, \chi)| + \sum_{s=2}^q' \sum_{r=1}^q' e\left(\frac{r(1-\bar{s})m + \bar{r}(1-s)n}{q}\right) \sum_{\chi \neq \chi_0} \chi(s) |L(1, \chi)|. \end{aligned} \quad (3.1)$$

From (3.1) and [Lemmas 2.2](#) and [2.3](#) we get

$$\begin{aligned} & \sum_{\chi \neq \chi_0} |L(1, \chi)| \cdot |S(m, n, \chi, q)|^2 \\ &= C \cdot \phi^2(q) + O(q^{3/2} \cdot \ln q) \\ &+ O\left(q^{1/2} d(q) \sum_{s=2}^q' ((s-1)m, (\bar{s}-1)n, q)^{1/2} \left| \sum_{\chi \neq \chi_0} \chi(s) |L(1, \chi)| \right| \right). \end{aligned} \quad (3.2)$$

Note that $(m, q) = (n, q) = (s, q) = 1$, and $(1-s, q) = (s\bar{s} - s, q) = (s(\bar{s}-1), q) = (\bar{s}-1, q)$, so we have $(s-1, \bar{s}-1, q) = (s-1, q)$. Thus from (3.2) and [Lemma 2.1](#) we obtain the asymptotic formula

$$\begin{aligned} & \sum_{\chi \neq \chi_0} |L(1, \chi)| \cdot |S(m, n, \chi, q)|^2 \\ &= C \cdot \phi^2(q) + O(q^{3/2} \cdot \ln q) \\ &+ O\left(q^{1/2} d(q) \sum_{s=2}^q' (s-1, q)^{1/2} \left| \sum_{\chi \neq \chi_0} \chi(s) |L(1, \chi)| \right| \right) \\ &= C \cdot \phi^2(q) + O(q^{3/2} \cdot \ln q) \\ &+ O\left(q^{1/2} d(q) \sum_{d|q} \sum_{t=1}^{q/d-1} d^{1/2} \left| \sum_{\chi \neq \chi_0} \chi(td+1) |L(1, \chi)| \right| \right) \\ &= C \cdot \phi^2(q) + O(q^{3/2} \cdot d^2(q) \cdot \ln^2 q). \end{aligned} \quad (3.3)$$

This completes the proof of the theorem.

ACKNOWLEDGMENT. This work is supported by the National Natural Science Foundation of China (NSFC) and the Shaanxi Province Natural Science Foundation of China (PNSF).

REFERENCES

- [1] S. Chowla, *On Kloosterman's sum*, Norske Vid. Selsk. Forh. (Trondheim) **40** (1967), 70–72.
- [2] T. Estermann, *On Kloosterman's sum*, Mathematika **8** (1961), 83–86.
- [3] A. V. Malyšev, *A generalization of Kloosterman sums and their estimates*, Vestnik Leningrad. Univ. **15** (1960), no. 13, 59–75 (Russian).

ZHANG WENPENG: RESEARCH CENTER FOR BASIC SCIENCE, XI'AN JIAOTONG UNIVERSITY, XI'AN, SHAANXI, CHINA

E-mail address: wpzhang@nwu.edu.cn

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.oter@usc.es