

## NOTES ON WHITEHEAD SPACE OF AN ALGEBRA

M. ARIAN-NEJAD

Received 2 August 2001

Let  $R$  be a ring, and denote by  $[R, R]$  the group generated additively by the additive commutators of  $R$ . When  $R_n = M_n(R)$  (the ring of  $n \times n$  matrices over  $R$ ), it is shown that  $[R_n, R_n]$  is the kernel of the regular trace function modulo  $[R, R]$ . Then considering  $R$  as a simple left Artinian  $F$ -central algebra which is algebraic over  $F$  with  $\text{Char } F = 0$ , it is shown that  $R$  can decompose over  $[R, R]$ , as  $R = Fx + [R, R]$ , for a fixed element  $x \in R$ . The space  $R/[R, R]$  over  $F$  is known as the Whitehead space of  $R$ . When  $R$  is a semisimple central  $F$ -algebra, the dimension of its Whitehead space reveals the number of simple components of  $R$ . More precisely, we show that when  $R$  is algebraic over  $F$  and  $\text{Char } F = 0$ , then the number of simple components of  $R$  is greater than or equal to  $\dim_F R/[R, R]$ , and when  $R$  is finite dimensional over  $F$  or is locally finite over  $F$  in the case of  $\text{Char } F = 0$ , then the number of simple components of  $R$  is equal to  $\dim_F R/[R, R]$ .

2000 Mathematics Subject Classification: 12E15, 16K40.

**1. Introduction.** Additive commutator elements of a ring  $R$  and the groups and structures they make have a great role in the general specification of a ring, and their study is one of the approaches to recognize rings in noncommutative ring theory [2, 3, 4, 5]. The reason is clear, they have covered the secrets of noncommutative behaviour of the structure. In recent years, these elements are returned once again under a full consideration, and a lot of wonderful works has been done on them [1, 10, 11, 12, 13]. Our study here is also among these studies, and it reveals some of bilateral relations between substructure given by additive commutators (the additive commutator group  $[R, R]$ , the additive Whitehead group, and the space  $R/[R, R]$ ) and some characteristics of the ring. In what follows let  $R$  be a ring. By  $[R, R]$  we denote the group generated additively by the additive commutators of  $R$ . Following [2], the additive group  $R/[R, R]$  is called the additive Whitehead group of  $R$ . This group is an  $F$ -vector space when  $R$  is a central  $F$ -algebra, and is called the Whitehead space of  $R$ .

**2. Results.** Our first result is about the additive commutator subgroup of a matrix ring over a given ring.

**PROPOSITION 2.1.** *Let  $R$  be a unitary ring and let  $R_n = M_n(R)$  be the ring of  $n \times n$  matrices over  $R$ . Consider the regular trace function on  $R_n$ , as  $\text{tr}: R_n \rightarrow R$ , then*

$$[R_n, R_n] = \{A \in R_n \mid \text{tr}(A) \in [R, R]\}. \quad (2.1)$$

**PROOF.** The inclusion “ $\subseteq$ ” follows by the fact that  $\text{tr}(AB - BA) \in [R, R]$ . In order to show the reverse inclusion, let  $\{E_{ij}\}$  be the matrix units and note that if  $i \neq j$ , we have  $E_{ij} = E_{ii}E_{ij} - E_{ij}E_{ii} \in [R_n, R_n]$  and  $E_{ii} - E_{jj} = E_{ij}E_{ji} - E_{ji}E_{ij} \in [R_n, R_n]$ . For any

$A = (a_{ij}) \in R_n$ , we have the following congruence:

$$A = \sum a_{ij}E_{ij} \equiv \sum a_{ii}E_{ii} \equiv \sum a_{ii}E_{11} \pmod{[R_n, R_n]}. \quad (2.2)$$

In particular, if  $\text{tr}(A) \in [R, R]$ , then  $A \in [R_n, R_n]$ .  $\square$

**COROLLARY 2.2.** *Consider the trace function on  $R_n$  module of  $[R, R]$ . Clearly the group isomorphism  $R_n/[R_n, R_n] \cong R/[R, R]$  can be derived.*

**THEOREM 2.3.** *Let  $R$  be a left Artinian central simple  $F$ -algebra which is algebraic over  $F$  with  $\text{Char}F = 0$ . Then  $R$  decomposes over  $[R, R]$  as  $R = Rx + [R, R]$ , for a fixed  $x \in R$ .*

**PROOF.** By Wedderburn-Artin theorem,  $R = M_n(D)$  for a division ring  $D$  and suitable  $n \in \mathbb{N}$  [6, 14]. We divide our proof into two parts.

(i) Let  $n = 1$ , in other words let  $R = M_1(D) = D$  be a division ring. Let  $a \in R$  and let  $f(t) = t^r + b_1t^{r-1} + \cdots + b_r$  be the minimal polynomial of  $a$  over  $F$ , where  $b_i \in F$ ,  $i = 1, 2, \dots, r$  and  $r = \dim_F F(a)$ . By the Wedderburn theorem [9, page 265],  $f(t)$  splits completely in  $R[t]$ , this means that there exists  $c_i \in R^* = D - \{0\}$ ,  $i = 1, 2, \dots, r-1$ , such that  $f(t) = (t-a)(t-c_1ac_1^{-1}) \cdots (t-c_{r-1}ac_{r-1}^{-1})$ . Then we have

$$\begin{aligned} \text{Tr}_{F(a)/F}(a) &= a + c_1ac_1^{-1} + c_2ac_2^{-1} + \cdots + c_{r-1}ac_{r-1}^{-1} \\ &= ra + (c_1ac_1^{-1} - a) + \cdots + (c_{r-1}ac_{r-1}^{-1} - a) \\ &= ra + (c_1(ac_1^{-1}) - (ac_1^{-1})c_1) + \cdots + (c_{r-1}(ac_{r-1}^{-1}) - (ac_{r-1}^{-1})c_{r-1}) \\ &= ra + d_1 + d_2 + \cdots + d_{r-1} = ra + d, \end{aligned} \quad (2.3)$$

where  $d_1, \dots, d_{r-1}, d \in [R, R]$ . This simply yields  $a \in F + [R, R]$  which imply that  $R = F + [R, R]$ ,  $x = 1$ .

(ii) Let  $n \in \mathbb{N}$  be an arbitrary positive integer. We have  $R = M_n(D)$ , where  $D$  is a division ring. By (i),  $D = F + [D, D]$ , so

$$R = M_n(D) = M_n(F + [D, D]) = M_n(F) + M_n([D, D]) \subseteq M_n(F) + [R, R] \subseteq R. \quad (2.4)$$

This implies that  $R = M_n(F) + [R, R]$ . By this formula, given  $A \in R$ , there exist  $B \in M_n(F)$  and  $C \in [R, R]$  such that  $A = B + C$ , hence  $A = (B - (\text{tr}B/n)I) + (\text{tr}B/n)I + C$ , where  $I$  is the identity matrix of size  $n$ . By [Proposition 2.1](#),  $(B - (\text{tr}B/n)I) \in [R, R]$ , and  $A = (\text{tr}B/n)I + ((B - (\text{tr}B/n)I) + C)$ , consequently

$$R = FI + [R, R], \quad x = I. \quad (2.5)$$

$\square$

To see a different statements and initial ideas of these theorems we refer the reader to [1, 2]. Also a multiplicative version of [Theorem 2.3](#) could be found in [11].

Now, we are going to state our main result, which is about the Whitehead space of a semisimple ring. This theorem is a generalization of a nice theorem due to R. Brauer [8, page 130].

**THEOREM 2.4.** *Let  $R$  be a left Artinian semisimple central  $F$ -algebra and let  $k$  be the number of left simple components of  $R$ . Then,*

- (i) *if  $R$  is algebraic over  $F$  and  $\text{Char } F = 0$ , then  $k \geq \dim_F R/[R, R]$ ;*
- (ii) *if  $R$  is finite dimensional over  $F$ , or is locally finite over  $F$ , and  $\text{Char } F = 0$ , then  $k = \dim_F R/[R, R]$ .*

**PROOF.** Consider the following chain of functions:

$$R \xrightarrow{f_1} M_{n_1}(D_1) \times \cdots \times M_{n_k}(D_k) \xrightarrow{f_2} D_1/[D_1, D_1] \times \cdots \times D_k/[D_k, D_k], \quad (2.6)$$

where  $f_1$  is the isomorphism given by the Wedderburn-Artin theorem for the decomposition of a semisimple left Artinian ring into a direct product of simple ring [6, 14], and  $f_2$  is the  $F$ -algebra homomorphisms, by considering component-wise the trace function on  $M_{n_i}(D_i) \text{ mod } [D_i, D_i]$ ,  $i = 1, \dots, k$ .

By [Proposition 2.1](#) we have,  $\ker(f_2 \circ f_1) = [R, R]$ , noting that  $[R, R] \cong [R_1, R_1] \times \cdots \times [R_k, R_k]$ , where  $R_{n_i} = M_{n_i}(D_i)$ ,  $i = 1, \dots, k$ . Therefore the following  $F$ -isomorphism holds:

$$R/[R, R] \cong D_1/[D_1, D_1] \times \cdots \times D_k/[D_k, D_k]. \quad (2.7)$$

It remains to compute the dimension of Whitehead space of a division ring in the two cases (i) and (ii) above.

First let  $D$  be algebraic over  $F$  and  $\text{Char } F = 0$ . We show that any two elements  $\bar{a}, \bar{b} \in D/[D, D]$  are linearly dependent. By [Theorem 2.3](#), there exist elements  $\alpha, \beta \in F$  and  $d_1, d_2 \in [D, D]$ , such that  $a = \alpha + d_1$  and  $b = \beta + d_2$ . In other words,  $\beta\bar{a} - \alpha\bar{b} = \bar{0}$  in  $D/[D, D]$ . Hence in this case  $\dim_F D/[D, D] \leq 1$ .

Now let  $D$  be finite dimensional  $F$ -central algebra. Let  $\text{RT}_{D/F} : D \rightarrow F$  be the reduced trace function which is surjective by [7, page 148]. Furthermore, by a theorem of Amitsur and Rowen [5, page 171] its kernel is equal to  $[D, D]$  and so it is a hyperplane over  $F$ , in this case  $\dim_F D/[D, D] = 1$ .

As a latter case let  $D$  be a locally finite division ring over its center  $F$  and  $\text{Char } F = 0$ . Now consider the function  $\text{TR} : D \rightarrow F$  defined by

$$\text{TR}(x) = \frac{1}{\deg_F(x)} \text{Tr}_{F(x)/F}(x), \quad (2.8)$$

we show that this function is an  $F$ -linear surjective map, whose kernel is  $[D, D]$ . The claim then is clear.

First note that in this case  $1 \notin [D, D]$ , for if  $1 \in [D, D]$ , then there exist some  $x_i$ 's and  $y_i$ 's in  $D$ , such that  $1 = \sum(x_i y_i - y_i x_i)$ . Let  $D_1$  be the division ring generated by  $F$  together with  $x_i$ 's and  $y_i$ 's. Taking the reduced trace of  $D_1$  over its centre of both sides of  $1 = \sum(x_i y_i - y_i x_i)$ , we get a contradicting result. Therefore  $[D, D] \cap F = \{0\}$ . Now, by considering the trace formula (given in the proof of [Theorem 2.3](#)) for elements  $a$ ,  $b$  and  $\lambda a + b$  ( $\lambda \in F$ ) in  $D$ , it is readily verified that

$$\frac{1}{r} \text{Tr}(\lambda a + b) = \frac{\lambda}{n} \text{Tr}(a) + \frac{1}{m} \text{Tr}(b), \quad (2.9)$$

where  $r$ ,  $n$ , and  $m$  are degrees of  $\lambda a + b$ ,  $a$  and  $b$ . So  $\text{TR}$  is  $F$ -linear. The surjectivity is clear. In order to specify the kernel of  $\text{TR}$ , consider the trace formula for elements of  $[D, D]$ . Suppose that  $a \in [D, D]$ . Now, we have  $\text{Tr}_{F(a)/F}(a) = na + d \in [D, D] \cap F$ , where  $n$  is the degree of  $a$  over  $F$  and  $d \in [D, D]$ . Therefore  $\text{TR}(a) = 0$ . By the same argument we can see that if  $\text{TR}(a) = 0$ , then  $a \in [D, D]$ .  $\square$

**ACKNOWLEDGMENT.** The author is indebted in part to the Research Council of the University of Zanjan for support.

#### REFERENCES

- [1] S. Akbari and M. Arian-Nejad, *Left Artinian algebraic algebras*, Algebra Colloq. **8** (2001), no. 4, 463–470.
- [2] S. Akbari, M. Arian-Nejad, and M. L. Mehrabadi, *On additive commutator groups in division rings*, Results Math. **33** (1998), no. 1-2, 9–21.
- [3] S. Akbari and M. Mahdavi-Hezavehi, *Normal subgroups of  $\text{GL}_n(D)$  are not finitely generated*, Proc. Amer. Math. Soc. **128** (2000), no. 6, 1627–1632.
- [4] S. Akbari, M. Mahdavi-Hezavehi, and M. G. Mahmudi, *Maximal subgroups of  $\text{GL}_1(D)$* , J. Algebra **217** (1999), no. 2, 422–433.
- [5] S. A. Amitsur and L. H. Rowen, *Elements of reduced trace 0*, Israel J. Math. **87** (1994), no. 1-3, 161–179.
- [6] P. M. Cohn, *Algebra. Vol. 3*, 2nd ed., John Wiley & Sons, Chichester, 1991.
- [7] P. K. Draxl, *Skew Fields*, London Mathematical Society Lecture Note Series, vol. 81, Cambridge University Press, Cambridge, 1983.
- [8] I. N. Herstein, *Noncommutative Rings*, The Carus Mathematical Monographs, vol. 15, The Mathematical Association of America, Washington, DC, 1971.
- [9] T. Y. Lam, *A First Course in Noncommutative Rings*, Graduate Texts in Mathematics, vol. 131, Springer-Verlag, New York, 1991.
- [10] M. Mahdavi-Hezavehi, *Extension of valuations on derived groups of division rings*, Comm. Algebra **23** (1995), no. 3, 913–926.
- [11] ———, *Commutator subgroups of finite dimensional division algebras*, Rev. Roumaine Math. Pures Appl. **43** (1998), no. 9-10, 853–867.
- [12] M. Mahdavi-Hezavehi and S. Akbari, *Some special subgroups of  $\text{GL}_n(D)$* , Algebra Colloq. **5** (1998), no. 4, 361–370.
- [13] M. Mahdavi-Hezavehi, S. Akbari-Feyzaabadi, M. Mehraabaadi, and H. Hajie-Abolhassan, *On derived groups of division rings. II*, Comm. Algebra **23** (1995), no. 8, 2881–2887.
- [14] L. H. Rowen, *Ring Theory*, Academic Press, Massachusetts, 1991, Student edition.

M. ARIAN-NEJAD: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZANJAN, ZANJAN, IRAN  
*E-mail address:* [arian@mail.znu.ac.ir](mailto:arian@mail.znu.ac.ir)

## Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

### Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from "Qualitative Theory of Differential Equations," allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

### Guest Editors

**José Roberto Castilho Piqueira**, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; [piqueira@lac.usp.br](mailto:piqueira@lac.usp.br)

**Elbert E. Neher Macau**, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; [elbert@lac.inpe.br](mailto:elbert@lac.inpe.br)

**Celso Grebogi**, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; [grebogi@abdn.ac.uk](mailto:grebogi@abdn.ac.uk)