

RESEARCH NOTES

CONTINUITY OF MULTIPLICATION OF DISTRIBUTORS

JAN KUCERA and KELLY MCKENNON

Department of Pure and Applied Mathematics
Washington State University
Pullman, Washington 99164 U.S.A.

(Received February 20, 1981)

ABSTRACT. In a reference book for distributions [1], it is shown that the multiplication $(u, f) \mapsto uf$ on $C^\infty \times \mathcal{D}'$, as well as on $\mathcal{C}_M \times \mathcal{S}'$, is hypocontinuous. We show here that in both cases it is discontinuous.

KEY WORDS AND PHRASES. Distribution, temperate distribution, dual space, strong topology, inductive limit.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. Primary 46F10, Secondary 46A09.

1. INTRODUCTION.

The discontinuity of multiplication on $C^\infty \times \mathcal{D}'$, seems to be part of general folklore, but no proof has yet been published. As far as we know the second result is new. The presented proofs are simple enough that they can be included in any future textbook on distributions.

THEOREM 1. Let β be the strong and σ the weak* topology, both on $\mathcal{D}'(\mathbb{R}^n)$, and γ the usual topology on $C^\infty(\mathbb{R}^n)$. Then the multiplication $(u, f) \mapsto uf : C_\gamma^\infty \times \mathcal{D}'_\beta \rightarrow \mathcal{D}'_\sigma$ is not jointly continuous.

PROOF. The family \mathfrak{F} of all increasing sequences of positive integers is a directed set under the induced product ordering from \mathbb{N}^N . Let Γ be the directed product $\mathbb{N} \times \mathfrak{F}$, and let $d = \partial^n / \partial x_1 \partial x_2 \dots \partial x_n$. For each $(m, s) \in \Gamma$, put $f_{m,s}(t) = (mt^{m+1})^{-1}$ if $t \in [1, \exp(m \cdot s(m))]$, with $f_{m,s}(t) = 0$ otherwise, and $F_{m,s} : \mathcal{D} \rightarrow C : g \mapsto s(m)^{-1/2} d^m g(0)$.

If $g_{m,s}$ is the Fourier transform of $\prod_{j=1}^n f_{m,s}(x_j)$, then the inequalities

$$\int_{-\infty}^{\infty} t^k f_{m,s}(t) dt \leq m^{-1}, \quad k = 1, 2, \dots, m-1, \quad \int_{-\infty}^{\infty} t^m f_{m,s}(t) dt = s(m) \text{ imply,}$$

respectively, $\|d^k g_{m,s}\|_{\infty} \leq m^{-n}$ and $d^m g_{m,s}(0) = i^{nm} s(m)^n$, where i is the imaginary unit. Thus $\lim_{(m,s) \in \Gamma} g_{m,s} = 0$ in C_{γ}^{∞} . For any $\psi \in \mathcal{D}(R^n)$, which equals 1 in some neighborhood of the origin, $\lim_{(m,s) \in \Gamma} |(g_{m,s} F_{m,s}) \psi| = \lim_{(m,s) \in \Gamma} |s(m)^{-\frac{1}{2}} d_m g_{m,s}(0)| = \lim_{(m,s) \in \Gamma} s(m)^{n-\frac{1}{2}} = +\infty$ and $\{g_{m,s} F_{m,s}\}_{(m,s) \in \Gamma}$ does not converge to 0 in \mathcal{D}'_0 .

It remains to show that $\{F_{m,s}\}$ converges to 0 uniformly on every set \mathbb{B} bounded in \mathbb{R} . For each such \mathbb{B} , there exists $r \in \mathbb{R}$ such that $|d^m g(0)| \leq r(m)$ for all $m \in \mathbb{N}$ and $g \in \mathbb{B}$. Choose $\varepsilon > 0$ and $s \in \mathbb{R}$ such that $s(m) > \varepsilon^{-2} r^2(m)$ for all $m \in \mathbb{N}$. Then

$$|F_{m,s}(g)| = |s(m)^{-\frac{1}{2}} d^m g(0)| \leq s(m)^{-\frac{1}{2}} r(m) < \varepsilon \quad \text{for all } g \in \mathbb{B}.$$

In the sequel, we need a weight function $W(x) = (1 + |x|^2)^{\frac{1}{2}}$, and Hilbert spaces $H_k = \{f : R^n \rightarrow C; \|f\|_k^2 = \sum_{|\alpha| \leq k} \int_{R^n} |W^{k-|\alpha|} D^\alpha f|^2 dx < +\infty\}$, $k \in \mathbb{N}$. The

space \mathbb{G} of rapidly decreasing functions equals the proj lim H_k . For every $p, q \in \mathbb{N}$, the space $\mathbb{G}_{p,q} = \{u : R^n \rightarrow C; f \mapsto uf : H_p \rightarrow H_q \text{ continuous}\}$ equipped with the operator norm $\|\cdot\|_{p,q}$ is Banach. If $\mathbb{G}_q = \text{ind lim}_{p \rightarrow \infty} \mathbb{G}_{p,q}$, then the

space \mathbb{G}_M of rapidly increasing functions equals proj lim \mathbb{G}_q , [4]. Finally, denote by $\|\cdot\|_{\infty}$ the supremum norm of $L^{\infty}(R^n)$ and by d_{ε} the dilation operator $(d_{\varepsilon} f)(x) = f(\varepsilon x)$.

LEMMA 1. For each $k \in \mathbb{N}$ and multi-index $\alpha \in \mathbb{N}^n$,

$$\lim_{\varepsilon \rightarrow 0+} \|\varepsilon^{\alpha} |W^k(x) D^\alpha \exp(-|x|^2)|\|_{\infty} = \|D^\alpha \exp(-|x|^2)\|_{\infty}, \text{ where } |x|^2 = x_1^2 + x_2^2 + \dots + x_n^2.$$

PROOF. $\lim_{\varepsilon \rightarrow 0+} \|\varepsilon^{\alpha} |W^k(x) D^\alpha \exp(-\frac{|x|^2}{\varepsilon})|\|_{\infty} = \lim_{\varepsilon \rightarrow 0+} \|\varepsilon^{\alpha} |W^k(\varepsilon x) D^\alpha \exp(-|\varepsilon x|^2)|\|_{\infty} = \|\varepsilon^{\alpha} |W^k(\varepsilon x) D^\alpha \exp(-|\varepsilon x|^2)|\|_{\infty} = \|\varepsilon^{\alpha} |W^k(x) D^\alpha \exp(-|x|^2)|\|_{\infty} = \|D^\alpha \exp(-|x|^2)\|_{\infty}.$

LEMMA 2. If $p, q \in \mathbb{N}$, $0 \leq q \leq p$, and $r = 1 + [\frac{1}{2}n]$, then there is a sequence $\{f_m\}$ in $\mathcal{S}(\mathbb{R}^n)$ such that $\sup_m \|f_m\|_{p,q} \leq 1$ and

$$\lim_{m \rightarrow \infty} \|f_m(x) \exp(-|x|^2)\|_{q+r} = \infty.$$

PROOF. By Prop. 8 of [4] and Lemma 1, there exists $A > 0$ such that

$$\limsup_{\varepsilon \rightarrow 0^+} \|\varepsilon^q \exp(-\frac{|x|}{\varepsilon}^2)\|_{p,q} \leq \limsup_{\varepsilon \rightarrow 0^+} \varepsilon^q A \sum_{|\alpha| \leq q} \|w^{q-p|\alpha|}(x) D^\alpha \exp(-\frac{|x|}{\varepsilon}^2)\|_\infty =$$

$$A \sum_{|\alpha|=q} \|D^\alpha \exp(-|x|^2)\|_\infty.$$

Define $h_m(x) = m^{-q} \exp(-|mx|^2)$, $m \in \mathbb{N}$, $x \in \mathbb{R}^n$.

Then $S = \sup_m \|h_m\|_{p,q} < +\infty$. If we put $f_m = S^{-1}h_m$, $m \in \mathbb{N}$, then

$$\begin{aligned} \sup_m \|f_m\|_{p,q} &\leq 1 \text{ and } \|f_m(x) \exp(-|x|^2)\|_{q+r}^2 = \\ S^{-2} m^{-2q} \sum_{|\alpha| \leq q+r} \int_{\mathbb{R}^n} &|w^{q+r-|\alpha|}(x) D^\alpha \exp(-(1+m^2)|x|^2)|^2 dx = \\ S^{-2} m^{-2q} (1+m^2)^{-\frac{1}{2}n} \sum_{|\alpha| \leq q+r} &(1+m^2)^{|\alpha|} \int_{\mathbb{R}^n} |w^{q+r-|\alpha|}((1+m^2)^{-\frac{1}{2}}x) D^\alpha \exp(-|x|^2)|^2 dx. \end{aligned}$$

Take a multi-index β such that $|\beta| = q+r$. Then

$$\begin{aligned} \limsup_{m \rightarrow \infty} &\|f_m(x) \exp(-|x|^2)\|_{q+r}^2 \geq \\ \limsup_{m \rightarrow \infty} &S^{-2} m^{-2q} (1+m^2)^{-\frac{1}{2}n+q+r} \int_{\mathbb{R}^n} |D^\beta \exp(-|x|^2)|^2 dx = \\ S^{-2} &\int_{\mathbb{R}^n} |D^\beta \exp(-|x|^2)|^2 dx \cdot \limsup_{m \rightarrow \infty} m^{-2q} (1+m^2)^{q+r-\frac{1}{2}n} = +\infty. \end{aligned}$$

THEOREM 2. Let β be the strong and σ the weak* topology on $\mathcal{S}'(\mathbb{R}^n)$. Then the multiplication $(u, f) \mapsto (u, f) : \mathcal{G}_M \times \mathcal{S}'_\beta \rightarrow \mathcal{S}'_\sigma$ is not jointly continuous.

PROOF. The polar P of the singleton $\{\exp(-|x|^2)\} \subset \mathcal{S}$ is a σ -neighborhood of 0 in \mathcal{S}' . If the multiplication was continuous, there would be neighborhoods of 0, $U \subset \mathcal{G}_M$ and $V \subset \mathcal{S}'_\beta$, such that $UV \subset P$. For some $q \in \mathbb{N}$, there exists a

a neighborhood G of 0 in \mathfrak{G}_q such that $G \cap \mathfrak{G}_M \subset U$, and there exists a ball $B(\varepsilon)$ of radius ε about the origin in $\mathfrak{G}_{q,p}$ such that $B(\varepsilon) \subset G$. Since $\mathfrak{g} \subset \mathfrak{G}_M$, Lemma 2 implies existence of a sequence $\{f_m\}$ in $B(\varepsilon) \cap \mathfrak{G}_M$ such that

$$\lim_{m \rightarrow \infty} \|f_m(x) \exp(-|x|^2)\|_{q+r} = +\infty, \text{ where } r = 1 + [\frac{1}{2}n].$$

For any $g \in V$, we have $f_m g \in UV \subset P$, which implies

$|g(f_m \exp(-|x|^2))| = |(f_m g) \exp(-|x|^2)| \leq 1$. Hence $f_m \exp(-|x|^2)$ is contained in the polar V^0 of V . Since V^0 is bounded in \mathfrak{g} , the sequence $\{f_m \exp(-|x|^2)\}$ is bounded in \mathfrak{g} , too; i.e., $\sup_m \|f_m \exp(-|x|^2)\|_{q+r} < +\infty$, which is a contradiction.

REFERENCES

1. SCHWARTZ, L., Théorie des distributions, Nouvelle édition, Hermann, Paris, (1966).
2. SCHAEFER, H. H., Topological Vector Spaces, 3rd printing, Springer-Verlag, New York, Heidelberg, Berlin (1971).
3. RUDIN, W., Functional Analysis, McGraw-Hill, New York (1973).
4. KUČERA, J., On Multipliers of Temperate Distributions, Czech. Math. J. 21(96) (1971), pp. 610-618.
5. TRÈVES, F., Topological Vector Spaces, Distributions and Kernels, Academic Press, New York (1967).

Special Issue on Time-Dependent Billiards

Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	March 1, 2009
First Round of Reviews	June 1, 2009
Publication Date	September 1, 2009

Guest Editors

Edson Denis Leonel, Department of Statistics, Applied Mathematics and Computing, Institute of Geosciences and Exact Sciences, State University of São Paulo at Rio Claro, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil; edleonel@rc.unesp.br

Alexander Loskutov, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; loskutov@chaos.phys.msu.ru