

ON THE SPECTRUM OF WEAKLY ALMOST PERIODIC SOLUTIONS OF CERTAIN ABSTRACT DIFFERENTIAL EQUATIONS

ARIBINDI SATYANARAYAN RAO

Department of Mathematics
Sir George Williams Campus
Concordia University
Montreal, Quebec, Canada

and

L.S. DUBE

Department of Mathematics
Vanier College
821 Ste-Mis Croix Blvd.
St.-Laurent
Quebec, H4L 3x9, Canada

(Received October 15, 1984)

ABSTRACT. In a sequentially weakly complete Banach space, if the dual operator of a linear operator A satisfies certain conditions, then the spectrum of any weakly almost periodic solution of the differential equation $u' = Au + f$ is identical with the spectrum of f except at the origin, where f is a weakly almost periodic function.

KEY WORDS AND PHRASES. *Strongly (weakly) almost periodic function, sequentially weakly complete Banach space, densely defined linear operator, dual operator, Hilbert space, nonnegative self-adjoint operator.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 34C25, 34G05; 43A60

1. INTRODUCTION.

Suppose X is a Banach space and X^* is the dual space of X . Let J be the interval $-\infty < t < \infty$. A continuous function $f : J \rightarrow X$ is said to be strongly almost periodic if, given $\epsilon > 0$, there is a positive real number $\lambda = \lambda(\epsilon)$ such that any interval of the real line of length λ contains at least one point τ for which

$$\sup_{t \in J} ||f(t+\tau) - f(t)|| \leq \epsilon. \quad (1.1)$$

We say that a function $f : J \rightarrow X$ is weakly almost periodic if the scalar-valued function $\langle x^*, f(t) \rangle = x^* f(t)$ is almost periodic for each $x^* \in X^*$.

It is known that, if X is sequentially weakly complete, $f : J \rightarrow X$ is weakly almost periodic, and λ is a real number, then the weak limit

$$m(e^{-i\lambda t}f(t)) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T e^{-i\lambda t}f(t)dt \quad (1.2)$$

exists in X and is different from the null element 0 of X for at most a countable set $\{\lambda_n\}_{n=1}^\infty$, called the spectrum of $f(t)$ (see Theorem 6, p. 43, Amerio-Prouse [1]). We denote by $\sigma(f(t))$ the spectrum of $f(t)$.

2. RESULTS

Our first result is as follows (see Theorem 9, p. 79, Amerio-Prouse [1] for the spectrum of an S^1 -almost periodic function).

THEOREM 1. Suppose X is a sequentially weakly complete Banach space, A is a densely defined linear operator with domain $D(A)$ and range $R(A)$ in X , and the dual operator A^* is densely defined in X^* , with $R(i\lambda - A^*)$ being dense in X^* for all real $\lambda \neq 0$. Further, suppose $f: J \rightarrow X$ is a weakly almost periodic (or an S^1 -almost periodic continuous) function. If a differentiable function $u: J \rightarrow D(A)$ is a weakly almost periodic solution of the differential equation

$$u'(t) = Au(t) + f(t) \quad (1.3)$$

on J , with u' being weakly continuous on J , then $\sigma(u(t)) \setminus \{0\} = \sigma(f(t)) \setminus \{0\}$.

PROOF OF THEOREM 1. First we note that u is bounded on J , since u is weakly almost periodic. Hence, for $x^* \in X^*$, we have

$$\begin{aligned} \frac{1}{T} \int_0^T e^{-i\lambda t} x^* u'(t) dt &= x^* \frac{1}{T} \{ [e^{-i\lambda t} u(t)]_0^T + \frac{i\lambda}{T} \int_0^T e^{-i\lambda t} u(t) dt \} \\ &\rightarrow i\lambda x^* m(e^{-i\lambda t} u(t)) \text{ as } T \rightarrow \infty. \end{aligned} \quad (2.1)$$

So, for $x^* \in D(A^*)$, it follows from (1.3) that

$$\begin{aligned} \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T e^{-i\lambda t} x^* Au(t) dt &= \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T e^{-i\lambda t} (A^* x^*) u(t) dt \\ &= \lim_{T \rightarrow \infty} (A^* x^*) \left[\frac{1}{T} \int_0^T e^{-i\lambda t} u(t) dt \right] \\ &= (A^* x^*) m(e^{-i\lambda t} u(t)) \\ &= i\lambda x^* m(e^{-i\lambda t} u(t)) - x^* m(e^{-i\lambda t} f(t)). \end{aligned} \quad (2.2)$$

Consequently, we have

$$x^* m(e^{-i\lambda t} f(t)) = (i\lambda x^* - A^* x^*) m(e^{-i\lambda t} u(t)). \quad (2.3)$$

Now suppose that $\lambda \in \sigma(f(t)) \setminus \{0\}$. Then, since $D(A^*)$ is dense in X^* , there exists $x_1^* \in D(A^*)$ such that

$$0 \neq x_1^* m(e^{-i\lambda t} f(t)) = (-\lambda x_1^* - A^* x_1^*) m(e^{-i\lambda t} u(t)). \quad (2.4)$$

Therefore $m(e^{-i\lambda t}u(t)) \neq 0$ and so $\lambda \in \sigma(u(t)) \setminus \{0\}$.

Thus we have

$$\sigma(f(t)) \setminus \{0\} \subset \sigma(u(t)) \setminus \{0\}. \quad (2.5)$$

Now assume that $\lambda \in \sigma(u(t)) \setminus \{0\}$. Then, since $R(i\lambda - A^*)$ is dense in X^* , there exists $x_2^* \in D(A^*)$ such that

$$0 \neq (-i\lambda x_2^* - A^* x_2^*)m(e^{-i\lambda t}u(t)) = x_2^* m(e^{-i\lambda t}f(t)). \quad (2.6)$$

Therefore $m(e^{-i\lambda t}f(t)) \neq 0$ and so $\lambda \in \sigma(f(t)) \setminus \{0\}$.

Consequently, we have

$$\sigma(u(t)) \setminus \{0\} \subset \sigma(f(t)) \setminus \{0\}. \quad (2.7)$$

It follows from (2.5) and (2.7) that $\sigma(u(t)) \setminus \{0\} = \sigma(f(t)) \setminus \{0\}$, which completes the proof of the theorem.

REMARK 1. The conclusion of Theorem 1 remains valid if $D(A^*)$ is total and $R(i\lambda - A^*)$ is total for all real $\lambda \neq 0$, instead of dense in X^* .

We indicate the proof of the following result.

THEOREM 2. In a sequentially weakly complete Banach space X , suppose A is a densely defined linear operator, the dual operator A^* is densely defined in X^* , with $R(\lambda^2 + A^*)$ being dense in X^* for all real $\lambda \neq 0$, and $f : J \rightarrow X$ is a weakly almost periodic (or an S^1 -almost periodic continuous) function. If a twice differentiable function $u : J \rightarrow D(A)$ is a weakly almost periodic solution of the differential equation

$$u''(t) = Au(t) + f(t) \quad (3.1)$$

on J , with u'' being weakly continuous and u' bounded on J , then

$$\sigma(u(t)) \setminus \{0\} = \sigma(f(t)) \setminus \{0\}.$$

PROOF. For $x^* \in D(A^*)$, we have

$$\begin{aligned} \frac{1}{T} \int_0^T e^{-i\lambda t} x^* u''(t) dt &= x^* \left\{ \frac{1}{T} [e^{-i\lambda t} u'(t)]_0^T + \frac{i\lambda}{T} \int_0^T e^{-i\lambda t} u'(t) dt \right\} \\ &= x^* \left\{ \frac{1}{T} [e^{-i\lambda t} u'(t)]_0^T + \frac{i\lambda}{T} [e^{-i\lambda t} u(t)]_0^T - \frac{\lambda^2}{T} \int_0^T e^{-i\lambda t} u(t) dt \right\} \\ &\rightarrow -\lambda^2 x^* m(e^{-i\lambda t} u(t)) \quad \text{as } T \rightarrow \infty. \end{aligned} \quad (3.2)$$

Hence it follows from (3.1) that

$$\begin{aligned} \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T e^{-i\lambda t} x^* Au(t) dt &= (A^* x^*) m(e^{-i\lambda t} u(t)) \\ &= -\lambda^2 x^* m(e^{-i\lambda t} u(t)) - x^* m(e^{-i\lambda t} f(t)). \end{aligned} \quad (3.3)$$

Thus we have

$$-x^* m(e^{-i\lambda t} f(t)) = (\lambda^2 x^* + A^* x^*) m(e^{-i\lambda t} u(t)). \quad (3.4)$$

Now the rest of the proof parallels that of Theorem 1.

REMARK 2. The conclusion of Theorem 2 also remains valid if $D(A^*)$ is total and $R(\lambda^2 + A^*)$ is total for all real $\lambda \neq 0$, instead of dense in X^* .

REMARK 3. If X is a Hilbert space and A is a nonnegative self-adjoint operator, then the hypotheses on A in Theorem 2 are verified (see Corollary 2, p. 208, Yosida [2]) and so Theorem 2 is a generalization of a result of Zaidman [3].

NOTE. As a consequence of our Theorem 1, we have the following result:

THEOREM 3. In a Hilbert space H , suppose A is a self-adjoint operator and $f : J \rightarrow H$ is a weakly almost periodic (or an S^1 -almost periodic continuous) function. If a differentiable function $u : J \rightarrow D(A)$ is a weakly almost periodic solution of the differential equation

$$u'(t) = Au(t) + f(t)$$

on J , with u' being weakly continuous on J , then

$$\sigma(u(t)) \setminus \{0\} = \sigma(f(t)) \setminus \{0\}.$$

PROOF. By Example 4, p. 210, Yosida [2], $R(i\lambda - A) = H$ for all real $\lambda \neq 0$.

ACKNOWLEDGEMENT. This work was supported by the National Research Council of Canada Grant Nos. 4056 and A-9085.

REFERENCES

1. AMERIO, L. and PROUSE, G. Almost Periodic Functions and Functional Equations, Van Nostrand Reinhold Company (1971).
2. YOSIDA, K. Functional Analysis, Springer-Verlag New York Inc. (1971).
3. ZAIDMAN, S. Spectrum of Almost Periodic Solutions for some abstract Differential Equations, Math. Anal. Appl., 28 (1969), pp. 336-338

Special Issue on Boundary Value Problems on Time Scales

Call for Papers

The study of dynamic equations on a time scale goes back to its founder Stefan Hilger (1988), and is a new area of still fairly theoretical exploration in mathematics. Motivating the subject is the notion that dynamic equations on time scales can build bridges between continuous and discrete mathematics; moreover, it often reveals the reasons for the discrepancies between two theories.

In recent years, the study of dynamic equations has led to several important applications, for example, in the study of insect population models, neural network, heat transfer, and epidemic models. This special issue will contain new researches and survey articles on Boundary Value Problems on Time Scales. In particular, it will focus on the following topics:

- Existence, uniqueness, and multiplicity of solutions
- Comparison principles
- Variational methods
- Mathematical models
- Biological and medical applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Authors should follow the Advances in Difference Equations manuscript format described at the journal site <http://www.hindawi.com/journals/ade/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	April 1, 2009
First Round of Reviews	July 1, 2009
Publication Date	October 1, 2009

Lead Guest Editor

Alberto Cabada, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; alberto.cabada@usc.es

Guest Editor

Victoria Otero-Espinar, Departamento de Análise Matemática, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; mvictoria.oter@usc.es