

RINGS DECOMPOSED INTO DIRECT SUMS OF J-RINGS AND NIL RINGS

HISAO TOMINAGA

Department of Mathematics
Okayama University
Okayama 700, Japan

(Received April 25, 1984)

ABSTRACT. Let R be a ring (not necessarily with identity) and let E denote the set of idempotents of R . We prove that R is a direct sum of a J -ring (every element is a power of itself) and a nil ring if and only if R is strongly π -regular and E is contained in some J -ideal of R . As a direct consequence of this result, the main theorem of [1] follows.

KEY WORDS AND PHRASES. *Periodic, potent, J -ring, nil ring, strongly π -regular ring, direct sum.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. Primary 16A15; Secondary 16A70.

1. INTRODUCTION.

Throughout the present note, R will represent a ring (not necessarily with identity), N the set of nilpotent elements of R , and E the set of idempotents of R . We say that R is periodic if for each $r \in R$, there exist distinct positive integers h, k for which $r^h = r^k$. According to Chacron's theorem (see, e.g., [2, Theorem 1]), R is periodic if and only if for each $r \in R$, there exists a polynomial $f(\lambda)$ with integer coefficients such that $r - r^2 f(r) \in N$. An element r of R is called potent if there is an integer $n > 1$ such that $r^n = r$. We denote by I the set of potent elements of R . If R coincides with I , R is called a J -ring. As is well known, every J -ring is commutative (Jacobson's theorem). An ideal of R is called a J -ideal if it is a J -ring. Also, we denote by I_0 the set $\{r \in R \mid r \text{ generates a subring with identity}\}$. It is clear that $E \subseteq I \subseteq I_0$. Furthermore, if I_0 is a subring of R then I_0 coincides with I . In fact, if r is an arbitrary element of I_0 then there exists a polynomial $f(\lambda)$ with integer coefficients such that $r = r^2 f(r)$. This proves that I_0 is a reduced periodic ring, and therefore a J -ring. Especially, R is a J -ring if and only if $R = I_0$. If R is the direct sum of a J -ideal I' and a nil ideal N' , then it is easy to see that $I' = I = I_0$ and $N' = N$.

2. MAIN THEOREM.

Now, the main theorem of this note is stated as follows:

THEOREM 1. The following conditions are equivalent:

- 1) R is right (or left) π -regular and N is contained in some J -ideal A of R .
- 2) R is periodic and N is contained in some reduced ideal A of R .
- 3) R is a direct sum of a J -ring and a nil ring.

More precisely, if 1) or 2) is satisfied, then N is an ideal of R , $R = A \oplus N$, and $A = I_0$. In particular, if R is right (or left) s -unital, that is, $r \in rR$ (or $r \in Rr$) for all $r \in R$, then each of 1), 2) is equivalent to

- 4) R is a J -ring.

PROOF. Obviously, 3) \Rightarrow 2) \Rightarrow 1).

1) \Rightarrow 3). By a result of Dischinger (see, e.g., [3, Proposition 2]), R is strongly π -regular. Let r be an arbitrary element of R . Then there exists a positive integer n and elements s' , s'' of R such that $r^{2n}s' = s''r^{2n} = r^n$. We put $s = r^n s'^2$. As is easily seen,

$$s = s''r^n s' = s''r^{2n}$$

and

$$r^n s' r^n = s''r^{2n} = r^n = r^{2n}s' = r^n s'' r^n.$$

Hence,

$$r^n s = r^n s'' r^n s' = r^n s' = s''r^{2n}s' = s''r^n = s''r^n s' r^n = s r^n$$

and

$$r^{2n}s = r^n s r^n = r^n s' r^n = r^n.$$

Since $e = r^n s$ is an idempotent with $re = er$ ($\in A$) and $r^n e = r^n$, we see that

$$(r - re)^n = r^n(1 - e)^n = 0.$$

This together with $r = re + (r - re)$ proves that r is represented as a sum of an element in A and a nilpotent element. Now, let $a, b \in A$, and $x, y \in N$. Noting that $xa \cdot yb = xyba$ and $ax \cdot by = baxy$, we can easily see that $xa \in N \cap A = 0$ and $ax = 0$; $NA = AN = 0$. Set $xy = c + u$ and $x + y = d + v$ ($c, d \in A$ and $u, v \in N$), where we may assume that $u^\ell = v^\ell = 0$. In view of $NA = 0$, we obtain

$$(xy)^2 = xy(c + u) = xyu$$

and

$$(x + y)^2 = (x + y)(d + v) = (x + y)v,$$

and therefore

$$(xy)^{\ell+1} = xyu^\ell = 0$$

and

$$(x + y)^{\ell+1} = (x + y)v^\ell = 0.$$

We have thus seen that N forms an ideal of R and $R = A \oplus N$.

Given an integer $n > 1$, we denote by I_n the set $\{r \in R \mid r^n = r\}$. In [1], Abu-Khuzam and Yaqub proved that if R is a periodic ring with N commutative and for which I_n forms an ideal, then R is a subdirect sum of finite fields of at most n elements and a nil commutative ring. The next corollary includes this result.

COROLLARY 1. If R is periodic and I_n forms an ideal of R for some integer $n > 1$ then $R = I \oplus N$ and I is a subdirect sum of finite fields of at most n elements.

REFERENCES

1. ABU-KHUZAM, H. and YAQUB, A. Periodic rings with commuting nilpotents, Internat. J. Math. Math. Sci. 7 (1984) 403-406.
2. BELL, H.E. On commutativity of periodic rings and near rings, Acta Math. Acad. Sci. Hungar. 36 (1980), 293-302.
3. HIRANO, Y. Some studies on strongly π -regular rings, Math. J. Okayama Univ. 20 (1978), 141-149.

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br